



Cite this: *CrystEngComm*, 2016, 18, 2780

DOI: 10.1039/c6ce90050j

www.rsc.org/crystengcomm

Correction: Preferential site substitution of Eu^{3+} ions in $\text{Ca}_{10}(\text{PO}_4)_6\text{Cl}_2$ nanoparticles obtained using a microwave stimulated wet chemistry technique

Robert Pazik,^{*a} Jean-Marie Nedelec^{bc} and Rafal J. Wiglusz^{*a}

Correction for 'Preferential site substitution of Eu^{3+} ions in $\text{Ca}_{10}(\text{PO}_4)_6\text{Cl}_2$ nanoparticles obtained using a microwave stimulated wet chemistry technique' by Robert Pazik *et al.*, *CrystEngComm*, 2014, 16, 5308–5318.

"In accordance with the rule of $2J + 1$ at C_s symmetry a maximum of five sublevels should be present for the $^5\text{D}_0 \rightarrow ^7\text{F}_1$ and eight in the case of the $^5\text{D}_0 \rightarrow ^7\text{F}_2$ transitions whereas at C_3 symmetry the $^5\text{D}_0 \rightarrow ^7\text{F}_1$ splits into two and the $^5\text{D}_0 \rightarrow ^7\text{F}_2$ into three Stark components."

Should have read:

In accordance with the rule of $2J + 1$ at C_s symmetry a maximum of three sublevels should be present for the $^5\text{D}_0 \rightarrow ^7\text{F}_1$ and five in the case of the $^5\text{D}_0 \rightarrow ^7\text{F}_2$ transitions whereas at C_3 symmetry the $^5\text{D}_0 \rightarrow ^7\text{F}_1$ splits into two and the $^5\text{D}_0 \rightarrow ^7\text{F}_2$ into three Stark components.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Institute of Low Temperature and Structure Research, PAS, Okólna 2, 50–422 Wrocław, Poland. E-mail: R.Wiglusz@int.pan.wroc.pl; Fax: +48 71 344 10 29; Tel: +48 71 395 41 59

^b Clermont Université, ENSCCF, ICCF, BP10448, 63000 Clermont-Ferrand, France

^c CNRS, UMR 6296, ICCF, 63171 Aubière, France

