



Cite this: *Phys. Chem. Chem. Phys.*, 2016, **18**, 18614

DOI: 10.1039/c6cp90156e

www.rsc.org/pccp

Correction: Gd(III)–Gd(III) EPR distance measurements – the range of accessible distances and the impact of zero field splitting

Arina Dalaloyan,^a Mian Qi,^b Sharon Ruthstein,^c Shimon Vega,^a Adelheid Godt,^{*b} Akiva Feintuch^{*a} and Daniella Goldfarb^{*a}

Correction for 'Gd(III)–Gd(III) EPR distance measurements – the range of accessible distances and the impact of zero field splitting' by Arina Dalaloyan *et al.*, *Phys. Chem. Chem. Phys.*, 2015, **17**, 18464–18476.

The authors would like to make the readers aware that eqn (4) in the original version of the paper is incorrect. The correct version of eqn (4) is as follows:

$$H_{\text{ZFS}}^{(2)} = - \sum_{i=1}^2 \frac{(D^i)^2}{\omega_e} \left[\sin^2 \theta_{\text{ZFS}}^i \cos^2 \theta_{\text{ZFS}}^i \left[2S_z^i \left\{ \left((S_z^i)^2 + \frac{1}{4} \right) - S(S+1) \right\} \right] + \frac{1}{4} \sin^4 \theta_{\text{ZFS}}^i \left[S_z^i \left\{ \left((S_z^i)^2 + \frac{1}{2} \right) - S(S+1) \right\} \right] \right]$$

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Department of Chemical Physics, Weizmann Institute of Science, Rehovot, 76100, Israel. E-mail: Daniella.Goldfarb@weizmann.ac.il

^b Bielefeld University, Faculty of Chemistry and Center for Molecular Materials, D-33615 Bielefeld, Germany. E-mail: godt@uni-bielefeld.de

^c Department of Chemistry, Faculty of Exact Sciences, Bar-Ilan University, Ramat Gan, 5290002, Israel

