


 Cite this: *Phys. Chem. Chem. Phys.*,
2016, **18**, 19975

Correction: Determination of Kamlet–Taft parameters for selected solvate ionic liquids

 Daniel J. Eyckens,^{ab} Baris Demir,^a Tiffany R. Walsh,^a Tom Welton^c and
Luke C. Henderson^{*ab}

DOI: 10.1039/c6cp90177h

 Correction for 'Determination of Kamlet–Taft parameters for selected solvate ionic liquids' by Daniel J. Eyckens *et al.*, *Phys. Chem. Chem. Phys.*, 2016, **18**, 13153–13157.

www.rsc.org/pccp

The authors would like to make the following corrections to their article:

- (a) In the abstract and conclusion, β values for the **G3TFSI** and **G4TFSI** should be amended to $\beta = 0.31$ and $\beta = 0.28$, respectively.
 (b) In Table 1, the β values and footnote 'b' should be amended as shown below:

 Table 1 Polarity (E_T^N) and Kamlet–Taft parameters for **G3TFSI**, **G4TFSI**, **G3**, **G4**, and **[bmim]TFSI**

Solvent	E_T^N	α^a	β^b	π^*
G3TFSI	1.03	1.32	0.31	0.94
G4TFSI	1.03	1.35	0.28	0.90
G3 Glyme	0.30	0.01	0.71	0.65
G4 Glyme	0.28	0.05	0.72	0.67
[bmim][TFSI]	0.59	0.59	0.22	0.96
[bmim][TFSI]^c	0.55	0.61	0.24	0.98

^a Normalised against methanol. ^b Normalised against HMPA. ^c Previously published polarity and Kamlet–Taft values for this IL, ref. 26 and 31.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Institute for Frontier Materials, Deakin University, Waurn Ponds Campus, Geelong, Victoria, 3216, Australia. E-mail: luke.henderson@deakin.edu.au

^b Strategic Research Centre for Chemistry and Biotechnology, Deakin University, Waurn Ponds Campus, Geelong, Victoria, 3216, Australia

^c Department of Chemistry, Imperial College London, South Kensington, London, SW7 2AZ, UK
