CORRECTION

[View Article Online](#)
[View Journal](#) | [View Issue](#)

CrossMark
click for updates

Cite this: *Catal. Sci. Technol.*, 2016, **6**, 5754

Correction: A Pd–bisphosphine complex and organic functionalities immobilized on the same SiO₂ surface: detailed characterization and its use as an efficient catalyst for allylation

Ken Motokura,^a Koki Saitoh,^a Hiroto Noda,^a Wang-Jae Chun,^b Akimitsu Miyaji,^a Sho Yamaguchi^a and Toshihide Baba^{*a}

DOI: 10.1039/c6cy90069k

www.rsc.org/catalysis

Correction for 'A Pd–bisphosphine complex and organic functionalities immobilized on the same SiO₂ surface: detailed characterization and its use as an efficient catalyst for allylation' by Ken Motokura *et al.*, *Catal. Sci. Technol.*, 2016, DOI: 10.1039/c6cy00593d.

In the above article, the chemical structure of the nucleophile displayed in entry 6 of Table 5 is incorrect. A corrected version of Table 5 is given below.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Department of Chemical Science and Engineering, Tokyo Institute of Technology, 4259 Nagatsuta-cho, Midori-ku, Yokohama, 226-8502, Japan.

E-mail: tbaba@chemenv.titech.ac.jp

^b Graduate School of Arts and Sciences, International Christian University, Mitaka, Tokyo, 181-8585, Japan

Table 5 The allylation catalyzed by SiO₂/DABCO/PP-Pd under solvent-free conditions^a

Reaction scheme: Nucleophile + Allyl methyl carbonate (2.5 equiv.) $\xrightarrow[\text{neat}]{\text{SiO}_2/\text{DABCO/PP-Pd}, \text{K}_2\text{CO}_3}$ Mono-allylated product + Di-allylated product

Entry	Nucleophile	(mmol)	Pd (μmol)	K ₂ CO ₃ (mmol)	Time (h)	Yield (mono/di) (%)	TON ^b
1 ^c		12.6	0.21	0.025	216	3/87	106 000
2		9.0	1.5	0.13	31	<1/>99	12 000
3		9.0	1.5	0.13	5	<1/93	11 200
4		9.0	1.5	0.13	48	<1/98	11 800
5		9.0	1.5	0.13	54	83/12	6400
6		9.0	1.5	0.13	24	<1/98	11 700
7 ^d		9.0	1.5	0.13	31	<1/95	11 400
8		18.0	1.5	0.13	113	98/-	11 700
9		18.0	1.5	0.13	124	84/-	10 100

^a Reaction conditions: nucleophile, allyl methyl carbonate (2.5 equiv.), K₂CO₃, SiO₂/DABCO/PP-Pd, 70 °C, neat. Yield was determined by ¹H NMR analysis using an internal standard. ^b TON was calculated as follows: TON = (moles of nucleophile) × [(yield of the mono-product) + 2 × (yield of the di-product)]/(moles of Pd). ^c 80 °C. ^d 100 °C.