Dalton Transactions

An international journal of inorganic chemistry

www.rsc.org/dalton

ISSN 1477-9226

COMMUNICATION

W. Buchowicz *et al.*

Half-sandwich nickel complexes with ring-expanded NHC ligands – synthesis, structure and catalytic activity in Kumada–Tamao–Corriu coupling

175 YEARS

COMMUNICATION

View Article Online
View Journal | View Issue Cite this: *Dalton Trans.*, 2016, **45**, 8688Received 27th November 2015,
Accepted 26th January 2016

DOI: 10.1039/c5dt04663g

www.rsc.org/dalton

Half-sandwich nickel complexes with ring-expanded NHC ligands – synthesis, structure and catalytic activity in Kumada–Tamao–Corriu coupling†

Ł. Banach, P. A. Guńka and W. Buchowicz*

The general synthesis of $[\text{Ni}(\text{Cp})(\text{X})(\text{NHC})]$ complexes from a nickel halide, CpLi , and a carbene solution is reported. This procedure yields unprecedented complexes with ring-expanded NHC ligands (RE-NHC) of six- (1a, 1b), seven- (1c), and eight-membered (1d) heterocycles. The NMR spectra of 1a–1d are consistent with the hindered rotation of $\text{Ni}-\text{C}_{\text{carbene}}$ and $\text{N}-\text{C}_{\text{Mes}}$ bonds, while X-ray analyses of 1b, 1c, and 1d reveal a pronounced *trans* influence of the RE-NHC ligands. Complexes 1a–1e are efficient pre-catalysts in Kumada–Tamao–Corriu coupling with the maximum efficiency observed for complexes bearing the six-membered NHC.

Half-sandwich nickel complexes of the general formula $[\text{Ni}(\eta^5\text{-C}_5\text{H}_4\text{R})(\text{X})(\text{NHC})]$ ($\text{R} = \text{H}$, alkyl, alkenyl; $\text{X} = \text{Cl}$, Br , I ; $\text{NHC} = \text{N-heterocyclic carbene}$) are known only for five-membered NHCs. These complexes are usually synthesized by reactions of nickelocene with imidazolium halides (Scheme 1a).¹ However, for compounds with bulky substituents on the cyclopentadienyl (Cp) ligand, another protocol has to be utilized (Scheme 1b).²

Owing to their catalytic activity in many organic transformations, these air- and moisture-stable complexes are currently the object of intensive research.^{2–5} Efforts for tuning the properties of $[\text{Ni}(\text{Cp})(\text{X})(\text{NHC})]$ are focused on appropriate ligand selection within the scope of synthetic methodologies presented in Scheme 1, *i.e.* for five-membered NHCs only. Since the halide^{1a,3b–d,4} or Cp ligands⁵ can be readily displaced from the title complexes, NHC is the ligand of choice for modifications aimed at enhancing their catalytic activity. Recently, it has been shown that increasing the steric demand of NHCs by means of bulky substituents on the nitrogen atoms improved the efficiency of the arylation of aromatic halides catalysed by $[\text{Ni}(\text{Cp})(\text{X})(\text{NHC})]$.³ⁱ Another way to enhance spatial requirements as well as the electron-donating strength of NHC

Scheme 1 Reported synthetic routes to complexes $[\text{Ni}(\eta^5\text{-C}_5\text{H}_4\text{R}_{5-n})(\text{X})(\text{NHC})]$.

ligands is the expansion of the carbenic heterocycle.⁶ Both of these properties are presumed to influence the reactivity of complexes by facilitating the formation of a catalytically active intermediate and increasing its stability. On the other hand, ring-expanded NHC (RE-NHC) may enable an intramolecular C–H activation, which can be detrimental to catalytic performance.⁷

Although RE-NHC nickel complexes are known, these studies are limited to only a few papers.^{7,8} To the best of our knowledge, to date there have been no reports on $[\text{Ni}(\text{Cp})(\text{X})(\text{RE-NHC})]$. Driven by scientific curiosity to see if $[\text{Ni}(\text{Cp})(\text{X})(\text{RE-NHC})]$ complexes are of a greater catalytic activity than their congeners with five-membered “classical” NHCs, we decided to develop the synthesis of such complexes.

Our initial attempts to prepare the desired RE-NHC nickel complexes according to the procedure shown in Scheme 1a, *i.e.* from nickelocene and an appropriate carbene precursor, failed. Reactions involving the use of $[\text{Ni}(\text{acac})_2]$ as the starting material were also unsuccessful (Scheme 1b). A probable

Faculty of Chemistry, Warsaw University of Technology, Noakowskiego 3, 00-664 Warsaw, Poland. E-mail: wbuch@ch.pw.edu.pl

† Electronic supplementary information (ESI) available. CCDC 1429974–1429976. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5dt04663g

Scheme 2 Synthesis of complexes $[\text{Ni}(\text{Cp})(\text{X})(\text{NHC})]$.

reason for this lack of reactivity is the weak acidity of carbene precursors⁹ and steric congestion around the reacting centre.⁶ Inspired by the synthesis of related complexes with Cp*-functionalized NHC ligands,^{3g} we decided to test a similar approach involving the addition of the two ligands (Cp and RE-NHC) to a nickel halide. Despite the plausible formation of different products, *e.g.* nickelocene and $[\text{Ni}(\text{X})_2(\text{RE-NHC})_2]$, the desired $[\text{Ni}(\text{Cp})(\text{X})(\text{RE-NHC})]$ complexes were obtained in remarkably good yields as the only isolable products. More specifically, $[\text{Ni}(\text{DME})(\text{X})_2]$ (X = Br or Cl, DME = 1,2-dimethoxyethane) was treated first with a CpLi solution in THF to form what we believe is a dinuclear complex $[\text{Ni}(\text{Cp})(\mu\text{-X})_2]$.¹⁰ Subsequently, a freshly prepared RE-NHC solution, obtained by deprotonation of the carbene precursor with BuLi, was injected. After stirring overnight and conventional purification complexes **1a–1d** were obtained in 32–41% yields (two steps from the Ni halide) as deep purple solids (Scheme 2).¹¹

Moreover, to further expand the scope of our method, we have applied it to prepare a complex with a five-membered NHC. Thus, $[\text{Ni}(\text{Cp})(\text{Br})(\text{SIMes})]^{3c}$ (SIMes = 1,3-bis(2,4,6-trimethylphenyl)-4,5-dihydroimidazol-2-ylidene) (**1e**) was obtained *via* this simple procedure in 61% yield (two steps from $[\text{Ni}(\text{DME})\text{Br}_2]$).

The NMR spectra of complexes **1a–1d** were conventionally recorded in CDCl_3 at ambient temperature. Signals originating from NHC ligands imply restricted Ni–C_{carbene} and N–C_{Mes} bond rotation. Owing to the inequivalence of hydrogen atoms in each NCH₂ group their signals appear in the ¹H NMR spectra as unresolved multiplets. Moreover, as a result of the hindered rotation, all complexes **1a–1d** have a characteristic set of signals for mesityl wingtips, showing two separated equal singlets (2:2 ratio) for *meta* hydrogens from phenyl rings and three singlets (6:6:6 ratio) for methyl groups, *i.e.* *ortho* methyls and *meta* hydrogens are nonequivalent within each mesityl moiety. Another interesting feature of ¹H NMR spectra concerns the Cp group. While for $[\text{Ni}(\text{Cp})(\text{X})(\text{NHC})]$ with five-membered NHCs the Cp singlet appears from 5.2 to 4.6 ppm, its chemical shift for **1a–1d** is unusual and strongly influenced by the size of the NHC as it varies from 4.21 ppm for **1b** to 3.68 ppm for **1d**. Moreover, all signals in the ¹H NMR spectrum of **1d** are strongly broadened. We tentatively inter-

Fig. 1 Molecular structure of complex **1b**. Since there are no significant differences in the conformations of the four molecules present in the asymmetric unit only one is shown. Thermal ellipsoids are drawn at the 50% probability level. Selected bond lengths in Å: Ni1–Br1 2.3371(5), Ni1–C1 2.054(3), Ni1–C2 2.193(3), Ni1–C3 2.197(3), Ni1–C4 2.165(3), Ni1–C5 2.151(3), Ni1–C6 1.905(3), C1–C2 1.429(5), C1–C5 1.433(5), C2–C3 1.379(4), C3–C4 1.440(4), C4–C5 1.387(5).

pret these phenomena in terms of spin equilibrium,^{4f,12} *i.e.* increasing content of the high-spin form of the complex which is correlated to the NHC's ring size.

Crystal structure analysis of complexes **1b–1d** revealed that they crystallize in the monoclinic crystal system and their molecular geometries are typical for nickel(II) NHC complexes (Fig. 1–3).¹³ The Ni–C_{Cp} bond lengths exhibit a pronounced *trans* influence on the RE-NHC ligands while the C–C bond lengths in Cp (see figure captions for selected bond lengths) show variations characteristic of the so-called “diene distortion”.^{14,15}

Complex $[\text{Ni}(\text{Cp})(\text{Cl})(\text{IMes})]$ (IMes = 1,3-bis(2,4,6-trimethylphenyl)imidazol-2-ylidene) was reported as an active catalyst in the Kumada–Tamao–Corriu (KTC) coupling between *p*-tolyl-

Fig. 2 Molecular structure of complex **1c**. Thermal ellipsoids are drawn at the 50% probability level. Selected bond lengths in Å: Ni1–Br1 2.3281(3), Ni1–C1 2.0529(17), Ni1–C2 2.1914(17), Ni1–C3 2.1847(17), Ni1–C4 2.1722(18), Ni1–C5 2.1422(17), Ni1–C6 1.8972(15), C1–C2 1.431(3), C1–C5 1.436(3), C2–C3 1.384(3), C3–C4 1.444(2), C4–C5 1.389(3).

Fig. 3 Molecular structure of complex **1d**. Thermal ellipsoids are drawn at the 50% probability level. Selected bond lengths in Å: Ni1–Br1 2.3466(4), Ni1–C1 2.074(2), Ni1–C2 2.195(2), Ni1–C3 2.178(2), Ni1–C4 2.150(2), Ni1–C5 2.161(2), Ni1–C6 1.918(2), C1–C2 1.438(3), C1–C5 1.425(4), C2–C3 1.380(3), C3–C4 1.439(3), C4–C5 1.393(4).

magnesium bromide and phenyl *O*-sulfamate.^{3a} To our surprise, further applications of [Ni(Cp)(X)(NHC)] complexes in this reaction are limited.^{31,p,16} In order to get an insight into the influence of NHC ring expansion on the activity of [Ni(Cp)(X)(NHC)] in KTC coupling, we tested complexes **1a–1e** in reactions of PhMgBr with halotoluenes (3-bromotoluene, 4-chlorotoluene). The expansion of the NHC ligand from a five- to six-membered ring proved to be beneficial for the catalytic activity (Table 1, entries 1–3, 9 and 10). A catalyst loading of only 1 mol% of **1a** or **1b** is sufficient to achieve conversions over 95% with very good selectivity at room temperature in 90 minutes.¹⁷ Also, the nature of the halogen ligand in the

Table 1 Kumada–Tamao–Corriu coupling of PhMgBr with halotoluenes catalysed by **1a–1e**^a

Entry	Catalyst	<i>n</i>	X'	Conversion ^b (%)	Ratio A : B : C ^b
1	1e	0	3-Br	78	91 : 6 : 3
2	1a	1	3-Br	97	92 : 6 : 2
3	1b	1	3-Br	98	92 : 7 : 1
4	1c	2	3-Br	36	89 : 11 : 0
5 ^c	1c	2	3-Br	86	85 : 10 : 5
6	1d	3	3-Br	Traces	—
7 ^c	1d	3	3-Br	97	70 : 20 : 10
8	1a	1	4-Cl	Traces	—
9 ^c	1a	1	4-Cl	68	65 : 22 : 13
10 ^c	1e	0	4-Cl	42	54 : 36 : 10

^a All reactions in duplicate; reaction conditions: THF (2.8 mL), halotoluene (0.66 mmol), phenylmagnesium bromide solution in THF (0.81 mL, 1.00 mmol, 1.24 mol L^{−1}), nickel complex (6.7 μmol, 1 mol%), room temperature, 90 min; see the ESI for details.

^b Determined with GC. ^c Reaction time = 21 h.

nickel complex appeared to have no influence on the catalytic performance (Table 1, entries 2 and 3). Further expansion of the NHC ligand to seven- or eight-membered rings resulted in diminished activity. However, high conversions could be achieved at longer reaction times (Table 1, entries 4–7). Selectivity is another parameter affected by ring expansion beyond the six-membered system (Table 1, entries 4, 5 and 7).

Conclusions

We report here a new general synthesis of [Ni(Cp)(X)(NHC)] complexes that is applicable to ring-expanded NHC ligands. Complexes with a six-membered NHC provided the best efficiency in KTC coupling. Further studies will focus on the application of [Ni(Cp)(X)(RE-NHC)] in other catalytic reactions, including arylamination of aromatic halides.

Acknowledgements

The authors thank the Warsaw University of Technology for financial support.

Notes and references

- (a) C. D. Abernethy, A. H. Cowley and R. A. Jones, *J. Organomet. Chem.*, 2000, **596**, 3; (b) R. A. Kelly III, N. M. Scott,

- S. Díez-González, E. D. Stevens and S. P. Nolan, *Organometallics*, 2005, **24**, 3442.
- 2 (a) V. Ritleng, C. Barth, E. Brenner, S. Milosevic and M. J. Chetcuti, *Organometallics*, 2008, **27**, 4223; (b) W. Buchowicz, J. Conder, D. Hryciuk and J. Zachara, *J. Mol. Catal. A: Chem.*, 2014, **381**, 16.
 - 3 For catalytic applications of $[\text{Ni}(\text{Cp})(\text{X})(\text{NHC})]$, see: (a) T. K. Macklin and V. Snieckus, *Org. Lett.*, 2005, **7**, 2519; (b) D. A. Malyshev, N. M. Scott, N. Marion, E. D. Stevens, V. P. Ananikov, I. P. Beletskaya and S. P. Nolan, *Organometallics*, 2006, **25**, 4462; (c) W. Buchowicz, A. Koziół, L. B. Jerzykiewicz, T. Lis, S. Pasynkiewicz, A. Pęcherzewska and A. Pietrzykowski, *J. Mol. Catal. A: Chem.*, 2006, **257**, 118; (d) W. Buchowicz, W. Wojtczak, A. Pietrzykowski, A. Lupa, L. B. Jerzykiewicz, A. Makal and K. Woźniak, *Eur. J. Inorg. Chem.*, 2010, 648; (e) V. Ritleng, A. M. Oertel and M. J. Chetcuti, *Dalton Trans.*, 2010, **39**, 8153; (f) A. M. Oertel, V. Ritleng and M. J. Chetcuti, *Organometallics*, 2012, **31**, 2829; (g) L. Postigo and B. Royo, *Adv. Synth. Catal.*, 2012, **354**, 2613; (h) L. P. Bheeter, M. Henrion, L. Brelot, C. Darcel, M. J. Chetcuti, J.-B. Sortais and V. Ritleng, *Adv. Synth. Catal.*, 2012, **354**, 2619; (i) A. R. Martin, Y. Makida, S. Meiries, A. M. Z. Slawin and S. P. Nolan, *Organometallics*, 2013, **32**, 6265; (j) O. R. Luca, D. L. Huang, M. K. Takase and R. H. Crabtree, *New J. Chem.*, 2013, **37**, 3402; (k) L. P. Bheeter, M. Henrion, M. J. Chetcuti, C. Darcel, V. Ritleng and J.-B. Sortais, *Catal. Sci. Technol.*, 2013, **3**, 3111; (l) S. Tamba, K. Fujii, H. Meguro, S. Okamoto, T. Tendo, R. Komobuchi, A. Sugie, T. Nishino and A. Mori, *Chem. Lett.*, 2013, **42**, 281; (m) M. Henrion, M. J. Chetcuti and V. Ritleng, *Chem. Commun.*, 2014, **50**, 4624; (n) A. Włodarska, A. Koziół, M. Dranka, J. Jurkowski and A. Pietrzykowski, *J. Mol. Catal. A: Chem.*, 2014, **395**, 481; (o) Y. Wei, A. Petronilho, H. Mueller-Bunz and M. Albrecht, *Organometallics*, 2014, **33**, 5834; (p) P. Buchalski, R. Pacholski, K. Chodkiewicz, W. Buchowicz, K. Suwińska and A. Shkurenko, *Dalton Trans.*, 2015, **44**, 7169; (q) H. Valdés, M. Poyatos, G. Ujaque and E. Peris, *Chem. – Eur. J.*, 2015, **21**, 1578.
 - 4 For other examples of studies involving $[\text{Ni}(\text{Cp})(\text{X})(\text{NHC})]$, see: (a) E. F. Hahn, B. Heidrich, A. Hepp and T. Pape, *J. Organomet. Chem.*, 2007, **692**, 4630; (b) C. Radloff, F. E. Hahn, T. Pape and R. Fröhlich, *Dalton Trans.*, 2009, 7215; (c) A. M. Oertel, J. Freudenreich, J. Gein, V. Ritleng, L. F. Veiros and M. J. Chetcuti, *Organometallics*, 2011, **30**, 3400; (d) A. M. Oertel, V. Ritleng, L. Burr and M. J. Chetcuti, *Organometallics*, 2011, **30**, 6685; (e) A. M. Oertel, V. Ritleng, A. Busiah, L. F. Veiros and M. J. Chetcuti, *Organometallics*, 2011, **30**, 6495; (f) W. Buchowicz, L. Banach, J. Conder, P. A. Guńka, D. Kubicki and P. Buchalski, *Dalton Trans.*, 2014, **43**, 5847; (g) S. Pelties, D. Herrmann, B. de Bruin, F. Hartl and R. Wolf, *Chem. Commun.*, 2014, **50**, 7014.
 - 5 M. Henrion, A. M. Oertel, V. Ritleng and M. J. Chetcuti, *Chem. Commun.*, 2013, **49**, 6424.
 - 6 For selected examples, see: (a) P. Bazinet, G. P. A. Yap and D. S. Richeson, *J. Am. Chem. Soc.*, 2003, **125**, 13314; (b) C. C. Scarborough, B. V. Popp, I. A. Guzei and S. S. Stahl, *J. Organomet. Chem.*, 2005, **690**, 6143; (c) M. Iglesias, D. J. Beetstra, A. Stasch, P. N. Horton, M. B. Hursthouse, S. J. Coles, K. J. Cavell, A. Dervisi and I. A. Fallis, *Organometallics*, 2007, **26**, 4800; (d) M. Iglesias, D. J. Beetstra, J. C. Knight, L.-L. Ooi, A. Stasch, S. Coles, L. Male, M. B. Hursthouse, K. J. Cavell, A. Dervisi and I. A. Fallis, *Organometallics*, 2008, **27**, 3279; (e) M. Iglesias, D. J. Beetstra, B. Kariuki, K. J. Cavell, A. Dervisi and I. A. Fallis, *Eur. J. Inorg. Chem.*, 2009, 1913; (f) J. J. Dunsford and K. J. Cavell, *Dalton Trans.*, 2011, **40**, 9131; (g) J. J. Dunsford, K. J. Cavell and B. M. Kariuki, *Organometallics*, 2012, **31**, 4118; (h) E. L. Kolychev, A. F. Asachenko, P. B. Dzhevakov, A. A. Bush, V. V. Shuntikov, V. N. Khrustalev and M. S. Nechaev, *Dalton Trans.*, 2013, **42**, 6859; (i) O. S. Morozov, A. V. Lunchev, A. A. Bush, A. A. Tukov, A. F. Asachenko, V. N. Khrustalev, S. S. Zaleskiy, V. P. Ananikov and M. S. Nechaev, *Chem. – Eur. J.*, 2014, **20**, 6162; (j) J. J. Dunsford and K. J. Cavell, *Organometallics*, 2014, **33**, 2902.
 - 7 C. J. E. Davies, M. J. Page, C. E. Ellul, M. F. Mahon and M. K. Whittlesey, *Chem. Commun.*, 2010, **46**, 5151.
 - 8 (a) P. D. Newman, K. J. Cavell and B. M. Kariuki, *Organometallics*, 2010, **29**, 2724; (b) R. C. Poulten, M. J. Page, A. G. Algarra, J. J. LeRoy, I. López, E. Carter, A. Llobet, S. A. Macgregor, M. F. Mahon, D. M. Murphy, M. Murugesu and M. K. Whittlesey, *J. Am. Chem. Soc.*, 2013, **135**, 13640; (c) M. J. Page, W. Y. Lu, R. C. Poulten, E. Carter, A. G. Algarra, B. M. Kariuki, S. A. Macgregor, M. F. Mahon, K. J. Cavell, D. M. Murphy and M. K. Whittlesey, *Chem. – Eur. J.*, 2013, **19**, 2158; (d) R. C. Poulten, I. López, A. Llobet, M. F. Mahon and M. K. Whittlesey, *Inorg. Chem.*, 2014, **53**, 7160.
 - 9 E. M. Higgins, J. A. Sherwood, A. G. Lindsay, J. Armstrong, R. S. Massey, R. W. Alder and A. M. C. O'Donoghue, *Chem. Commun.*, 2011, **47**, 1559.
 - 10 Complexes of this type have been reported for multi-substituted Cp ligands, see: (a) U. Kölle, B. Fuss, F. Khouzami and J. Gersdorf, *J. Organomet. Chem.*, 1985, **290**, 77; (b) M. Schär, D. Saurenz, F. Zimmer, I. Schädlich, G. Wolmershäuser, S. Demeshko, F. Meyer, H. Sitzmann, O. M. Heigl and F. H. Köhler, *Organometallics*, 2013, **32**, 6298.
 - 11 See the ESI† for details.
 - 12 M. E. Smith and R. A. Andersen, *J. Am. Chem. Soc.*, 1996, **118**, 11119.
 - 13 The ESI† and CCDC 1429974–1429976 contain the supplementary crystallographic data for this paper.
 - 14 P. L. Holland, M. E. Smith, R. A. Andersen and R. G. Bergman, *J. Am. Chem. Soc.*, 1997, **119**, 12815.
 - 15 L. Banach, P. A. Guńka, D. Górka, M. Podlowska, J. Zachara and W. Buchowicz, *Eur. J. Inorg. Chem.*, 2015, 5677.
 - 16 A. E. Goetz and N. K. Garg, *Nat. Chem.*, 2013, **5**, 54.

17 We note that the activity of $[\text{Ni}(\text{Cp})(\text{X})(\text{RE-NHC})]$ in KTC coupling is comparable to that reported for the corresponding Pd complexes with standard five-membered

NHCs, see: Z. Jin, X.-P. Gu, L.-L. Qiu, G.-P. Wu, H.-B. Song and J.-X. Fang, *J. Organomet. Chem.*, 2011, **696**, 859.