



Cite this: *New J. Chem.*, 2016, 40, 6564

Received (in Montpellier, France)
27th January 2016,
Accepted 6th June 2016

DOI: 10.1039/c6nj00281a

www.rsc.org/njc

Visible light-induced monofluoromethylation of benzofurans and benzothiophenes with ethyl bromofluoroacetate was developed. This method provided a convenient access to novel α -fluoro- α -heteroarylesters under mild reaction conditions.

Since Fried and Sabo's discovery that the incorporation of a fluorine atom into a corticosteroid derivative led to valuable enhanced biological activity in 1954,¹ fluorinated organic compounds have attracted increasing attention for use in pharmaceuticals and agrochemicals.² The α -fluoro- α -arylcarboxylic acid moiety is found in many active bioactive compounds³ and has been widely used as a building block in organic synthesis.⁴ Consequently, lots of efforts have been made in the preparation of α -aryl- α -fluorocarbonyl units. There are mainly two strategies for the construction of α -aryl- α -fluorocarbonyl units. The first strategy involves the direct fluorination of α -arylcarbonyl derivatives using nucleophilic⁵ or electrophilic⁶ fluorinating reagents. The second strategy involves the condensation reaction of aromatic substrates and fluorine-containing building blocks.^{7,8} Compared to the direct fluorination strategy, the "building block" strategy has been much less explored despite its potential importance. Recently, Sanford^{7a} as well as Wang and Zhou,^{7b} respectively, reported the S_NAr reaction of *ortho*-fluoronitrobenzenes and 2-fluoromalones for the preparation of α -aryl- α -fluoroesters (Scheme 1a). Palladium-^{8a,b} and nickel-catalyzed^{8c} cross-coupling reactions of aromatic substrates and fluorine-containing building blocks have provided an efficient strategy to incorporate monofluoromethylene units into aromatic compounds (Scheme 1b). However, both of the methods require the use of prefunctionalized aromatic substrates. The direct monofluoromethylation of aryl C–H bonds remains a big challenge.


^a Key Laboratory of Organofluorine Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Science, Shanghai 200032, China. E-mail: flq@mail.sioc.ac.cn

^b College of Chemistry, Chemical Engineering and Biotechnology, Donghua University, Shanghai 201620, China

† Electronic supplementary information (ESI) available: Detailed experimental procedures and analytical data for all new compounds. See DOI: 10.1039/c6nj00281a

Visible light-induced monofluoromethylation of heteroarenes with ethyl bromofluoroacetate†

Wei Yu,^a Xiu-Hua Xu^a and Feng-Ling Qing*^{ab}




Scheme 1 Monofluoromethylation of (hetero)aromatic compounds.

Recently, visible light photoredox catalysis has emerged as an efficient and eco-friendly tool in organic synthesis⁹ and has been applied in the fluoroalkylation of organic compounds.¹⁰ In 2013, we reported a mild and versatile approach for the direct introduction of an ethoxycarbonyldifluoromethyl group into heteroarenes with ethyl bromodifluoroacetate *via* visible light photocatalysis.¹¹ On the basis of that work, we anticipated that a similar transformation with ethyl bromofluoroacetate might allow for the direct introduction of the ethoxycarbonylmonofluoromethyl group into heteroarenes. In continuation of our research interest in visible light-induced fluoroalkylation reactions,^{10f,i,w,11} herein we describe the reactions of benzofurans and benzothiophenes with ethyl bromofluoroacetate under visible light photoredox conditions (Scheme 1c). This reaction provides a convenient approach to the synthesis of novel α -fluoro- α -heteroarylesters.

Initially, we investigated the model reaction of benzofuran **1a** with ethyl bromofluoroacetate using blue LEDs as the source of visible light (Table 1). Among the commonly used photocatalysts, only *fac*-Ir(ppy)₃ could promote the reaction giving the desired product **2a** in 55% yield, and no reaction occurred in the presence of Ru(bpy)₃Cl₂·6H₂O, methylene blue, or eosin




Table 1 Optimization of reaction conditions^a


Entry	Photocat.	Base	Solvent	Yield ^b (%)
1	Ru(bpy) ₃ Cl ₂ ·6H ₂ O	K ₂ HPO ₄	MeCN	0
2	<i>fac</i> -Ir(ppy) ₃	K ₂ HPO ₄	MeCN	55
3	Methylene Blue	K ₂ HPO ₄	MeCN	0
4	Eosin Y	K ₂ HPO ₄	MeCN	0
5	[Ir(ppy) ₂ Cl] ₂	K ₂ HPO ₄	MeCN	29
6	[Ir(dtbbpy)(ppy) ₂][PF ₆]	K ₂ HPO ₄	MeCN	12
7	<i>fac</i> -Ir(ppy) ₃	K ₂ CO ₃	MeCN	43
8	<i>fac</i> -Ir(ppy) ₃	K ₃ PO ₄	MeCN	52
9	<i>fac</i> -Ir(ppy) ₃	KOAc	MeCN	23
10	<i>fac</i> -Ir(ppy) ₃	<i>t</i> -BuONa	MeCN	12
11	<i>fac</i> -Ir(ppy) ₃	DBU	MeCN	21
12	<i>fac</i> -Ir(ppy) ₃	DIPEA	MeCN	13
13	<i>fac</i> -Ir(ppy) ₃	K ₂ HPO ₄	DMF	54
14	<i>fac</i> -Ir(ppy) ₃	K ₂ HPO ₄	DCM	46
15	<i>fac</i> -Ir(ppy) ₃	K ₂ HPO ₄	THF	49
16	<i>fac</i> -Ir(ppy) ₃	K ₂ HPO ₄	DMF/H ₂ O	74

^a Reaction conditions: **1a** (0.1 mmol), ethyl bromofluoroacetate (0.2 mmol), photocat. (0.005 mmol), base (0.2 mmol), solvent (1.0 mL), visible light, rt, under N₂, 12 h. ^b Yields determined by ¹⁹F NMR spectroscopy using fluorobenzene as an internal standard.

Y (entries 1–4). Other iridium photocatalysts including [Ir(ppy)₂Cl]₂ and [Ir(dtbbpy)(ppy)₂][PF₆] were found to be less effective than *fac*-Ir(ppy)₃ (entries 5 and 6). To increase the yield, various inorganic and organic bases including K₂CO₃, K₃PO₄, KOAc, *t*-BuONa, DBU, and DIPEA were tested (entries 7–12). However, none of them gave higher yield. Further screening of different solvents disclosed that DMF, DCM, and THF gave the desired product in slightly lower yields than MeCN (entries 13–15). To our delight, when the reaction was performed in DMF/H₂O, the yield of **2a** was improved to 74% (entry 16). It is noteworthy that this optimized yield is much higher than the one obtained in the coupling reaction of benzofuran-2-ylboronic acid with ethyl bromofluoroacetate.^{8b}

With the optimized reaction conditions in hand, we next examined the substrate scope of the visible light-promoted direct monofluoromethylation of heteroarenes (Table 2). A variety of benzofurans **1a–s** were employed to give the corresponding 2-monofluoromethylated products **2a–s** in moderate to excellent yields. The selective formation of 2-substituted isomers is probably due to the generation of a more stable benzylic radical as an intermediate. Different functional groups including ethers, esters, and nitriles were well tolerated. Notably, substrates bearing fluoro, chloro, and bromo substituents on the arene rings are also compatible, thus providing opportunities for additional transformations (**1m–1q**). With substrates (**1r** and **1s**) containing electron-withdrawing groups, the desired products were isolated in low yields. It is noteworthy that the reaction could be extended to benzothiophenes **1t** and **1u**, furnishing products **2t** and **2u** in moderate yields, respectively. The bromofluoromethylation of indole **1v** gave the desired product **2v** in 47% yield. However, the electron-rich arene **1w** displayed poor reactivity, with only 22% of the product being obtained. In the cases of furan and thiophene, low conversion was observed.

Table 2 Substrate scope of photocatalytic monofluoromethylation of heteroarenes^a



2a , 62%	2b , 76%	2c , 77%
2d , 60%	2e , 68%	2f , 63%
2g , 68%	2h , 50%	2i , 77%
2j , 70%	2k , 50%	2l , 55%
2m , 48%	2n , 70%	2o , 74%
2p , 51%	2q , 36%	2r , 40%
2s , 31%	2t , 34%	2u , 51%
2v , 47%	2w , 22%	

^a Reaction conditions: **1** (0.5 mmol), ethyl bromofluoroacetate (1.0 mmol), *fac*-Ir(ppy)₃ (0.025 mmol), K₂HPO₄ (1.0 mmol), DMF/H₂O (5.0 mL), visible light, rt, under N₂, 12 h, isolated yields.

Finally, the control experiments were carried out to gain insights into the reaction mechanism. No reaction took place in the absence of visible light or *fac*-Ir(ppy)₃, and the addition of 2,2,6,6-tetramethylpiperidine-1-oxyl (TEMPO), a known radical scavenger, could totally inhibit the reaction. Based on the above results and the reported work,^{10a,12} a plausible reaction mechanism was proposed as in Scheme 2. Initially, the irradiation of visible light excited *fac*-Ir^{III}(ppy)₃ to *fac*-Ir^{III}(ppy)₃*. Then, a single-electron-transfer (SET) from *fac*-Ir^{III}(ppy)₃* to ethyl bromofluoroacetate generated the radical intermediate **A**, which was subsequently added to benzofuran **1** for the formation of radical intermediate **B**. Intermediate **B** was then oxidized to cation intermediate **C**, which underwent deprotonation affording the final product **2**.

In conclusion, we have developed a visible light-promoted monofluoromethylation of heteroarenes with ethyl bromofluoroacetate. This protocol does not require prefunctionalization





Scheme 2 Proposed reaction mechanism.

of the heteroaremetics and proceeds under mild reaction conditions, which makes it attractive in the preparation α -fluoro- α -heteroarylcarbonyl compounds. The investigation of the present method in the synthesis of biologically active compounds is in progress.

Experimental section

General procedure for the photocatalytic monofluoromethylation of heteroarenes

A 50 mL Schlenk flask equipped with a rubber septum and a magnetic stir bar was charged with *fac*-Ir(ppy)₃ (16.4 mg, 0.025 mmol, 5 mol%), K₂HPO₄ (174.1 mg, 1.0 mmol, 2.0 equiv.), ethyl bromodifluoroacetate (184.9 mg, 1.0 mmol, 2.0 equiv.), and heteroarene (0.5 mmol, 1.0 equiv.). DMF (1.0 mL) and H₂O (4.0 mL) were added to the mixture. Then, the reaction mixture was degassed three times by the freeze–pump–thaw procedure. The flask was placed at a distance of 2 cm from the blue LEDs. The mixture was stirred under a nitrogen atmosphere and irradiated by blue LEDs for 12 h. After the reaction was complete, the reaction mixture was extracted by Et₂O, and the combined organic phase was dried over anhydrous Na₂SO₄. The solvent was removed under vacuum and the residue was purified by column chromatography on silica gel to give the corresponding product.

Acknowledgements

We thank the National Natural Science Foundation of China (21272036, 21332010 and 21421002) and the National Basic Research Program of China (2012CB21600) for financial support.

Notes and references

- 1 J. Fried and E. F. Sabo, *J. Am. Chem. Soc.*, 1954, **76**, 1455.
- 2 (a) K. Muller, C. Faeh and F. Diederich, *Science*, 2007, **317**, 1881; (b) S. Purser, P. R. Moore, S. Swallow and V. Gouverneur, *Chem. Soc. Rev.*, 2008, **37**, 320; (c) W. K. Hagmann, *J. Med. Chem.*, 2008, **51**, 4359; (d) N. A. Meanwell, *J. Med. Chem.*, 2011, **54**, 2529; (e) E. P. Gillis, K. J. Eastman, M. D. Hill, D. J. Donnelly and N. A. Meanwell, *J. Med. Chem.*, 2015, **58**, 8315.
- 3 *Fluorine in Medicinal Chemistry and Chemical Biology*, ed. I. Ojima, Wiley, Chichester, 2009.
- 4 For recent examples, see: (a) A. J. Grenning and J. A. Tunge, *Org. Lett.*, 2010, **12**, 740; (b) L. Wu, L. Falivene, E. Drinkel, S. Grant, A. Linden, L. Cavallo and R. Dorta, *Angew. Chem., Int. Ed.*, 2012, **51**, 2870; (c) S. Mizuta, I. S. R. Stenhagen, M. O'Duill, J. Wolstenhulme, A. K. Kirjavainen, S. J. Forsback, M. Tredwell, G. Sandford, P. R. Moore, M. Huiban, S. K. Luthra, J. Passchier, O. Solin and V. Gouverneur, *Org. Lett.*, 2013, **15**, 2648; (d) K. Araki, T. Katagiri and M. Inoue, *J. Fluorine Chem.*, 2014, **157**, 41; (e) J. Alvarado, A. T. Herrmann and A. Zakarian, *J. Org. Chem.*, 2014, **79**, 6206; (f) D. W. Manley and J. C. Walton, *Org. Lett.*, 2014, **16**, 5394.
- 5 (a) S. Hamman, M. Barrelle, F. Tetaz and C. G. Beguin, *J. Fluorine Chem.*, 1987, **37**, 85; (b) E. Laurent, B. Marquet, R. Tardivel and H. Thiebault, *Tetrahedron Lett.*, 1987, **28**, 2359; (c) S. Watanabe, T. Fujita, M. Sakamoto, H. Endo and T. Kitazume, *J. Fluorine Chem.*, 1990, **47**, 187; (d) C.-S. Tsai, Y.-K. Li and L.-C. Lo, *Org. Lett.*, 2002, **4**, 3607; (e) V. Dinoiu, T. Fukuhara, K. Miura and N. Yoneda, *J. Fluorine Chem.*, 2003, **121**, 227; (f) J. Yin, D. S. Zarkowsky, D. W. Thomas, M. M. Zhao and M. A. Huffman, *Org. Lett.*, 2004, **6**, 1465; (g) T.-H. Shie, Y.-L. Chiang, J.-J. Lin, Y.-K. Li and L.-C. Lo, *Carbohydr. Res.*, 2006, **341**, 443; (h) K.-Y. Kim, B. C. Kim, H. B. Lee and H. Shin, *J. Org. Chem.*, 2008, **73**, 8106; (i) X. Zhao, W. Zhuang, D. Fang, X. Xue and J. Zhou, *Synlett*, 2009, 779; (j) I. E. Kopka, L. S. Lin, J. P. Jewell, T. J. Lanza, T. M. Fong, C.-P. Shen, Z. J. Lao, S. Ha, L. G. Castonguay, L. V. der Ploeg, M. T. Goulet and W. K. Hagmann, *Bioorg. Med. Chem. Lett.*, 2010, **20**, 4757; (k) T. Sawamura, S. Kuribayashi, S. Inagi and T. Fuchigami, *Org. Lett.*, 2010, **12**, 644.
- 6 (a) W. J. Middleton and E. M. Bingham, *J. Am. Chem. Soc.*, 1980, **102**, 4845; (b) T. Umemoto, K. Kawada and K. Tomita, *Tetrahedron Lett.*, 1986, **27**, 4465; (c) S. T. Purrington and D. L. Woodard, *J. Org. Chem.*, 1990, **55**, 3423; (d) F. A. Davis, W. Han and C. K. Murphy, *J. Org. Chem.*, 1995, **60**, 4730; (e) S. Rozen, A. Hagooley and R. Harduf, *J. Org. Chem.*, 2001, **66**, 7464; (f) H. Fujisawa and Y. Takeuchi, *J. Fluorine Chem.*, 2002, **117**, 173; (g) F. Zhang and J. Z. Song, *Tetrahedron Lett.*, 2006, **47**, 7641; (h) T. Suzuki, Y. Hamashima and M. Sodeoka, *Angew. Chem., Int. Ed.*, 2007, **46**, 5435; (i) D. H. Paull, M. T. Scerba, E. Alden-Danforth, L. R. Widger and T. Lectka, *J. Am. Chem. Soc.*, 2008, **130**, 17260; (j) I. T. Schiefer, S. Abdul-Hay, H. Wang, M. Vanni, Z. Qin and G. R. J. Thatcher, *J. Med. Chem.*, 2011, **54**, 2293; (k) F. Li, Z. Wu and J. Wang, *Angew. Chem., Int. Ed.*, 2015, **54**, 656.
- 7 (a) A. Harsanyi, G. Sandford, D. S. Yufit and J. A. K. Howard, *Beilstein J. Org. Chem.*, 2014, **10**, 1213; (b) F. Zhu, P.-W. Xu, F. Zhou, C.-H. Wang and J. Zhou, *Org. Lett.*, 2015, **17**, 972.
- 8 (a) N. A. Beare and J. F. Hartwig, *J. Org. Chem.*, 2002, **67**, 541; (b) C. Guo, X. Yue and F.-L. Qing, *Synthesis*, 2010, 1837;



- (c) Y.-M. Su, G.-S. Feng, Z.-Y. Wang, Q. Lan and X.-S. Wang, *Angew. Chem., Int. Ed.*, 2015, **54**, 6003.
- 9 (a) T. P. Yoon, M. A. Ischay and J. Du, *Nat. Chem.*, 2010, **2**, 527; (b) J. M. R. Narayanam and C. R. J. Stephenson, *Chem. Soc. Rev.*, 2011, **40**, 102; (c) J. Xuan and W.-J. Xiao, *Angew. Chem., Int. Ed.*, 2012, **51**, 6828; (d) C. K. Prier, D. A. Rankic and D. W. C. MacMillan, *Chem. Rev.*, 2013, **113**, 5322; (e) D. A. Nicewicz and T. M. Nguyen, *ACS Catal.*, 2014, **4**, 355.
- 10 For recent examples, see: (a) J. Jung, E. Kim, Y. You and E. J. Cho, *Adv. Synth. Catal.*, 2014, **356**, 2741; (b) N. Iqbal, J. Jung, S. Park and E. J. Cho, *Angew. Chem., Int. Ed.*, 2014, **53**, 539; (c) P. Xu, A. Abdulkader, K. Hu, Y. Cheng and C. Zhu, *Chem. Commun.*, 2014, **50**, 2308; (d) W. Li, X. Zhu, H. Mao, Z. Tang, Y. Cheng and C. Zhu, *Chem. Commun.*, 2014, **50**, 7521; (e) M. Rueda-Becerril, O. Mahé, M. Drouin, M. B. Majewski, J. G. West, M. O. Wolf, G. M. Sammis and J.-F. Paquin, *J. Am. Chem. Soc.*, 2014, **136**, 2637; (f) Q.-Y. Lin, X.-H. Xu and F.-L. Qing, *J. Org. Chem.*, 2014, **79**, 10434; (g) Y. Yasu, Y. Arai, R. Tomita, T. Koike and M. Akita, *Org. Lett.*, 2014, **16**, 780; (h) D. Cantillo, O. de Frutos, J. A. Rincón, C. Mateos and C. O. Kappe, *Org. Lett.*, 2014, **16**, 896; (i) A. Carboni, G. Dagousset, E. Magnier and G. Masson, *Org. Lett.*, 2014, **16**, 1240; (j) S. H. Oh, Y. R. Malpani, N. Ha, Y.-S. Jung and S. B. Han, *Org. Lett.*, 2014, **16**, 1310; (k) J. Xie, X. Yuan, A. Abdulkader, C. Zhu and J. Ma, *Org. Lett.*, 2014, **16**, 1768; (l) X. Sun and S. Yu, *Org. Lett.*, 2014, **16**, 2938; (m) Y.-M. Su, Y. Hou, F. Yin, Y.-M. Xu, Y. Li, X. Zheng and X.-S. Wang, *Org. Lett.*, 2014, **16**, 2958; (n) L. Wang, X.-J. Wei, W.-L. Jia, J.-J. Zhong, L.-Z. Wu and Q. Liu, *Org. Lett.*, 2014, **16**, 5842; (o) X.-J. Tang and W. R. Dolbier, Jr., *Angew. Chem., Int. Ed.*, 2015, **54**, 4246; (p) R. Tomita, T. Koike and M. Akita, *Angew. Chem., Int. Ed.*, 2015, **54**, 12923; (q) X.-J. Tang, Z. Zhang and W. R. Dolbier, *Chem. – Eur. J.*, 2015, **21**, 18961; (r) A. Singh, J. J. Kubik and J. D. Weaver, *Chem. Sci.*, 2015, **6**, 7206; (s) J. W. Beatty, J. J. Douglas, K. P. Cole and C. R. J. Stephenson, *Nat. Commun.*, 2015, **6**, 7979; (t) Q.-Y. Lin, X.-H. Xu and F.-L. Qing, *Org. Biomol. Chem.*, 2015, **13**, 8740; (u) Z. Zhang, X. Tang, C. S. Thomason and W. R. Dolbier, Jr., *Org. Lett.*, 2015, **17**, 3528; (v) Z. Zhang, X. Tang and W. R. Dolbier, Jr., *Org. Lett.*, 2015, **17**, 4401; (w) Q.-Y. Lin, X.-H. Xu, K. Zhang and F.-L. Qing, *Angew. Chem., Int. Ed.*, 2016, **55**, 1479; (x) Y. Arai, R. Tomita, G. Ando, T. Koike and M. Akita, *Chem. – Eur. J.*, 2016, **22**, 1262.
- 11 Q.-Y. Lin, L. Chu and F.-L. Qing, *Chin. J. Chem.*, 2013, **31**, 885.
- 12 J. D. Nguyen, J. W. Tucker, M. D. Konieczynska and C. R. J. Stephenson, *J. Am. Chem. Soc.*, 2011, **133**, 4160.

