



Cite this: *Nanoscale*, 2016, **8**, 16787

## Correction: Composition-dependent Raman modes of $\text{Mo}_{1-x}\text{W}_x\text{S}_2$ monolayer alloys

Yanfeng Chen,<sup>a</sup> Dumitru O. Dumcenco,<sup>b</sup> Yiming Zhu,<sup>a</sup> Xin Zhang,<sup>c</sup> Nannan Mao,<sup>a,d</sup> Qingliang Feng,<sup>a,d</sup> Mei Zhang,<sup>a</sup> Jin Zhang,<sup>d</sup> Ping-Heng Tan,<sup>c</sup> Ying-Sheng Huang<sup>\*b</sup> and Liming Xie<sup>\*a</sup>

DOI: 10.1039/c6nr90195f

[www.rsc.org/nanoscale](http://www.rsc.org/nanoscale)

Correction for 'Composition-dependent Raman modes of  $\text{Mo}_{1-x}\text{W}_x\text{S}_2$  monolayer alloys' by Yanfeng Chen *et al.*, *Nanoscale*, 2014, **6**, 2833–2839.

The authors would like to bring a corrected Table 1 to the attention of the readers:

**Table 1** MREI-model parameters for  $\text{Mo}_{1-x}\text{W}_x\text{S}_2$  monolayer and bulk alloys (note  $F$  is in  $10^6 \text{ amu cm}^{-2}$ )

Fitted parameters	$A'_1$ (1 L) or $A_{1g}$ (bulk)			$E'(1 \text{ L})$ or $E_{2g}^1$ (bulk)				
	$F_{10}$	$F_{20}$	$\theta_1$	$F_{30}$	$F_{40}$	$F_{50}$	$\theta_2$	$\theta_3$
Monolayer	5.162	5.371	−0.039	2.840	4.803	12.08	0.375	−0.052
Bulk	5.297	4.916	−0.147	2.812	4.801	12.06	0.387	−0.064

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup>CAS Key Laboratory of Standardization and Measurement for Nanotechnology, National Center for Nanoscience and Technology, Beijing 100190, P. R. China. E-mail: xielm@nanocr.cn; Tel: +86-010 8254 5692

<sup>b</sup>Department of Electronic Engineering, National Taiwan University of Science and Technology, Taipei 106, Taiwan, Republic of China. E-mail: ysh@mail.ntust.edu.tw

<sup>c</sup>State Key Laboratory of Superlattices and Microstructures, Institute of Semiconductors, Chinese Academy of Sciences, Beijing 100083, China

<sup>d</sup>Center for Nanochemistry, College of Chemistry and Molecular Engineering, Peking University, Beijing 100871, China

