Organic & Biomolecular Chemistry

PAPER

Cite this: Org. Biomol. Chem., 2016, **14**, 7956

Aluminium ion-promoted radical-scavenging reaction of methylated hydroquinone derivatives

Ikuo Nakanishi,*^a Kei Ohkubo,*^{a,b,c} Yukihiro Ogawa,^{a,d} Ken-ichiro Matsumoto,^{a,d} Toshihiko Ozawa^e and Shunichi Fukuzumi^{c,f}

The effect of the aluminium ion (Al^{3+}) on the scavenging reaction of a 2,2-diphenyl-1-picrylhydrazyl radical (DPPH⁺), as a reactivity model of reactive oxygen species, with hydroquinone (QH_2) and its methylated derivatives (Me_nQH₂, n = 1-4) was investigated using stopped-flow and electrochemical techniques in a hydroalcoholic medium. The second-order rate constants (*k*) for the DPPH⁺-scavenging reaction of the hydroquinones increased with the increasing number of methyl substituents. Upon addition of Al³⁺, the *k* values significantly increased depending on the concentration of Al³⁺. Such an accelerating effect of Al³⁺ on the DPPH⁺-scavenging rates of the hydroquinones results from the remarkable positive shift of the one-electron reduction potential (E_{red}) of DPPH⁺ in the presence of Al³⁺. These results demonstrate that Al³⁺, a strong Lewis acid, can act as a radical-scavenging promoter by stabilising the one-electron reduced species of the radical, although Al³⁺ is reported not only to act as a pro-oxidant but also to strongly interact with biomolecules, showing toxicities.

Received 11th July 2016, Accepted 29th July 2016 DOI: 10.1039/c6ob01470d

www.rsc.org/obc

Introduction

Metal ions are known to play important roles in controlling biological redox reactions.¹ It is also well known that metal ions accelerate electron-transfer reactions by binding to electron acceptors.² Furthermore, they are reported to form transient complexes with the superoxide anion $(O_2^{\bullet-})$, which is one of the reactive oxygen species (ROS).^{3,4} Among them, a complex of Al³⁺ with $O_2^{\bullet-}$ contributes to promote the Fenton reaction by reducing Fe³⁺ to Fe²⁺ to produce hydroxyl radicals ('OH), the strongest ROS.⁵ In addition, Al³⁺ as a strong Lewis acid forms a variety of stable complexes with proteins and biomolecules, such as citrate, glutamate and α -ketoglutarate.⁶

^aQuantitative RedOx Sensing Team (QRST), Department of Basic Medical Sciences for Radiation Damages, National Institute of Radiological Sciences (NIRS), National Institutes for Quantum and Radiological Science and Technology (QST), Inage-ku, Chiba 263-8555, Japan. E-mail: nakanishi.ikuo@qst.go.jp; Fax: +81-43-255-6819; Tel: +81-43-206-3131

^cDepartment of Chemistry and Nano Science, Ewha Womans University,

Al³⁺ also interacts with molecules containing phosphate groups, such as AMP, ATP, RNA and DNA.⁷ These complexes disrupt the cell metabolic processes, leading to neurological diseases.⁸

Ubiquinol (UQH₂), the reduced form of coenzyme Q_{10} , has a hydroquinone (QH₂) structure as an active centre, showing an efficient scavenging activity against ROS.9 In human beings, ROS are generated as side products during the energy production by mitochondrial electron-transport systems or the elimination of virus and bacteria in macrophages and neutrophils. However, excessive ROS cause many diseases, such as cancer, diabetes, cardiovascular disease, Alzheimer's and Parkinson's disease by oxidising proteins, lipids and DNA.^{10,11} Thus, antioxidants are very useful to protect against oxidative stress and may lead to prevention of these diseases. UQH₂ is one of the lipid-soluble antioxidants and efficiently scavenges lipid peroxyl radicals.^{12,13} Furthermore, UQH₂ reacts with the α -tocopheroxyl radical to regenerate α -tocopherol inside the biomembrane.^{14–16} On the other hand, QH_2 is a major benzene metabolite and is widely applied in human and industrial activities. QH₂ is used as an antioxidant in the rubber and food industries due to its antioxidative activity.¹⁷ QH₂ also has a long history of use in pigmentary disorders and is applied in the commercially available cosmetic skin lightning formulations.18-20

The radical-scavenging reactions of phenolic antioxidants, such as flavonoids and α -tocopherol, are affected by redox-inactive metal ions, such as Mg²⁺, Al³⁺ and Sc³⁺.²¹⁻²³ However, little is known about the effects of metal ions on the radical-scavenging reaction of QH₂ and its derivatives.

View Article Online

^bDivision of Applied Chemistry, Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan

Seoul 120-750, Korea

^dGraduate School of Advanced Integration Science, Chiba University, Inage-ku, Chiba 263-8522, Japan

^eDivision of Oxidative Stress Research, Showa Pharmaceutical University, Machida, Tokyo 194-8543, Japan

^fFaculty of Science and Technology, Meijo University, SENTAN, Japan Science and Technology Agency (JST), Nagoya, Aichi 468-8502, Japan



Fig. 1 DPPH as a model of ROS.



Fig. 2 Structures of UQH₂, QH₂ and its methylated derivatives.

Particularly, whether Al³⁺ promotes the radical-scavenging reactions of antioxidants or acts as a pro-oxidant is yet to be clarified.

We report herein the effects of Al³⁺ on the scavenging reaction of a 2,2-diphenyl-1-picrylhydrazyl radical (DPPH') (Fig. 1) with QH₂ and its methylated derivatives, which are methylhydroquinone (MeQH₂), 2,6-dimethylhydroquinone (Me₂QH₂), trimethylhydroquinone (Me₃QH₂) and tetramethylhydroquinone (Me₄QH₂) (Fig. 2).²⁴ DPPH' is frequently used as a reactivity model of peroxyl radicals, which are relatively less reactive ROS compared to hydroxyl and alkoxyl radicals.²⁵⁻³⁰ In addition, we investigated the effects of the number of methyl substituents of the hydroquinones on the DPPH'-scavenging activity. Furthermore, the electron-donating ability of the hydroquinones and the electron-accepting ability of DPPH' were also evaluated by cyclic voltammetry (CV) and secondharmonic alternating current voltammetry (SHACV) in the absence and presence of Al3+.31-35 The strong Lewis acidity of Al³⁺ enhanced the electron-accepting ability of DPPH[•], leading to an acceleration of the DPPH'-scavenging reaction by the hydroquinones.

Experimental

Materials

 QH_2 and its methylated derivatives were purchased from Tokyo Chemical Industry Co., Ltd, Japan. DPPH[•] was purchased from

Wako Pure Chemical industries, Ltd. Tetra-*n*-butylammonium perchlorate (Bu_4NClO_4), used as a supporting electrolyte for the electrochemical measurements, was purchased from Tokyo Chemical Industry Co., Ltd, Japan. Bu_4NClO_4 was recrystallised from ethanol (EtOH) and dried under vacuum at 313 K. Aluminium perchlorate nonahydrate [$Al(ClO_4)_3 \cdot 9H_2O$] was purchased from Wako Pure Chemical Ind. Ltd, Japan. EtOH (spectral grade) used as a solvent was purchased from Nacalai Tesque, Inc., Japan.

Spectral and kinetic measurements

To avoid the effect of molecular oxygen (O_2) , the reactions were carried out under strictly deaerated conditions. A continuous flow of argon gas was bubbled through an EtOH-H₂O (9:1 v/v) solution (3.0 mL) containing DPPH[•] (5.0–8.9 \times 10⁻⁵ M) or the hydroquinones $(1.1-6.6 \times 10^{-3} \text{ M})$ in a reservoir of the spectrophotometer for 7 min. The rates of the DPPH'-scavenging reactions of the hydroquinones were determined by monitoring the absorbance change at 517 nm due to DPPH' using a stoppedflow technique on a UNISOKU RSP-1000-02NM spectrophotometer. The pseudo-first-order rate constants (k_{obs}) were determined by least-squares curve fit using an Apple MacBook Pro personal computer. The first-order plots of $\ln(A - A_{\infty})$ vs. time (A and A_{∞} denoted as the absorbance at the reaction time and the final absorbance, respectively) were linear until three or more half-lives with the correlation coefficient $\rho > 0.999$. The reaction of hydroquinones with DPPH' in the presence of various concentrations of $Al(ClO_4)_3$ was carried out in the same manner.

Electrochemical measurements

The cyclic voltammetry (CV) and second-harmonic alternating current voltammetry (SHACV)³⁶ measurements were performed on an ALS-630A electrochemical analyser in deaerated EtOH- H_2O (9:1 v/v) containing 0.1 M Bu₄NClO₄ as a supporting electrolyte. The glassy carbon-working electrode (BAS) was polished with BAS polishing alumina suspension and rinsed with methanol before use. The counter electrode was a platinum wire. The measured potentials were recorded with respect to an Ag/AgNO₃ (0.01 M) reference electrode. One-electron oxidation potentials (E_{rox}) of hydroquinones and one-electron reduction potentials (E_{red}) of DPPH[•] (vs. Ag/AgNO₃) were converted to those vs. SCE by adding 0.29 V.³⁷ All electrochemical measurements were carried out at room temperature under 1 atm Ar.

pH measurements

The pH values of the solutions of hydroquinones in the presence or absence of the metal ion in EtOH–H₂O (9:1 v/v) were measured by using a METTLER TOLEDO SevenEasy pH meter.

Results and discussion

Kinetics of the DPPH'-scavenging reaction of the hydroquinones in the presence or absence of Al³⁺

The rate of the DPPH-scavenging reaction of the hydroquinones was measured by monitoring the absorbance change



Fig. 3 Plots of k_{obs} vs. $[Me_nQH_2]$ (n = 0-4) for the reaction of QH₂ (closed circles), MeQH₂ (open circles), Me₂QH₂ (closed squares), Me₃QH₂ (open squares) and Me₄QH₄ (closed triangles) with DPPH⁻ in deaerated EtOH-H₂O (9 : 1 v/v) at 298 K.

Table 1Second-order rate constants (k) for the DPPH⁻-scavengingreaction of the hydroquinones in the presence or absence of $Al(ClO_4)_3$ in deaerated $EtOH-H_2O$ (9:1 v/v) at 298 K

	$k/M^{-1} s^{-1}$						
[Al(ClO ₄) ₃]/M	QH_2	MeQH ₂	Me ₂ QH ₂	Me ₃ QH ₂	Me ₄ QH ₂		
0	1.3×10^{2}	2.0×10^{2}	3.4×10^2	1.1×10^{3}	1.4×10^{3}		
0.05	3.5×10^{2}	7.0×10^{2}	1.0×10^{3}	2.7×10^{3}	3.0×10^{3}		
0.1	3.9×10^{2}	9.3×10^{2}	1.7×10^{3}	3.4×10^{3}	3.5×10^{3}		
0.2	3.4×10^{2}	$1.1 imes 10^3$	2.8×10^{3}	5.3×10^{3}	5.4×10^3		
0.3	$5.8 imes 10^2$	1.3×10^{3}	3.2×10^{3}	6.3×10^{3}	$5.8 imes 10^3$		
0.5	5.2×10^2	1.1×10^{3}	4.4×10^{3}	7.5×10^{3}	8.2×10^{3}		

at 517 nm due to DPPH' using a stopped-flow technique in deaerated EtOH-H₂O (9:1 v/v) at 298 K. The decay of the absorbance at 517 nm obeyed pseudo-first-order kinetics, when the concentration of the hydroquinones ($[Me_nQH_2]$) (n = 0-4) was maintained at a more than 10-fold excess of the DPPH' concentration. The pseudo-first-order rate constants (k_{obs}) increase with increasing [Me_nQH₂], exhibiting the first order dependence on $[Me_nQH_2]$ (Fig. 3). From the slope of the linear plots of k_{obs} vs. [Me_nQH₂], the second-order rate constants (k) were determined for the DPPH⁻-scavenging reaction (Table 1). The k values of the hydroquinones increased with the increasing number of methyl substituents in the absence of metal ions. Upon addition of $Al(ClO_4)_3$ (0.05–0.50 M) to the hydroquinone-DPPH' system, the k values for all the hydroquinones significantly increased with the increasing $Al(ClO_4)_3$ concentration (Fig. 4, Table 1).

Effect of Al³⁺ on the redox behaviour of the hydroquinones and DPPH'

The one-electron oxidation potentials (E_{ox}) of the hydroquinones were determined by cyclic voltammetry (CV) and



Fig. 4 Plots of *k vs.* $[Al(ClO_4)_3]$ for the reaction of QH₂ (closed circles), MeQH₂ (open circles), Me₂QH₂ (closed squares), Me₃QH₂ (open squares) and Me₄QH₄ (closed triangles) with DPPH⁻ in deaerated EtOH-H₂O (9 : 1 v/v) at 298 K.

second-harmonic alternating current voltammetry (SHACV)³⁶ measurements in the presence or absence of Al³⁺ in EtOH– H₂O (9:1 v/v) to investigate the detailed effects of Al³⁺ on the DPPH⁻-scavenging reaction of the hydroquinones. The $E_{\rm ox}$ values of the hydroquinones were determined from the intersection of SHACV waves (Fig. 5). In the absence of the metal ion, the $E_{\rm ox}$ values of the hydroquinones were shifted to the negative direction with the increasing number of methyl substituents (Table 2). The electron-donating methyl substituents contribute to an increase in the electron density, leading to an enhancement of the electron-donating ability of the hydroquinones.

Upon addition of Al(ClO₄)₃ (0.05–0.5 M), the E_{ox} values of hydroquinones were significantly shifted to the positive direction



Fig. 5 Cyclic voltammogram (CV) (dashed line) and second-harmonic alternating current voltammogram (SHACV) (solid line) of QH₂ (2.4 × 10^{-3} M) recorded at the scan rate of 0.1 V s⁻¹ and 4 mV s⁻¹, respectively, on a glassy carbon working electrode in EtOH–H₂O (9:1 v/v) (0.1 M Bu₄NClO₄).

[Al(ClO ₄) ₃]/M	$E_{\rm ox}$ (vs. SCE	$E_{\rm red}$ (vs. SCE)/V				
	QH ₂	$MeQH_2$	Me ₂ QH ₂	Me ₃ QH ₂	Me ₄ QH ₂	DPPH.
0	+0.39	+0.36	+0.29	+0.23	+0.19	+0.23
0.05	+0.48	+0.42	+0.36	+0.29	+0.24	$+0.67^{a}$
0.1	+0.54	+0.47	+0.39	+0.33	+0.28	$+0.69^{a}$
0.2	+0.55	+0.48	+0.40	+0.34	+0.28	$+0.70^{a}$
0.3	+0.56	+0.50	+0.42	+0.36	+0.30	$+0.70^{a}$
0.5	+0.56	+0.50	+0.42	+0.36	+0.32	$+0.69^{a}$

Table 2 The E_{ox} values of hydroquinones and the E_{red} values of DPPH in EtOH-H₂O (9:1 v/v) in the presence or absence of Al(ClO₄)₃

^{*a*} Determined by SHACV at the rate of 4 mV s⁻¹ on a glassy carbon working electrode (0.1 M Bu₄NClO₄).

depending on the concentration of Al³⁺ (Table 2). Thus, the one-electron donating ability of the hydroquinones was decreased in the presence of Al^{3+} . The pK_a values of the hydroquinones in H₂O and DMSO are listed in Table 3.^{38,39} The pH values of the hydroquinone solutions in the presence or absence of Al(ClO₄)₃ in EtOH-H₂O (9:1 v/v) are shown in Table 4. The pK_a values of the hydroquinones used in this study in aqueous solution are in the range of 9.9-11.3 (Table 3). Thus, all the hydroquinones do not undergo deprotonation and exist in their neutral forms. The one-electron reduction potential (E_{red}) of DPPH[•] was also determined using CV in the presence or absence of Al^{3+} . In the absence of Al^{3+} , two well-defined reversible redox waves were observed as shown in Fig. 6 (dashed line), from which the one-electron reduction potential of DPPH' was determined to be +0.23 V vs. SCE. In the presence of 0.1 M $Al(ClO_4)_3$, however, the cyclic voltammogram of DPPH' was remarkably changed and the E_{red} value of DPPH' was significantly shifted to the positive direction (Fig. 6, Table 2). These results indicated that the DPPH anion (DPPH⁻), the one-electron reduced species of DPPH⁺, strongly interacted with Al³⁺ to enhance the electron-accepting ability of DPPH'. The effects of metal ions on the Ered of DPPH' differed by the Lewis acidity.40

Table 3 T	he p K_a	values	of hydr	oquinones
-----------	------------	--------	---------	-----------

	QH_2	MeQH ₂	Me_2QH_2	Me ₃ QH ₂	Me ₄ QH ₂
p <i>K</i> _a	9.9^{a} 19.1^{b}	10.1^{a} 18.3 ^b	-19.0^{b}	-17.8 ^b	11.3^{a} 20.0^{b}

^{*a*} The pK_a values in aqueous solution.³⁸ ^{*b*} The pK_a values in DMSO.³⁹

Table 4 The pH values of hydroquinone solutions in $EtOH-H_2O$ (9:1 v/v) containing 0.1 M Bu_4NClO_4 in the presence or absence of $Al(ClO_4)$

$[Al(ClO_4)_3]/M$	$\rm QH_2$	$MeQH_2$	Me_2QH_2	$\mathrm{Me_3QH_2}$	Me_4QH_2
0	5.5	4.7	4.7	4.5	4.6
0.05	2.1	2.2	2.1	2.1	2.1
0.1	1.7	1.8	1.6	1.7	1.7
0.2	1.5	1.5	2.0	1.6	1.5
0.3	1.3	1.1	1.2	1.1	1.1
0.5	0.9	0.7	0.8	0.8	0.8



Fig. 6 Cyclic voltammograms of DPPH[•] $(2.0 \times 10^{-3} \text{ M})$ in the absence (dashed line) and presence (solid line) of 0.1 M Al(ClO₄)₃ recorded at the scan rate 0.1 V s⁻¹ on a glassy carbon working electrode in EtOH-H₂O (9 : 1 v/v) (0.1 M Bu₄NClO₄).

We previously reported that the radical-scavenging mechanism of a vitamin E model is significantly affected by solvents.^{22b} In a protic solvent, such as methanol, an electron transfer from the vitamin E model to radicals followed by proton transfer is favorable compared to one-step hydrogenatom transfer. Based on these results, the DPPH^{*}-scavenging reaction by hydroquinones in the hydroalcoholic medium would proceed *via* the electron transfer. The free energy change (ΔG_{et}) of the electron transfer from hydroquinones to DPPH^{*} is calculated by using eqn (1),

$$\Delta G_{\rm et} = F(E_{\rm ox} - E_{\rm red}) \tag{1}$$

where *F* is the Faraday constant, which is negative ($\Delta G_{\rm et} < 0$) in the presence of Al³⁺. Al³⁺ enhanced the electron transfer by strongly stabilizing DPPH⁻ and increased the *k* values of the hydroquinones depending on its concentration, in spite of decreasing the one-electron donor ability of hydroquinones.

Fig. 7 shows the free-energy-change dependence of $\log k$ for the radical-scavenging reaction of DPPH^{*} with a series of hydroquinones in the absence and presence of Al³⁺ (0.5 M).



Fig. 7 Plots of log *k vs.* ΔG_{et} for the DPPH⁻-scavenging reaction by the hydroquinone derivatives in the absence of metal ions (closed circles) and the presence of Al³⁺ (0.5 M) (open circles). The lines represent the curve fits to eqn (2) with $\lambda = 1.78$ and 2.19 eV for the reaction without and with Al³⁺ (0.5 M), respectively.

The dependence of outer-sphere electron transfer has been well established by Marcus as given by eqn (2),⁴¹

$$1/k = 1/k_{\rm diff} + 1/\{Z \exp[-(\lambda/4)(1 + \Delta G_{\rm et}/\lambda)^2/k_{\rm B}T]\}$$
(2)

in which k_{diff} is the diffusion rate constant $(10^{10} \text{ M}^{-1} \text{ s}^{-1})$,^{42,43} λ is the reorganization energy of electron transfer, k_{B} is the Boltzmann constant and Z is the collision frequency $(1 \times 10^{11} \text{ M}^{-1} \text{ s}^{-1})$. The λ value is determined to be 1.78 eV for the radical-scavenging reaction in the absence of metal ions. In the presence of Al³⁺, the rate constants increase with a decrease in the ΔG_{et} value as shown in Fig. 7. The λ value in the absence of metal ions, suggesting that trivalent Al³⁺ strongly binds to DPPH⁻ as an electron transfer product in electron transfer from the hydroquinones to DPPH⁺.^{2,44}

Conclusions

The DPPH'-scavenging activity of the hydroquinones increased with the increasing number of methyl substituents in the dearated hydroalcoholic medium. Upon addition of $Al(ClO_4)_3$ to the hydroquinone–DPPH' system, the *k* values of the hydroquinones were significantly increased depending on the concentration of Al^{3+} , in spite of decreasing the electron-donating ability of the hydroquinones. Al^{3+} , one of the strong Lewis acids, strongly stabilized the DPPH⁻ generated by the electron transfer from the hydroquinones to DPPH'. In addition, the electron transfer occurred voluntarily ($\Delta G_{et} < 0$) in the presence of Al^{3+} according to the electrochemical data. The strong Lewis acids, such as Al^{3+} , significantly accelerated the electron-transfer reaction and enhanced the radical-scavenging reaction of phenolic antioxidants. The results obtained in this study demonstrate that Al^{3+} , a strong Lewis acid, can act as a radicalscavenging promoter by stabilising the one-electron reduced species of the radical, although Al³⁺ is reported not only to act as a pro-oxidant but also to strongly interact with bio-molecules, showing toxicities.

Acknowledgements

This work was partially supported by Grant-in-Aid (No. 26460056 to I. N., 26620154, 266288037 and 16K13964 to K. O.) from the Ministry of Education, Culture, Sports, Science and Technology, Japan, Acenet Inc. Foundation (to K. O.) and SENTAN projects from JST (to S. F.). We thank Mr T. Waki (Chiba University) for his help in the measurements of redox potentials.

Notes and references

- 1 W. Kaim, B. Schwederski and A. Klein, *Bioinorganic Chemistry: Inorganic Elements in the Chemistry of Life*, John Wiley & Son, Chichester, 2nd edn, 2013.
- 2 (a) S. Fukuzumi and K. Ohkubo, *Coord. Chem. Rev.*, 2010,
 254, 372; (b) S. Fukuzumi, K. Ohkubo and Y. Morimoto,
 Phys. Chem. Chem. Phys., 2012, 14, 8472.
- 3 (a) S. Fukuzumi and K. Ohkubo, *Chem. Eur. J.*, 2000, 6, 4532; (b) K. Ohkubo, S. C. Menon, A. Orita, J. Otera and S. Fukuzumi, *J. Org. Chem.*, 2003, 68, 4720–4726.
- 4 J. I. Mujika, F. Ruipérez, I. Infante, J. M. Ugalde, C. Exley and X. Lopez, *J. Phys. Chem. A*, 2011, **115**, 6717.
- 5 F. Ruipérez, J. I. Mujika, J. M. Ugalde, C. Exley and Z. Lopez, *J. Inorg. Biochem.*, 2012, **117**, 118.
- 6 (a) J. I. Mujika, J. M. Ugalde and X. Lopez, *J. Phys. Chem. B*, 2014, **118**, 6680; (b) J. I. Mujika, J. M. Ugalde and X. Lopes, *Phys. Chem. Chem. Phys.*, 2012, **14**, 12465.
- 7 N. B. Luque, J. I. Mujika, E. Rezabal, J. M. Ugalde and X. Lopez, *Phys. Chem. Chem. Phys.*, 2014, 16, 20107.
- 8 C. A. Shaw, S. Seneff, S. D. Kette, L. Tomljenovic, J. W. Oller Jr. and R. M. Davidson, *J. Toxicol.*, 2014, 491316.
- 9 (a) A. Maroz, R. F. Anderson, R. A. Smith and M. P. Murphy, *Free Radical Biol. Med.*, 2009, 465, 105;
 (b) A. M. James, R. A. Smith and M. P. Murphy, *Arch. Biochem. Biophys.*, 2004, 423, 47;
 (c) T. Kawashima, K. Ohkubo and S. Fukuzumi, *Phys. Chem. Chem. Phys.*, 2011, 13, 3344.
- 10 I. F. Benzie, Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol., 2003, 136, 113.
- 11 K. Rahman, Clin. Interventions Aging, 2007, 2, 219.
- 12 (a) P. Forsmark-Andree, G. Dallner and L. Ernster, Free Radicals Biol. Med., 1995, 19, 749; (b) L. Ernster and G. Dallner, Biochim. Biophys. Acta, 1995, 1271, 195.
- 13 B. Frei, M. C. Kim and B. N. Ames, Proc. Natl. Acad. Scl. U. S. A., 1990, 87, 4879.
- 14 (a) K. Mukai, A. Ouchi, S. Nakaya and S. Nagaoka, J. Phys. Chem. B, 2013, 117, 8378; (b) A. Ouchi, S. Nagaoka and

K. Mukai, J. Phys. Chem. B, 2010, **114**, 6601; (c) S. Itoh, S. Nakaya and K. Mukai, J. Phys. Chem. A., 2008, **112**, 448.

- 15 (a) Y. Yamamoto, J. Dermatol. Sci., 2001, 27, S1;
 (b) Y. Yamamoto, E. Komuro and E. Niki, J. Nutr. Sci. Vitaminol., 1990, 36, 505.
- 16 K. U. Ingold, V. W. Bowry, R. Stocker and C. Walling, Proc. Natl. Acad. Sci. U. S. A., 1993, 90, 45.
- 17 IPCS, *Hydroquinone*, Environmental Health Criteria 157, 1994.
- 18 J. Ancerewicz, E. Migliavacca, P. A. Carrupt, B. Testa, F. Brée, R. Zini, J. P. Tillement, S. Labidalle, D. Guyot, A. M. Chauvet-Monges, A. Crevat and A. L. Ridant, *Free Radical Biol. Med.*, 1998, 25, 113.
- 19 M. Nishizawa, M. Kohono, M. Nishimura, A. Kitagawa and Y. Niwano, *Chem. Pharm. Bull.*, 2005, **53**, 714.
- 20 T. W. Tse, J. Dermatol. Treat., 2010, 21, 272.
- 21 T. Waki, S. Kobayashi, K. Matsumoto, T. Ozawa, T. Kamada and I. Nakanishi, *Chem. Commun.*, 2013, **49**, 9842.
- 22 (a) I. Nakanishi, T. Shimada, K. Ohkubo, S. Manda, T. Shimizu, S. Urano, H. Okuda, N. Miyata, T. Ozawa, K. Anzai, S. Fukuzumi, N. Ikota and K. Fukuhara, Chem. Lett., 2007, 36, 1276; (b) I. Nakanishi, T. Kawashima, K. Ohkubo, H. Kanazawa, K. Inami, M. Mochizuki, K. Fukuhara, H. Okuda, T. Ozawa, S. Itoh, S. Fukuzumi and N. Ikota, Org. Biomol. Chem., 2005, 3, 626; (c) I. Nakanishi, K. Ohkubo, K. Miyazaki, W. Hakamata, S. Urano, T. Ozawa, H. Okuda, S. Fukuzumi, N. Ikota and K. Fukuhara, Chem. Res. Toxicol., 2004, 17, 26; (d) I. Nakanishi, Y. Uto, K. Ohkubo, K. Miyazaki, H. Yakumaru, S. Urano, H. Okuda, J. Ueda, T. Ozawa, K. Fukuhara, S. Fukuzumi, H. Nagasawa, H. Hori and N. Ikota, Org. Biomol. Chem., 2003, 1, 1452; (e) I. Nakanishi, K. Miyazaki, T. Shimada, K. Ohkubo, S. Urano, N. Ikota, T. Ozawa, S. Fukuzumi and K. Fukuhara, J. Phys. Chem. A, 2002, 106, 11123; (f) I. Nakanishi, K. Fukuhara, T. Shimada, K. Ohkubo, Y. Iizuka, K. Inami, M. Mochizuki, S. Urano, S. Itoh, N. Miyata and S. Fukuzumi, J. Chem. Soc., Perkin Trans. 2, 2002, 1520.
- 23 (a) K. Mukai, M. Oi, A. Ouchi and S. Nagaoka, J. Phys. Chem. B, 2012, 116, 2615; (b) Y. Kohno, M. Fujii, C. Matsuoka, H. Hashimoto, A. Ouchi, S. Nagaoka and K. Mukai, J. Phys. Chem. B, 2011, 115, 9880; (c) A. Ouchi, S. Nagaoka, K. Abe and K. Abe, J. Phys. Chem. B, 2009, 113, 13322.
- 24 S. B. Kedrare and R. P. Singh, *J. Food Sci. Technol.*, 2011, 48, 412.
- 25 M. J. Ajitha, S. Mohanlal, C. H. Suresh and A. Jayalekshmy, J. Agric. Food Chem., 2012, 60, 3693.
- 26 M. Hidalgo, C. Sánchez-Moreno and S. Pascual-Teresa, *Food Chem.*, 2010, **121**, 691.

- 27 M. Locatelli, R. Gindro, F. Travaglia, J. Coïsson, M. Rinaldi and M. Arlorio, *Food Chem.*, 2009, **114**, 889.
- 28 O. P. Shama and T. K. Bhat, Food Chem., 2009, 113, 1202.
- 29 M. C. Foti and C. Daquino, *Chem. Commun.*, 2006, **30**, 3252.
- 30 W. Hakamata, I. Nakanishi, Y. Masuda, T. Shimizu, H. Higuchi, Y. Nakamura, S. Saito, S. Urano, T. Oku, T. Ozawa, N. Ikota, N. Miyata, H. Okuda and K. Fukuhara, *J. Am. Chem. Soc.*, 2006, **128**, 6524.
- 31 A. W. Taylor, S. Puttick and P. Licence, J. Am. Chem. Soc., 2012, 134, 15636.
- 32 I. Nakanishi, K. Ohkubo, K. Miyazaki, W. Hakamata, S. Urano, T. Ozawa, H. Okuda, S. Fukuzumi, N. Ikota and K. Fukuhara, *Chem. Res. Toxicol.*, 2004, **17**, 26.
- 33 (a) A. M. Bond and D. E. Smith, Anal. Chem., 1974, 46, 1946; (b) T. G. McCord and D. E. Smith, Anal. Chem., 1969, 41, 1423.
- 34 E. M. Arnett, K. Amarnath, N. G. Harvey and J. P. Cheng, J. Am. Chem. Soc., 1990, 112, 344.
- 35 M. R. Wasielewski and R. Breslow, J. Am. Chem. Soc., 1976, 98, 4222.
- 36 A. J. Bard and L. R. Faulkner, *Electrochemical Methods, Fundamental and Applications*, John Wiley & Sons, New York, 2001, ch. 10, pp. 368–416.
- 37 C. K. Mann and K. K. Barnes, *Electrochemical Reactions in Non-aqueous Systems*, Mercel Dekker, New York, 1970.
- 38 J. H. Baxendale and H. R. Hardy, *Trans. Faraday Soc.*, 1953, 49, 1140.
- 39 X. O. Zhu, C. H. Wang and H. Liang, *J. Org. Chem.*, 2010, 75, 7240.
- 40 (a) R. G. Pearson, J. Chem. Educ., 1968, 45, 581;
 (b) R. G. Peason, J. Chem. Educ., 1968, 45, 643.
- 41 (a) R. A. Marcus, Annu. Rev. Phys. Chem., 1964, 15, 155;
 (b) R. A. Marcus, Angew. Chem., Int. Ed. Engl., 1993, 32, 1111;
 (c) R. A. Marcus and N. Sutin, Biochim. Biophys. Acta, 1985, 811, 265.
- 42 (a) H. Shizuka, M. Nakamura and T. Morirta, J. Phys. Chem., 1980, 84, 989; (b) H. Shizuka, T. Saito and T. Morirta, Chem. Phys. Lett., 1978, 56, 519.
- 43 The term $1/k_{\text{diff}}$ is negligible where k_{diff} is significantly larger than *k* in this study.
- 44 Large reorganization energies of electron transfer in the presence of a strong Lewis acid have been reported and discussed therein. (a) K. Ohkubo, R. Garcia, P. J. Sintic, T. Khoury, M. J. Crossley, K. M. Kadish and S. Fukuzumi, *Chem.-Eur. J.*, 2009, 15, 10493; (b) K. Okamoto, H. Imahori and S. Fukuzumi, *J. Am. Chem. Soc.*, 2003, 125, 7014; (c) K. Okamoto, Y. Mori, H. Yamada, H. Imahori and S. Fukuzumi, *Chem. Eur. J.*, 2004, 10, 474.