Organic & Biomolecular Chemistry



CORRECTION

View Article Online



Cite this: *Org. Biomol. Chem.*, 2016, **14**, 1840

Correction: Copper-mediated cyanation of indoles and electron-rich arenes using DMF as a single surrogate

Lianpeng Zhang, Ping Lu* and Yanguang Wang*

DOI: 10.1039/c6ob90012g www.rsc.org/obc

Correction for 'Copper-mediated cyanation of indoles and electron-rich arenes using DMF as a single surrogate' by Lianpeng Zhang et al., Org. Biomol. Chem., 2015, **13**, 8322–8329.

The authors regret that Table 3 was given incorrectly. The correct Table 3 is shown below.

Table 3 Copper-mediated cyanation of aryl aldehydes^a

ArCHO 4	O CC	u(NO ₃) ₂ ·3H ₂ O (1.0 HOAc (8 equiv) TBHP (2 equiv) air, 130 °C, 48 h	→ ArCN
R ³ N H	CICN	CN	R ² N
2a , $R^3 = H$, 74% 2x , 55% 2h , $R^2 = Ph$, 85% 2b , 82%			
2u , $R^3 = OCH_3$, 52% 2i , $R^2 = 4-FC_6H_4$, 70%			
2v , R ³ = CN, 75%			
2w , $R^3 = CI$, 57%		CN	CN
CN	ÇN	Ĭ	
			H ₃ CO OCH ₃
S	Y)	1 0	OCH ₃
• • • • • • • • • • • • • • • • • • • •	0-/	0-	•
2y , 35%	2z , 60%	2A , 65%	2B , 90%
CN OCH ₃	CN CN NO ₂	CN	CN
2C, 70% 2D	2E , 20	% 2F , 45%	2G , 65%
(140 °C)			

 $[^]a$ Reaction conditions: 3 (0.5 mmol), Cu(NO₃₎₂·3H₂O (0.5 mmol), HOAc (4.0 mmol), TBHP (1.0 mmol), DMF (3.0 mL), air, 130 $^{\circ}$ C, 48 h.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.