

CORRECTION

[View Article Online](#)
[View Journal](#) | [View Issue](#)



Cite this: *Polym. Chem.*, 2016, **7**, 1682

DOI: 10.1039/c6py90024k

www.rsc.org/polymers

Correction: Chemistry of aromatic polythioesters and polydithioesters

Daisuke Abe, Yuichiro Fukuda and Yuji Sasanuma*

Correction for 'Chemistry of aromatic polythioesters and polydithioesters' by Daisuke Abe, et al., *Polym. Chem.*, 2015, **6**, 3131–3142.

The authors regret the wrong arrangement of the *trans* fractions derived from MO calculations for the S–CH₂ bond in Table 1 of the original paper. The corrected version of Table 1 is as shown below.

Table 1 Vicinal coupling constants and *trans* fractions (*p_t*'s) of 3DBS₂ and 3DBS₄

Medium	Temp. (°C)	³ J _{HH} ^a	³ J' _{HH} ^a	³ J _{CH} ^a	<i>p_t</i>		S–CH ₂ ^d	
					<i>CH₂–CH₂</i>			
					Set A ^b	Set B ^c		
3DBS ₂ (NMR expt)								
Cyclohexane-d ₁₂	15	6.61	7.50	4.55	0.42	0.44	0.14	
	25	6.62	7.47	4.53	0.42	0.44	0.15	
	35	6.66	7.47	4.52	0.42	0.43	0.15	
	45	6.66	7.47	4.46	0.42	0.43	0.16	
	55	6.72	7.46	4.46	0.41	0.43	0.16	
Benzene-d ₆	25	6.47	7.84	4.46	0.45	0.47	0.16	
Dimethyl-d ₆ sulfoxide	25	6.40	7.94	4.48	0.47	0.48	0.16	
3DBS ₂ (MO calc.)								
Gas	15				0.44		0.11	
	25				0.44		0.12	
	35				0.44		0.12	
	45				0.43		0.12	
	55				0.43		0.12	
Benzene	25				0.46		0.14	
3DBS ₄ (NMR expt)								
Cyclohexane-d ₁₂	15	6.48	7.89	5.40	0.47	0.47	0.20	
	25	6.50	7.89	5.37	0.46	0.47	0.20	
	35	6.50	7.88	5.34	0.46	0.47	0.21	
	45	6.54	7.84	5.30	0.46	0.46	0.21	
	55	6.54	7.83	5.27	0.46	0.46	0.22	
Benzene-d ₆	25	6.43	8.14	5.25	0.48	0.49	0.22	
Dimethyl-d ₆ sulfoxide	25	6.35	8.11	5.35	0.49	0.49	0.20	
3DBS ₄ (MO calc.)								
Gas	15				0.38		0.19	
	25				0.38		0.19	
	35				0.38		0.20	
	45				0.38		0.20	
	55				0.38		0.21	
Benzene	25				0.41		0.22	

^a In Hz. ^b From MO calculations: $J_T = 12.55$, $J_G = 3.81$, $J'_T = 11.58$, $J'_G = 2.40$, $J''_G = 3.59$, and $J'''_G = 2.79$ Hz for 3DBS₂; $J_T = 12.58$, $J_G = 3.77$, $J'_T = 11.76$, $J'_G = 2.39$, $J''_G = 3.81$, and $J'''_G = 2.70$ Hz for 3DBS₄ (this study). ^c From experimental J_T and J_G values of 2-*tert*-butyl-1,3-dithiane.²⁴ ^d From MO calculations: $J_G = 1.08$, $J'_T = 6.39$, and $J'_G = 3.86$ Hz for 3DBS₂; $J_G = 1.14$, $J'_T = 8.17$, and $J'_G = 4.69$ Hz for 3DBS₄ (this study).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

