



CrossMark  
 click for updates

Cite this: *RSC Adv.*, 2016, 6, 79236

## Correction: Synergic therapy of melanoma using GNRs-MUA-PEI/siIDO2-FA through targeted gene silencing and plasmonic photothermia

Yujuan Zhang,<sup>\*ab</sup> Na Song,<sup>ab</sup> Jiamin Fu,<sup>ab</sup> Yanling Liu,<sup>ab</sup> Xuelin Zhan,<sup>ab</sup>  
 Shanshan Peng,<sup>ab</sup> Zhi Yang,<sup>ab</sup> Xianfang Zhu,<sup>c</sup> Yiguo Chen,<sup>ab</sup> Zhigang Wang,<sup>ab</sup>  
 Yanrong Yu,<sup>ab</sup> Qiaofa Shi,<sup>a</sup> Yingyuan Fu,<sup>a</sup> Keng Yuan,<sup>ab</sup> Nanjin Zhou,<sup>ab</sup>  
 Thomas E. Ichim<sup>d</sup> and Weiping Min<sup>\*abe</sup>

DOI: 10.1039/c6ra90075e

[www.rsc.org/advances](http://www.rsc.org/advances)

Correction for 'Synergic therapy of melanoma using GNRs-MUA-PEI/siIDO2-FA through targeted gene silencing and plasmonic photothermia' by Yujuan Zhang *et al.*, *RSC Adv.*, 2016, 6, 77577–77589.

The authors regret that Fig. 6 in the original manuscript requires correction. The titles for subparts 'a' and 'd' within Fig. 6C should state 'PBS' and 'PBS+Laser' rather than 'BSA' and 'BSA+Laser', respectively. The amended figure is shown below.

<sup>a</sup>Institute of Immunotherapy and College of Basic Medicine, Nanchang University, Jiangxi Academy of Medical Sciences, Nanchang, China. E-mail: yujuanzhang@ncu.edu.cn


<sup>b</sup>Jiangxi Provincial Key Laboratory of Immunotherapy, Nanchang, China

<sup>c</sup>China-Australia Joint Laboratory for Functional Nanomaterials, School of Physics and Mechanical and Electrical Engineering, Xiamen University, Xiamen, China

<sup>d</sup>Batu Biologics Inc, San Diego, California, USA

<sup>e</sup>Department of Surgery, Pathology and Oncology, University of Western Ontario, London, Canada. E-mail: mweiping@uwo.ca





**Fig. 6** *In vitro* photothermal effects of GNR-MUA-PEI-FA and GNR-MUA-PEI/siIDO2-FA. B16-BL6 cells were incubated with a final concentration of  $15 \mu\text{g mL}^{-1}$  of GNR-MUA-PEI-FA or PBS overnight. Subsequently, the cells were irradiated at (A)  $2 \text{ W cm}^{-2}$  for various times (0 minutes, 0.5 minutes, 1 minute, 2 minutes, 5 minutes, 10 minutes) or for (B) 5 minutes with different power densities ( $0 \text{ W cm}^{-2}$ ,  $0.5 \text{ W cm}^{-2}$ ,  $1 \text{ W cm}^{-2}$ ,  $2 \text{ W cm}^{-2}$ ,  $4 \text{ W cm}^{-2}$ ). After 24 hours, the cell viabilities were measured by the MTT assay and the percentages of dead cells = (cell viabilities before laser irradiation – cell viabilities after laser irradiation)/(cell viabilities before laser irradiation). (C) Apoptosis of tumor cells induced by GNR-MUA-PEI/siIDO2-FA. B16-BL6 cells were incubated with PBS (a) or GNR-MUA-PEI/siGL2-FA (b), or GNR-MUA-PEI-FA-siIDO2 (c), at the final concentration of  $16 \mu\text{g mL}^{-1}$  of wt((GNR-MUA-PEI-FA) : wt(siIDO2) = 15 : 1) overnight. Cells were either not irradiated (a–c) or were irradiated (d–f) at  $2 \text{ W cm}^{-2}$  for 5 minutes using a laser with a wavelength of 808 nm. The apoptotic and necrotic cell populations were determined at 24 hours by Annexin V-FITC/PI Apoptosis Detection Kit and analyzed by flow cytometry. Error bars represent the standard deviation of 3 experiments.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

