


Cite this: *RSC Adv.*, 2017, 7, 25171

Received 28th January 2017

Accepted 4th May 2017

DOI: 10.1039/c7ra01229b

rsc.li/rsc-advances

Direct C(sp²)-H amination of aryl aldehyde-derived hydrazones *via* visible light promoted photoredox catalysis†

Xin Zhu,^a Zhi He,^a Qiu-Yan Li^b and Xiao-Jun Wang^{id}*^{bc}

An unprecedented visible-light-induced direct C(sp²)-H amination of aryl aldehyde-derived hydrazones was developed by using *N*-acyloxypthalimides as nitrogen-radical precursors. This reaction represents a new way to substituted hydrazones with amination of carbon–nitrogen π bonds. A radical C–H amination mechanism is proposed.

Aldehyde-derived hydrazones belong to an important class of chemicals because of their many applications in synthetic organic chemistry.¹ Notably, they are attractive as umpolung carbonyl reagents because of the presence of the electron-releasing amino component that activates the azomethine (CH=N) carbon atom position toward electrophilic substitution. Much research attention has been devoted to the preparation of functionalized aromatic aldehyde hydrazones.² Recent attention has shifted toward the achievement of using the hydrazones' C=N bond as various radical acceptors for their diverse functionalizations (Scheme 1).^{3–6} Along this line, Baudoin and co-workers have reported an efficient procedure for the CuCl catalyzed trifluoromethylation of *N,N*-dialkylhydrazones under Togni reagent.³ In 2016, the groups of Hashmi and Zhu presented a work of difluoroalkylation of aldehyde *N,N*-dialkylhydrazones by CF₂ radical.^{4,5} Then Zhu's group reported a procedure for the C–H bond phosphonation of aldehyde *N,N*-dialkylhydrazones with the commercially available diphenylphosphine oxide.⁶ For the limited work of presented studies, substantial effort need to be devoted to the development of different radical (such as N, Si and B) to activate the hydrazones' C–H bond.†

Due to the abundant, renewable, inexpensive and eco-friendly energy source, visible-light promoted photocatalysis

has recently been employed as a powerful strategy in synthetic organic chemistry.⁷ Many novel visible-light mediated reactions were developed for the efficient and rapid preparation of fine chemicals over the past years.^{7,8} In the context, the addition of nitrogen-central radical generated by photo-catalysis to a π acceptor has been developed as a straightforward method to access structurally diverse *N*-containing compounds.⁹ Nevertheless, despite these important progress, existing methods for radical-type amination are largely limited to carbon–carbon π bonds (alkene, alkyne, or arene). For example, MacMillan *et al.* developed an elegant *N*-radical-mediated enantioselective direct α -amination of aldehydes by rationally merging visible-light photoredox catalysis with asymmetric aminocatalysis in photoredox-catalyzed C–H amination of arenes and heteroarenes with the use of *N*-acyloxypthalimides as the *N*-radical precursor.¹⁰ Shortly thereafter, Yu *et al.* developed a similar strategy by using a readily accessible hydroxylamine derivative as an *N*-radical precursor, which proved to be an efficient and broadly applicable visible-light photoredox-catalyzed direct C–H bond amination of various heteroarenes, including indoles, pyrroles, and furans.¹¹

Inspired by the pioneering works, we wondered that a *N*-radical precursor with good leaving group could be easy to generate a *N*-mediated radical and here present a work of direct C–H amination of aldehyde *N,N*-dialkyl hydrazones by visible

^aCollege of Chemistry, Beijing Normal University, Beijing 100875, P. R. China


^bJiangsu Key Laboratory of Green Synthetic Chemistry for Functional Materials, School of Chemistry and Materials Science, Jiangsu Normal University, Xuzhou 221116, P. R. China. E-mail: xjwang@jsnu.edu.cn

^cKey Laboratory of Photochemical Conversion and Optoelectronic Materials, Technical Institute of Physics and Chemistry, Chinese Academy of Sciences, Beijing 100190, P. R. China


† Electronic supplementary information (ESI) available: Experiment details and compounds characterization. CCDC 1543507. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7ra01229b

‡ Crystal data for 7b: CCDC 1543507, C₂₀H₁₆N₃O₃F₃, *M*_r = 403.36, orthorhombic, space group *Pbcn*, *T* = 153 K, *a* = 19.520(4) Å, *b* = 13.946(3) Å, *c* = 13.312(3) Å, $\alpha = \beta = \gamma = 90^\circ$, *V* = 3623.8(13) Å³, λ = 0.71073 Å, *Z* = 8, independent reflections 4113, *R*_{int} = 0.0501, *R*₁ [*I* > 2 σ (*I*)] = 0.0669, *wR*₂ (all data) = 0.1341, GOF = 1.251.

a) Previous work



b) Present work


Scheme 1 Radical addition to carbon–nitrogen π bond.



light irradiation. To the best of our knowledge, the direct C(sp²)-H amination of benzaldehyde or its synthetic equivalents (aromatic aldehyde hydrazones) through a radical way is unexplored, but could complement existing strategies. Herein we reported an efficient and mild procedure for the amination of aromatic aldehyde hydrazones based on *N*-(acyloxy)phthalimides as aminating reagents under visible-light irradiation.

The generation of a *N*-central radical by visible-light photo-redox catalysis recently has emerged as a powerful tool in synthetic organic chemistry.^{7e} On the basis of these previous reports, we initiated the reaction of *N*-morpholino-1-phenylmethanimine (1a) and various nitrogen radical precursors by using *fac*-Ir(ppy)₃ as photocatalyst owing to its superior reduction capacity upon excitation by visible light (Scheme 2). After irradiation by fluorescent light bulb for 24 h at room temperature, it was found only precursor 6b with strong electron-deficient group gave a slight yield (7%).

Encouraged by this preliminary result, we further optimized the reaction condition. The desired product (7a) was obtained by employing aldehyde hydrazone (1a) and the *N*-(acyloxy)phthalimides as the model substrates and by using the photo-redox catalyst Ir(ppy)₃ with visible light in MeCN (Table 1). According to the Sanford group's report that the use of more electron-withdrawing R-substituents would enhance the leaving group ability of the carboxylate anion, thereby favoring fragmentation to release RCO²⁻ with generation of the *N*-centered phthalimidyl radical (PhthN[•]). The PhthN[•] then could participate in C-H amination of diverse substrates. Our initial efforts toward achieving this goal focused on the changing of leaving group of *N*-(acyloxy)phthalimides. To achieve this goal, we examined the C-H amination of aldehyde hydrazone (1a) with a series of electronically different *N*-(acyloxy)phthalimides in the presence of 2 mol% of Ir(ppy)₃ and visible light in MeCN. As shown in Table 1, *N*-(acyloxy)phthalimide 6a (R = CH₃, entry 1) did not react to afford C-H amination product 7a but with the starting materials consumed. However, the increase of electron-withdrawing ability of R group in 6 derivatives resulted in the improved yield. In particular, 6f gave the best yield (18%).

To optimize the reaction condition, the solvents were varied together with using different additions. After solvent screening, dimethyl sulfoxide (DMSO) was found to be the most suitable for this reaction (Table 2, entry 3). Contrary to expectations, the

Table 1 Visible light photocatalyzed C-H amination of hydrazones with *N*-(acyloxy)phthalimide derivatives^a



Entry	R	Yield ^b (%)
1	CH ₃ CO (6a)	nd ^c
2	CF ₃ CO (6b)	7
3	PhCO (6c)	<5
4	<i>p</i> -MeOPhCO (6d)	<5
5	<i>p</i> -CF ₃ PhCO (6e)	11
6	C ₆ F ₅ CO (6f)	18
7	Bs (6g)	8
8	Ns (6h)	<5


^a Reaction condition: 1a (0.1 mmol), 2 (0.3 mmol, 3 equiv.), Ir(ppy)₃ (0.002 mmol, 2 mol%), MeCN (1.0 mL), at room temperature, fluorescent light bulb, 24 h. ^b Yields determined by GC-mass, decahydronaphthalene used as internal standard. ^c Starting materials were consumed and no products were detected by GC-mass. nd = not detected. Bs = benzenesulfonyl, Ns = 4-nitrobenzenesulfonyl.

Table 2 Representative examples from the optimization of the reaction conditions^a



Entry	Base	Solvent	Yield ^b (%)
1	—	MeCN	8
2	—	DMF	<5
3	—	DMSO	52
4	—	DCM	<5
5	K ₂ HPO ₄	DMSO	<5
6	KOAc	DMSO	<5
7	NaHCO ₃	DMSO	<5
8	CsF	DMSO	<5
9 ^c	—	DMSO	0
10 ^d	—	DMSO	0
11 ^e	—	DMSO	56
12 ^f	—	DMSO	87

^a Reaction condition: 1a (0.1 mmol), 2a (0.3 mmol, 3 equiv.), base (0.15 mmol, 1.5 equiv.), Ir(ppy)₃ (0.002 mmol, 2 mol%), solvent (1.0 mL), at room temperature, fluorescent light bulb, 24 h. ^b Yields determined by GC-mass, decahydro-naphthalene used as internal standard. ^c In dark. ^d Without Ir(ppy)₃. ^e With blue LEDs (20 W). ^f With blue LEDs (20 W) and 48 h.



Scheme 2 Screening for an effective *N*-radical precursors. Reaction condition: 1a (0.1 mmol), nitrogen source (3 equiv.), Ir(ppy)₃ (0.002 mmol, 2 mol%), MeCN (1.5 mL), at room temperature, fluorescent light bulb, 24h.^aYields determined by GC-MS.


bases were bad for the reaction (Table 2, entries 5–8). We then speculated that the nitrogen precursor was easily decomposed under the inorganic bases and could not generate the *N*-radical. Control experiments showed that the reaction could not



proceed in the absence of either visible-light irradiation or the photoredox catalyst (Table 2, entries 9 and 10). And we also found that the reaction got a good yield under visible light irradiation from blue LEDs instead of fluorescent light bulb after 48 h (Table 2, entries 12).


With the optimized condition established, we next investigated the effect of the *N*-substituent (Scheme 3). The results showed that this reaction did not tolerate secondary amino groups, as a complex mixture of product was obtained in the case of 1a-c-1ab, even though the starting materials were consumed. These experimental results indicated that the *N,N*-dialkyl structural motif was crucial for this transformation.

Furthermore, a plausible photo-catalytic cycle is proposed in Scheme 4. First, upon visible-light irradiation, the photocatalyst Ir(ppy)₃ undergoes a metal to ligand charge-transfer (MLCT) process to produce the strongly reducing excited state Ir(ppy)₃^{*} (step a). Single electron transfer from Ir(ppy)₃^{*} to 6f results in fragmentation of the *N*-(acyloxy)phthalimide with the generation of an *N*-centered phthalimidyl radical (PhthN[•]) and Ir(ppy)₃⁺ (step b). Subsequent addition of PhthN[•] to the C=N π bond leads to an aminyl radical intermediate 9 (step c), which then undergoes oxidation and deprotonation of the aminyl cation, giving aminated product 7a. When 2,2,6,6-tetramethylpiperidine 1-oxyl (TEMPO) was added into the reaction mixture under the standard condition, only trace amount of product 7a can be




Scheme 5 Investigation into the reaction mechanism.


Table 3 Substrate scope of hydrazones^{a,b}



^a Reaction condition: hydrazone (0.1 mmol), 7f (0.3 mmol, 3 equiv.), Ir(ppy)₃ (0.002 mmol, 2 mol%), DMSO (1.0 mL), at room temperature, blue LEDs (20 W), 48 h. ^b Isolated yields.



Scheme 3 Investigation of the effect of the *N*-substituent.



Scheme 4 Proposed the photocatalyzed reaction mechanism.

detected (Scheme 5). It demonstrated that this reaction generated radical which could be trapped by TEMPO, which suggested the single electron transfer process during the reaction.

The amination of various aromatic *N,N*-dialkyl hydrazones under the optimized reaction conditions was next examined to gauge the scope of the reaction (Table 3). Aryl aldehyde-derived hydrazones bearing either electron-donating or electron-withdrawing substituents furnished the corresponding products. And electron-withdrawing (7b,7c) gave a slightly better yield. The C–H amination reaction of heterocyclic hydrazones or aliphatic aldehyde-derived hydrazones was discouraged in the present condition (7l and 7m).

In conclusion, we have developed an efficient and mild visible-light photoredox-catalyzed C–H amination of aryl aldehyde-derived hydrazones for the first time, where *N*-acyloxyphthalimides were employed as nitrogen-radical precursors. And it is anticipated that the present strategy could be used for more useful transformation of C–N π bond in the future.

Acknowledgements

This work was financially supported from Beijing Normal University, National Natural Science Foundation of China



(21302072) and Key Laboratory of Photochemical Conversion and Optoelectronic Materials, TIPC, CAS (PCOM201711).

Notes and references

- (a) R. Lazny and A. Nodzevska, *Chem. Rev.*, 2010, **110**, 1386–1434; (b) R. Brehme, D. Enders, R. Fernandez and J. M. Lassaletta, *Eur. J. Org. Chem.*, 2007, 5629–5660.
- (a) G. K. Friestad, *Tetrahedron*, 2001, **57**, 5461–5496; (b) G. K. Friestad, Y. Shen and E. L. Ruggles, *Angew. Chem., Int. Ed.*, 2003, **42**, 5061–5063; (c) K. Yamada and K. Tomioka, *Chem. Rev.*, 2008, **108**, 2874–2886; (d) H. Miyabe, *Synlett*, 2012, **23**, 1709–1724.
- E. Pair, N. Monteiro, D. Bouyssi and O. Baudoin, *Angew. Chem., Int. Ed.*, 2013, **52**, 5346–5349.
- J. Xie, T. Zhang, F. Chen, N. Mehrkens, F. Rominger, M. Rudolph and A. S. K. Hashmi, *Angew. Chem., Int. Ed.*, 2016, **55**, 2934–2938.
- P. Xu, G. Wang, Y. Zhu, W. Li, Y. Cheng, S. Li and C. Zhu, *Angew. Chem., Int. Ed.*, 2016, **55**, 2939–2943.
- P. Xu, Z. Wu, N. Zhou and C. Zhu, *Org. Lett.*, 2016, **18**, 1143–1145.
- (a) M. H. Shaw, J. Twilton and D. W. C. MacMillan, *J. Org. Chem.*, 2016, **81**, 6898–6926; (b) N. A. Romero and D. A. Nicewicz, *Chem. Rev.*, 2016, **116**, 10075–10166; (c) T. P. Yoon, M. A. Ischay and J. Du, *Nat. Chem.*, 2010, **2**, 527–532; (d) J. Xuan and W.-J. Xiao, *Angew. Chem., Int. Ed.*, 2012, **51**, 6828–6838; (e) J.-R. Chen, X.-Q. Hu, L.-Q. Lu and W.-J. Xiao, *Chem. Soc. Rev.*, 2016, **45**, 2044–2056.
- (a) Y.-W. Zheng, B. Chen, P. Ye, K. Feng, W. Wang, Q.-Y. Meng, L.-Z. Wu and C.-H. Tung, *J. Am. Chem. Soc.*, 2016, **138**, 10080–10083; (b) L.-M. Zhao, Q.-Y. Meng, X.-B. Fan, C. Ye, X.-B. Li, B. Chen, V. Ramamurthy, C.-H. Tung and L.-Z. Wu, *Angew. Chem., Int. Ed.*, 2017, **56**, 3020–3024; (c) G. Zhang, C. Liu, H. Yi, Q. Meng, C. Bian, H. Chen, J.-X. Jian, L.-Z. Wu and A. Lei, *J. Am. Chem. Soc.*, 2015, **137**, 9273–9280; (d) Q.-Y. Li, Z. Ma, W.-Q. Zhang, J.-L. Xu, W. Wei, H. Lu, X. Zhao and X.-J. Wang, *Chem. Commun.*, 2016, **52**, 11284–11287; (e) W.-Q. Zhang, Q.-Y. Li, Q. Zhang, Y. Lu, H. Lu, W. Wang, X. Zhao and X.-J. Wang, *Inorg. Chem.*, 2016, **55**, 1005–1007; (f) Q.-Y. Meng, J.-J. Zhong, Q. Liu, X.-W. Gao, H.-H. Zhang, T. Lei, Z.-J. Li, K. Feng, B. Chen, C.-H. Tung and L.-Z. Wu, *J. Am. Chem. Soc.*, 2013, **135**, 19052–19055.
- (a) S. Z. Zard, *Chem. Soc. Rev.*, 2008, **37**, 1603–1618; (b) L. Shi and W. Xia, *Chem. Soc. Rev.*, 2012, **41**, 7687–7697; (c) J. Hu, J. Wang, T. H. Nguyen and N. Zheng, *Beilstein J. Org. Chem.*, 2013, **9**, 1977–2001; (d) S. H. Cho, J. Yoon and S. Chang, *J. Am. Chem. Soc.*, 2011, **133**, 5996–6005; (e) M. Louillat-Habermeyer, R. Jin and F. W. Patureau, *Angew. Chem., Int. Ed.*, 2015, **54**, 4102–4104; (f) L. J. Allen, P. J. Cabrera, M. Lee and M. S. Sanford, *J. Am. Chem. Soc.*, 2014, **136**, 5607–5610; (g) H. Jiang, X. An, K. Tong, T. Zheng, Y. Zhang and S. Yu, *Angew. Chem., Int. Ed.*, 2015, **54**, 4055–4059; (h) J. Davies, S. G. Booth, S. Essafi, R. A. W. Dryfe and D. Leonori, *Angew. Chem., Int. Ed.*, 2015, **127**, 14223–14227; (i) T. W. Greulich, C. G. Daniliuc and A. Studer, *Org. Lett.*, 2015, **17**, 254–257; (j) K. Miyazawa, T. Koike and M. Akita, *Chem.–Eur. J.*, 2015, **21**, 11677–11680.
- G. Cecere, C. M. Konig, J. L. Allea and D. W. MacMillan, *J. Am. Chem. Soc.*, 2013, **135**, 11521–11524.
- Q. Qin and S. Yu, *Org. Lett.*, 2014, **16**, 3504–3507.

