



Correction: Controlled generation and use of CO in flow

Cite this: *React. Chem. Eng.*, 2016, 1, 340

Steffen V. F. Hansen,^{ab} Zoe E. Wilson,^a Trond Ulven^{*b} and Steven V. Ley^{*a}

DOI: 10.1039/c6re90010k

Correction for 'Controlled generation and use of CO in flow' by Steffen V. F. Hansen *et al.*, *React. Chem. Eng.*, 2016, DOI: 10.1039/c6re00020g.

rsc.li/reaction-engineering

In the above article, Table 3 did not include a yield for entry 5. The correct information can be found in the new version of Table 3 below.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, UKsvl1000@cam.ac.uk

^b Department of Physics, Chemistry and Pharmacy, University of Southern Denmark, Campusvej 55, 5230 Odense M, Denmark. E-mail: ulven@sdu.dk





Table 3 Alkoxy carbonylation of vinyl and aryl iodides with modified set up

#	Iodide ^a	Product	Yield ^b	Iodide ^a	Product	Yield ^b	Iodide ^a	Product	Yield ^b
1 ^c			71%	4		74%	7 ^a		79%
2			97%	5		78%	8 ^d		96% (1.47 g)
3			99%	6		78%			

EtOH
0.25 mL.min⁻¹
Vapourtec E-series

^a All reactions were carried out on 0.5 mmol scale apart from #8 which was run continuously on 8 mmol scale (320 minutes). ^b Isolated yield. ^c Methanol was used as the solvent for pump 3, the reaction coil was heated to 25 °C and sample loop filled with iodide 5 (0.1 M), Pd(OAc)₂ (0.01 eq.), XantPhos (0.012 eq.), NEt₃ (1.5 eq.), H₂NNH₂ (0.3 eq.) and THF/MeOH (1:1). ^d Continuous flow reaction.