

CrossMark  
click for updatesCite this: *Chem. Sci.*, 2016, 7, 2170

# Radical transfer in *E. coli* ribonucleotide reductase: a NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ mutant unmasks a new conformation of the pathway residue 731<sup>†</sup>

Müge Kasanmascheff,<sup>‡,ab</sup> Wankyu Lee,<sup>‡,c</sup> Thomas U. Nick,<sup>a</sup> JoAnne Stubbe<sup>\*c</sup> and Marina Bennati<sup>\*ab</sup>

Ribonucleotide reductases (RNRs) catalyze the conversion of ribonucleotides to deoxyribonucleotides in all living organisms. The catalytic cycle of *E. coli* RNR involves a long-range proton-coupled electron transfer (PCET) from a tyrosyl radical (Y<sub>122</sub><sup>•</sup>) in subunit  $\beta$ 2 to a cysteine (C<sub>439</sub>) in the active site of subunit  $\alpha$ 2, which subsequently initiates nucleotide reduction. This oxidation occurs over 35 Å and involves a specific pathway of redox active amino acids (Y<sub>122</sub>  $\leftrightarrow$  [W<sub>48</sub>?]  $\leftrightarrow$  Y<sub>356</sub> in  $\beta$ 2 to Y<sub>731</sub>  $\leftrightarrow$  Y<sub>730</sub>  $\leftrightarrow$  C<sub>439</sub> in  $\alpha$ 2). The mechanisms of the PCET steps at the interface of the  $\alpha$ 2 $\beta$ 2 complex remain puzzling due to a lack of structural information for this region. Recently, DFT calculations on the 3-aminotyrosyl radical (NH<sub>2</sub>Y<sub>731</sub><sup>•</sup>)- $\alpha$ 2 trapped by incubation of NH<sub>2</sub>Y<sub>731</sub>- $\alpha$ 2/ $\beta$ 2/CDP(substrate)/ATP(allosteric effector) suggested that R<sub>411</sub>- $\alpha$ 2, a residue close to the  $\alpha$ 2 $\beta$ 2 interface, interacts with NH<sub>2</sub>Y<sub>731</sub><sup>•</sup> and accounts in part for its perturbed EPR parameters. To examine its role, we further modified NH<sub>2</sub>Y<sub>731</sub>- $\alpha$ 2 with a R<sub>411</sub>A substitution. NH<sub>2</sub>Y<sub>731</sub><sup>•</sup>/R<sub>411</sub>A generated upon incubation of NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2/ $\beta$ 2/CDP/ATP was investigated using multi-frequency (34, 94 and 263 GHz) EPR, 34 GHz pulsed electron–electron double resonance (PELDOR) and electron–nuclear double resonance (ENDOR) spectroscopies. The data indicate a large conformational change in NH<sub>2</sub>Y<sub>731</sub><sup>•</sup>/R<sub>411</sub>A relative to the NH<sub>2</sub>Y<sub>731</sub><sup>•</sup> single mutant. Particularly, the inter-spin distance from NH<sub>2</sub>Y<sub>731</sub><sup>•</sup>/R<sub>411</sub>A in one  $\alpha\beta$  pair to Y<sub>122</sub><sup>•</sup> in a second  $\alpha\beta$  pair decreases by 3 Å in the presence of the R<sub>411</sub>A mutation. This is the first experimental evidence for the flexibility of pathway residue Y<sub>731</sub>- $\alpha$ 2 in an  $\alpha$ 2 $\beta$ 2 complex and suggests a role for R<sub>411</sub> in the stacked Y<sub>731</sub>/Y<sub>730</sub> conformation involved in collinear PCET. Furthermore, NH<sub>2</sub>Y<sub>731</sub><sup>•</sup>/R<sub>411</sub>A serves as a probe of the PCET process across the subunit interface.

Received 14th September 2015

Accepted 6th December 2015

DOI: 10.1039/c5sc03460d

www.rsc.org/chemicalscience

## Introduction

Coupling of electron and proton transfers between donors and acceptors in proteins are ubiquitous in biology and can occur in a stepwise or concerted fashion. The concerted case avoids high energy intermediates and is designated as proton coupled electron transfer (PCET).<sup>1</sup> The mechanisms of these couplings are fundamental to our understanding of photosynthesis, respiration, synthesis of DNA building blocks, and many other processes. Unresolved issues describing these mechanisms have been articulated in several recent comprehensive reviews, with different mechanisms dictated by transfer distances, protein environment and dynamics.<sup>2,3</sup> When the proton and

electron donor and acceptor are distinct, the mechanism involves orthogonal PCET; when the donor and acceptor are the same, it involves collinear PCET.<sup>4,5</sup> A different mechanism in which a proton is transferred through water chains over long distances in concert with electron transfer (ET) has also been recently studied and discussed extensively in model systems.<sup>6,7</sup> In all mechanistic cases, since the electrons and protons have very different masses, electrons tunnel over large distances (10–15 Å) while proton tunnelling is restricted to shorter distances, on the order of hydrogen bond lengths.<sup>1,8,9</sup> This distance dependence complicates the issue of proton management. One important representative of the diversity of PCET mechanisms in proteins is found in the class I ribonucleotide reductases (RNRs). These enzymes catalyze the conversion of nucleotides to deoxynucleotides, the monomeric precursors required for DNA replication and repair in all eukaryotic and some prokaryotic organisms.<sup>10,11</sup> In this paper, we use the *Escherichia coli* (*E. coli*) class Ia RNR as a model system to interrogate the PCET process across the interface of the two subunits of this enzyme, proposed to involve two redox active protein tyrosine residues, one on each subunit, and a water interface between the subunits.<sup>12</sup>

<sup>a</sup>Max Planck Institute for Biophysical Chemistry, 37077 Göttingen, Germany. E-mail: marina.bennati@mpibpc.mpg.de

<sup>b</sup>Department of Chemistry, University of Göttingen, 37077 Göttingen, Germany

<sup>c</sup>Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA. E-mail: stubbe@mit.edu

<sup>†</sup> Electronic supplementary information (ESI) available. See DOI: 10.1039/c5sc03460d

<sup>‡</sup> These authors contributed equally.



The *E. coli* RNR consists of two homodimeric subunits,  $\alpha 2$  and  $\beta 2$ .<sup>13</sup> The enzyme is active when a transient  $\alpha 2\beta 2$  complex is formed.<sup>14</sup>  $\alpha 2$  contains the active site for nucleotide reduction and two allosteric effector binding sites that regulate the specificity and the rate of reduction.<sup>15–19</sup>  $\beta 2$  harbors the essential di-iron tyrosyl radical cofactor ( $\text{Fe}_2^{\text{III}}\text{-Y}_{122}^{\cdot}$ ).<sup>20,21</sup> During each turnover,  $\text{Y}_{122}^{\cdot}\text{-}\beta 2$  oxidizes  $\text{C}_{439}\text{-}\alpha 2$  to a thiyl radical, which subsequently initiates dNDP production.<sup>11</sup> There are X-ray structures of the individual subunits, and a docking model of the  $\alpha 2\beta 2$  complex places  $\text{Y}_{122}^{\cdot}$  at a distance of about 35 Å from  $\text{C}_{439}$ .<sup>22,23</sup> These initial studies led to the first formulation of radical transfer (RT) in RNR *via* a radical hopping mechanism involving a pathway of conserved amino acids ( $\text{Y}_{122} \leftrightarrow [\text{W}_{48}^?] \leftrightarrow \text{Y}_{356}$  in  $\beta 2$  to  $\text{Y}_{731} \leftrightarrow \text{Y}_{730} \leftrightarrow \text{C}_{439}$  in  $\alpha 2$ ). Biochemical<sup>24</sup> and biophysical (EPR,<sup>25,26</sup> SAXS,<sup>27</sup> and cryoEM<sup>28</sup>) studies confirmed that the docking model provides a reasonable representation of *E. coli* RNR in its transient, active form and led to a detailed mechanism of RT over such a long distance.<sup>4,14,24</sup> Nevertheless, in wild-type (wt) *E. coli* RNR, the rate limiting step, conformational change(s) upon substrate and allosteric effector binding to  $\alpha 2$ , has prevented spectroscopic detection of any intermediates in this process.<sup>29</sup> The recent development of methods to site-specifically incorporate tyrosine analogs with altered  $\text{p}K_{\text{a}}$ s and reduction potentials has permitted the detection of pathway radical intermediates<sup>30–32</sup> and, combined with state-of-the-art EPR spectroscopy,<sup>12,26,33,34</sup> has started to reveal the molecular basis of the long-range RT in RNR.<sup>14</sup>

These experiments have led to the current model illustrated in Fig. 1, which involves orthogonal PCET<sup>35</sup> steps within subunit  $\beta 2$  and collinear PCET steps within the  $\alpha 2$  subunit.<sup>12,33</sup> However, the mechanism of the PCET process at the subunit interface between  $\text{Y}_{356}$  in  $\beta 2$  and  $\text{Y}_{731}$  in  $\alpha 2$  remains elusive, as structural information on the C-terminal 35 amino acids of  $\beta 2$ , including a putative proton acceptor  $\text{E}_{350}$  and  $\text{Y}_{356}$  (Fig. 1), is missing.<sup>22</sup>

Our recent high-field (HF) EPR/ENDOR and DFT investigations using the 3-aminotyrosine mutants  $\text{NH}_2\text{Y}_{730}\text{-}\alpha 2$  and

$\text{NH}_2\text{Y}_{731}\text{-}\alpha 2$ , which generate the corresponding  $\text{NH}_2\text{Y}^{\cdot}$  upon incubation with  $\beta 2$ , CDP (substrate) and ATP (allosteric effector), established that an unusual stacked conformation of residues 730 and 731, observed in some X-ray structures of  $\alpha 2$  (ref. 23 and 36) (see ESI, Fig. S1†), occurs in the  $\alpha 2\beta 2$  complex.<sup>12,33</sup> However, the X-ray structure of  $\text{NH}_2\text{Y}_{730}\text{-}\alpha 2$  (PDB 2XO4) alone exhibited multiple conformations for  $\text{Y}_{731}\text{-}\alpha 2$ , with one rotated away from  $\text{NH}_2\text{Y}_{730}\text{-}\alpha 2$  toward the  $\alpha 2\beta 2$  subunit interface.<sup>30</sup> This “flipped” conformation was accompanied by reorientations of  $\text{R}_{411}$  and  $\text{N}_{733}$  in  $\alpha 2$ . Further comparison of  $\text{NH}_2\text{Y}_{730}\text{-}\alpha 2$ ,  $\text{NH}_2\text{Y}_{731}\text{-}\alpha 2$  and  $\text{NH}_2\text{Y}_{356}\text{-}\beta 2$  by HF EPR indicated that the electrostatic environment of all three transient  $\text{NH}_2\text{Y}^{\cdot}$ s is strongly perturbed and that their hydrogen bond interactions are intrinsically different.<sup>12,33</sup> Interestingly, one of our DFT models of the protein environment for  $\text{NH}_2\text{Y}_{731}\text{-}\alpha 2$  required  $\text{R}_{411}\text{-}\alpha 2$  to explain the perturbed  $g_x$  value observed and suggested that  $\text{R}_{411}\text{-}\alpha 2$  approaches to  $\text{NH}_2\text{Y}_{731}\text{-}\alpha 2$  within 2.6 Å (Fig. S1†).<sup>12</sup> Therefore, to examine the role of  $\text{R}_{411}\text{-}\alpha 2$  during the PCET process in *E. coli* RNR, we generated two mutants:  $\text{R}_{411}\text{A-}\alpha 2$  and the double mutant  $\text{NH}_2\text{Y}_{731}/\text{R}_{411}\text{A-}\alpha 2$ . Here, we report the incubation of  $\text{NH}_2\text{Y}_{731}/\text{R}_{411}\text{A-}\alpha 2$  with  $\beta 2$ /CDP and ATP, which generates the  $\text{NH}_2\text{Y}_{731}^{\cdot}/\text{R}_{411}\text{A-}\alpha 2\beta 2$  complex. Using advanced EPR methods, including 263 GHz pulse EPR and 34 GHz PELDOR/DEER (pulsed electron–electron double resonance) and ENDOR (electron–nuclear double resonance) spectroscopies, we have provided evidence for a new conformation of  $\text{NH}_2\text{Y}_{731}^{\cdot}/\text{R}_{411}$  that is “flipped” towards the subunit interface in the  $\alpha 2\beta 2$  complex. This is the first time an alternative conformation of any pathway tyrosine ( $\text{NH}_2\text{Y}_{731}^{\cdot}$ ) has been observed and it provides a new probe of the PCET mechanism across the subunit interface, which remains unknown.

## Experimental

### Materials

4-(2-Hydroxyethyl)-1-piperazineethanesulfonic acid (Hepes) was purchased from EMD Bioscience. Adenosine-5'-triphosphate (ATP), cytidine-5'-diphosphate (CDP), reduced  $\beta$ -nicotinamide adenine dinucleotide phosphate (NADPH), hydroxyurea (HU), kanamycin (Km), chloramphenicol (Cm), 2XYT media, M9 Minimal Salts, L-arabinose (ara),  $\beta$ -mercaptoethanol ( $\beta$ -ME), streptomycin sulfate and  $\text{NH}_2\text{Y}$  were purchased from Sigma-Aldrich. Isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) and 1,4-dithiothreitol (DTT) were purchased from Promega. Tris(2-carboxyethyl)phosphine (TCEP) hydrochloride was purchased from Thermo Scientific. Nucleotide primers were purchased from Invitrogen, and Pfu Ultra II polymerase was purchased from Stratagene.

### Site-directed mutagenesis to generate $\text{R}_{411}\text{A-}\alpha 2$ and $\text{NH}_2\text{Y}_{731}/\text{R}_{411}\text{A-}\alpha 2$

The Quikchange kit (Stratagene) was used to generate each mutant according to the manufacturer's protocol. The templates pET28a-nrdA and pET28a-nrdA  $\text{Y}_{731}\text{Z}^{30}$  were amplified with primer 5'-G CAG GAA CGT GCG TCT ACC GGT GCG ATC TAT ATT CAG AAC GTT GAC-3' and its reverse complement




Fig. 1 Current model of the long-range PCET in *E. coli* RNR.  $\text{R}_{411}$  is shown along with the redox active amino acid residues that are involved in this process ( $\text{Y}_{122} \leftrightarrow [\text{W}_{48}^?] \leftrightarrow \text{Y}_{356}$  in  $\beta 2$  to  $\text{Y}_{731} \leftrightarrow \text{Y}_{730} \leftrightarrow \text{C}_{439}$  in  $\alpha 2$ ).  $\text{W}_{48}$ ,  $\text{D}_{237}$  and  $\text{E}_{350}$  are shown in grey, because currently there is no evidence for their role in this process.<sup>14</sup> The locations of  $\text{Y}_{356}$  and  $\text{E}_{350}$  are unknown, as they are within the flexible C-terminal tail (35 amino acids) of  $\beta 2$ . Red, blue and purple arrows represent electron transfer, proton transfer and collinear PCET pathways, respectively.



and used to insert a GCG (Ala) at position 411. The sequences were confirmed by QuintaraBio Laboratory. All constructs contain an N-terminal (His)<sub>6</sub>-tag with a 10 amino acid linker.<sup>30</sup>

### Expression, purification and activity assays of R<sub>411</sub>A- $\alpha$ 2 and NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2

(His)<sub>6</sub>-wt- $\alpha$ 2 (2750 nmol min<sup>-1</sup> mg<sup>-1</sup>) and wt- $\beta$ 2 (7000 nmol min<sup>-1</sup> mg<sup>-1</sup>), and 1.2 Y'/ $\beta$ 2 were expressed and purified by standard protocols.<sup>30,37,38</sup> All  $\alpha$ 2 mutants were pre-reduced with 30 mM DTT and 15 mM HU before use.<sup>29</sup> *E. coli* thioredoxin (TR, 40 U mg<sup>-1</sup>) and thioredoxin reductase (TRR, 1800 U mg<sup>-1</sup>) used in assays were isolated as previously described.<sup>39,40</sup> (His)<sub>6</sub>-NH<sub>2</sub>Y<sub>731</sub>- $\alpha$ 2 was purified as previously described.<sup>30</sup> Expression and purification of R<sub>411</sub>A- $\alpha$ 2 and NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 followed previous protocols,<sup>30</sup> except that the purification buffer (50 mM Tris, 5% glycerol, 1 mM PMSF, pH 7.6) for NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 contained 1 mM TCEP. The yields of purified R<sub>411</sub>A- $\alpha$ 2 and NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 were 10–12 mg g<sup>-1</sup> and 6–7 mg g<sup>-1</sup> cell paste, respectively. The activity of R<sub>411</sub>A- $\alpha$ 2 (0.2  $\mu$ M) and NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 (1  $\mu$ M) was determined in the presence of 50-fold excess of wt- $\beta$ 2 with 3 mM ATP, 1 mM [<sup>3</sup>H]-CDP (4850 cpm nmol<sup>-1</sup>), 30  $\mu$ M TR, 0.5  $\mu$ M TRR, and 1 mM NADPH in assay buffer (50 mM HEPES, 1 mM EDTA, 15 mM MgSO<sub>4</sub>, pH 7.6). The amount of dCDP was determined by the method of Steeper and Steuart.<sup>41</sup> For single turnover experiments, NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 (5  $\mu$ M) was incubated with wt- $\beta$ 2 (5  $\mu$ M), 3 mM ATP, and 1 mM [<sup>3</sup>H]-CDP (20 000 cpm nmol<sup>-1</sup>) in assay buffer. The dissociation constant (*K<sub>d</sub>*) for R<sub>411</sub>A- $\alpha$ 2 and wt- $\beta$ 2 was determined in H<sub>2</sub>O and D<sub>2</sub>O buffers by the competitive inhibition assay<sup>42</sup> (SI-2, Fig. S2†).

### Samples for HF EPR and PELDOR spectroscopy

NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 and wt- $\beta$ 2 were mixed 1:1 to a final concentration of 160–180  $\mu$ M in D<sub>2</sub>O assay buffer as previously described.<sup>32,34</sup> These protein concentrations resulted in >95% binding between subunits. The reaction was initiated at room temperature by adding CDP and ATP to final concentrations of 1 and 3 mM, respectively. The reactions were manually freeze-quenched in liquid N<sub>2</sub> within 10–23 s. The PELDOR sample was prepared by adding glycerol-(OD)<sub>3</sub> to a final concentration of 10% (v/v) 16 s after the initiation of the reaction. This reaction was manually freeze-quenched after 56 s as just described. The NH<sub>2</sub>Y<sub>731</sub>· accounted for 30–33% of the total spin for all the samples used in this work, which was similar to the yields reported previously.<sup>30,32</sup>

### HF pulsed EPR spectroscopy

Echo-detected (ESE:  $\pi/2 - \tau - \pi - \text{echo}$ ) EPR spectra at 263 GHz were recorded on a Bruker Elexsys E780 quasi optical spectrometer using a single mode (TE<sub>011</sub>) cylindrical resonator (E9501610 – Bruker BioSpin) with a typical quality factor of 500–1000. The maximum microwave power coupled to the resonator was about 15 mW. Samples for 263 GHz EPR were inserted in capillaries (0.33 mm OD, Vitrocom CV2033S) with typical volumes of ca. 50 nL. 94 GHz ESE spectra were recorded on a Bruker E680 spectrometer with a 400 mW W-band power setup (Bruker power

upgrade – 2). Samples for 94 GHz ESE contained typical volumes of 2  $\mu$ L in 0.84 mm OD capillaries (WilmaD S6X84). All manually freeze-quenched samples were immersed in liquid N<sub>2</sub> and loaded into pre-cooled EPR cryostats.

### 34 GHz PELDOR spectroscopy

34 GHz ESE and PELDOR spectra were recorded on a Bruker E580 X/Q-band spectrometer equipped with a Bruker EN 5107D2 pulse EPR/ENDOR resonator. The spectrometer was power-upgraded with a Q-band TWT amplifier, providing about 170 W output power at 34.1 GHz. PELDOR experiments were recorded with an overcoupled resonator. The center of the mode was chosen for the pump frequency for measurements at 20 K. However, for measurements at 50 K the detection frequency was set in the center of the cavity mode to enhance detection sensitivity. Q-band samples contained typical volumes of 10  $\mu$ L in 1.6 mm OD capillaries (WilmaD 222T-RB).

### Processing and simulation of EPR spectra

Spectra were processed by phasing and baseline correction. Derivatives of the absorption spectra were obtained by fitting every four points with a second order polynomial and differentiating the function in MATLAB\_R2014b.<sup>43</sup> EPR spectra were simulated using the EasySpin-4.5.5 “pepper”-routine which was run in MATLAB.<sup>44</sup>

### DFT calculations

DFT calculations were performed with the ORCA 3.0.0 program package.<sup>45</sup> The geometry optimization of the neutral NH<sub>2</sub>Y· was performed using the unrestricted B3LYP<sup>46–48</sup> hybrid density functional in combination with the def2-TZVPP basis set and def2-TZVPP/JK auxiliary basis set.<sup>49,50</sup> To take into account the electrostatic environment of the radical intermediate at the protein interface, a solvation model (COSMO<sup>51</sup>) with the polarity of ethanol ( $\epsilon = 24$ ) was used. Otherwise, Grimme's dispersion correction<sup>52,53</sup> and RIJCOSX<sup>54</sup> approximations were employed. The energy converged to 10<sup>-9</sup> E<sub>h</sub>. The hyperfine couplings and *g* values were calculated using NH<sub>2</sub>Y·-C<sub>4</sub> as the gauge origin.<sup>55,56</sup> The def2-TZVPP basis set was consistent with the geometry optimization step.<sup>50</sup> The C2–C1–C $\beta$ –C $\alpha$  dihedral angle of the NH<sub>2</sub>Y· was changed stepwise with a geometry optimization for each step. The xyz coordinates for one of the optimized models are given in the ESI.†

### PyMOL models

The docking model refers to the  $\alpha$ 2 $\beta$ 2 complex structure generated from the individual wt- $\alpha$ 2 and wt- $\beta$ 2 X-ray structures.<sup>22,23</sup> In order to predict distances, the mutant *E. coli* RNR structure (PDB 2XO4)<sup>30</sup> was overlaid with the wt- $\alpha$ 2 structure in the docking model<sup>23</sup> using PyMOL, which first performs a sequence alignment and then aligns the structures to minimize the root mean square deviation between the structures.



## Results and discussion

### Preparation and characterization of R<sub>411</sub>A- $\alpha$ 2, NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 and ND<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$

Our recent studies on NH<sub>2</sub>Y<sub>731</sub>- $\alpha$ 2 (ref. 12) suggested that R<sub>411</sub> might interact with NH<sub>2</sub>Y<sub>731</sub>, partially accounting for the measured EPR and ENDOR parameters. To investigate this proposal, R<sub>411</sub>A- $\alpha$ 2 was generated and characterized. Because the mutation is proposed to be at the interface of  $\alpha$ 2 and  $\beta$ 2, the dissociation constant ( $K_d$ ) for subunit interactions was also examined and was determined to be  $0.94 \pm 0.33 \mu\text{M}$  (Fig. S2A†),  $\sim 5$  fold higher than that for wt- $\alpha$ 2 ( $0.18 \mu\text{M}$ ).<sup>42</sup> Under these conditions, this mutant was shown to have a specific activity of  $467 \pm 22 \text{ nmol min}^{-1} \text{ mg}^{-1}$ , 17% of that of the wt enzyme ( $2750 \text{ nmol min}^{-1} \text{ mg}^{-1}$ ). The reduced activity and weaker subunit binding suggest that R<sub>411</sub> plays a functional role.

Furthermore, we characterized the role of R<sub>411</sub> in the oxidation of Y<sub>731</sub>- $\alpha$ 2 by generating the double mutant NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2. The  $K_d$  for subunit interactions between NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 and wt- $\beta$ 2 was determined to be  $8 \pm 1 \text{ nM}$  (Fig. S2C†), which is consistent with the formation of a tight complex when a NH<sub>2</sub>Y<sup>•</sup> is generated.<sup>28</sup> Its specific activity was  $13 \pm 3 \text{ nmol min}^{-1} \text{ mg}^{-1}$ , 0.4% of the specific activity of wt-RNR and in the range of contaminating wt- $\alpha$ 2 activity.<sup>32</sup> A more sensitive, one turnover assay was then employed to determine if this double mutant could generate any dCDP. When pre-reduced NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 was mixed with wt- $\beta$ 2, CDP, and ATP for 5 min, only  $0.036 \pm 0.018 \text{ dCDP}/\alpha$ 2 was observed, consistent with contaminating wt- $\alpha$ 2. Thus, the double mutant is unable to make detectable dCDP, which is not unexpected, given the specific activities of the R<sub>411</sub>A and the NH<sub>2</sub>Y<sub>731</sub>- $\alpha$ 2 mutants (see also SI-3 and Fig. S3†).

We next investigated whether NH<sub>2</sub>Y<sub>731</sub> could be generated by NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2, despite its inability to make dCDPs. NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2, wt- $\beta$ 2, CDP and ATP were studied by stopped-flow (SF) spectroscopy and the reaction was monitored at 320 nm, the absorption feature associated with the NH<sub>2</sub>Y<sup>•</sup> (Fig. S4,† red). The data were split into two time domains: 5 ms to 6 s and 25 s to 100 s. In the first time domain, NH<sub>2</sub>Y<sub>731</sub> formation was fit to a double exponential with  $k_{\text{fast}}$  of  $3.6 \pm 0.5 \text{ s}^{-1}$  (amplitude 8%) and  $k_{\text{slow}}$  of  $0.47 \pm 0.03 \text{ s}^{-1}$  (amplitude 21%) (Table S1†). The rate constants for NH<sub>2</sub>Y<sub>731</sub> in the single mutant control were similar:  $k_{\text{fast}}$  of  $9.6 \pm 0.6 \text{ s}^{-1}$  and  $k_{\text{slow}}$  of  $0.8 \pm 0.1 \text{ s}^{-1}$ . However, in this case, the fast phase accounted for 27% and the slow phase accounted for 13% of the NH<sub>2</sub>Y<sub>731</sub>. The biphasic kinetics of NH<sub>2</sub>Y<sub>731</sub> formation in both cases is attributed to multiple conformations that give rise to NH<sub>2</sub>Y<sub>731</sub>.<sup>32</sup> From 25 s to 100 s, NH<sub>2</sub>Y<sub>731</sub> in the double mutant reaction disappeared with a  $k_{\text{obs}}$  of  $0.02 \pm 0.003 \text{ s}^{-1}$ , while with the single mutant, disappearance occurred with a  $k_{\text{obs}}$  of  $0.005 \pm 0.002 \text{ s}^{-1}$ . Analysis of the Y<sub>122</sub>- $\beta$  disappearance kinetics was unsuccessful at early time points due to the detection limits, as described in SI-4.

Given the distinct kinetics of our double mutant relative to the NH<sub>2</sub>Y<sub>731</sub>- $\alpha$ 2, the 9 GHz EPR spectrum of the sample generated from the reaction of NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 with wt- $\beta$ 2, ATP, and

CDP quenched after 25 s was recorded and is shown in Fig. S5A and C.† Subsequent to subtraction of Y<sub>122</sub>, 32% of the total spin is associated with NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 with no spin loss. This result is similar to that of the single mutant, NH<sub>2</sub>Y<sub>731</sub>.<sup>30,32</sup> A comparison of their spectra, as shown in Fig. S5B,† revealed substantial differences in their hyperfine interactions, suggesting that further characterization of this radical might provide insight into the function of R<sub>411</sub>. Therefore, the role of R<sub>411</sub> in the RT pathway was further studied with advanced EPR spectroscopy.

### HF EPR of ND<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2

To examine the generated ND<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2, we took advantage of the proximity of Y<sub>122</sub> to the di-iron cluster and its altered relaxation properties. Pulsed EPR spectra of ND<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 at 34, 94 and 263 GHz were recorded in D<sub>2</sub>O buffer at 70 K and are shown in Fig. 2A. The use of D<sub>2</sub>O considerably simplifies the EPR spectra due to the absence of <sup>1</sup>H hyperfine (hf) splittings arising from the amino protons. The ND<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 EPR spectrum at 34 GHz is mainly dominated by the large hf couplings with the deuterons of the amino group and the two C $\beta$ -methylene protons.<sup>34</sup> On the other hand, the 94 and 263 GHz EPR spectra are dominated by  $g$ -anisotropy, and the relative contributions of  $g$ - and hf-anisotropy are strongly dependent on the operating magnetic field. The  $g$  values of ND<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 are best resolved at 263 GHz and are consistent with the values from our previous ND<sub>2</sub>Y<sup>•</sup> studies.<sup>12,33</sup> The 94 GHz spectra reveal differences in the hf splitting of the C $\beta$ -methylene protons (Fig. 2A, marked with an arrow): the large hf splitting of the C $\beta$ -methylene proton visible in the central line of ND<sub>2</sub>Y<sub>731</sub>- $\alpha$ 2 (red) is missing in ND<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2 (black). This splitting is also absent in the 263 GHz spectrum. The EPR spectra were simulated iteratively to find a global solution for the contributing hf couplings. All of the EPR data and simulations, in which the previously reported<sup>34</sup> hf coupling for <sup>14</sup>N is used, are consistent with the NH<sub>2</sub>Y<sub>731</sub> generated in the NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$ 2/ $\beta$ 2 complex being a single, well-oriented radical species with one set of magnetic parameters, which are listed in Table 1 (see also Fig. S7†). This finding is not self-evident, as our previous experiments with other double mutants, NH<sub>2</sub>Y<sub>731</sub>/Y<sub>730</sub>F- $\alpha$ 2 and NH<sub>2</sub>Y<sub>730</sub>/C<sub>439</sub>A- $\alpha$ 2, showed distributions in  $g$  values indicative of multiple radical environments and/or molecular orientations.<sup>12</sup>

Interestingly, we do not observe changes in the  $g$  values between ND<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha$  and ND<sub>2</sub>Y<sub>731</sub>- $\alpha$ 2. This is unexpected because the  $g_x$  value is affected by the electrostatic environment of a radical,<sup>57</sup> and the R<sub>411</sub>A mutation has changed the local environment of ND<sub>2</sub>Y<sub>731</sub>, as demonstrated by the substantial changes in the C $\beta$ -methylene <sup>1</sup>H couplings (Table 1). These couplings are related to the dihedral angle  $\theta_{\text{C}\beta}$  between the C $\beta$ -H bond and the  $p_z$  orbital axis of C<sub>1</sub> (Fig. 2B), and therefore provide information on the molecular orientation of the tyrosyl and 3-aminotyrosyl radicals.<sup>34</sup> The dihedral angle can be extracted from the McConnell equation ( $a_{\text{iso}(\text{C}\beta\text{-H})} = B_1 \times \rho_{\text{C}1} \times \cos^2 \theta_{\text{C}\beta}$ ),<sup>58</sup> which provides a semi-empirical relationship for the observed isotropic constant  $a_{\text{iso}}$ . The C2-C1-C $\beta$ -C $\alpha$  angle of






Fig. 2 (A) Derivative EPR spectra (solid black lines) at 34 GHz (top), 94 GHz (middle) and 263 GHz (bottom) of  $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A}-\alpha 2$  with the corresponding simulations (dashed black lines). The 94 GHz EPR spectrum of  $\text{ND}_2\text{Y}_{731}^*-\alpha 2$  in the single mutant (solid red line) with the corresponding simulation (dashed red line) is shown for comparison. The difference between the two spectra around  $g_y$  is marked with an arrow. The glass signal is marked with an asterisk. Exp. conditions (34 GHz):  $\pi/2 = 6$  ns,  $\tau = 280$  ns, shot repetition time = 6 ms, shots/point = 80, number of scans = 10; (94 GHz):  $\pi/2 = 30$  ns,  $\tau = 280$  ns, shot repetition time = 5 ms, shots/point = 100, number of scans = 50–100; (263 GHz):  $\pi/2 = 40$  ns,  $\tau = 270$  ns, shot repetition time = 6 ms, shots/point = 250, number of scans = 36. Structure of  $\text{NH}_2\text{Y}^*$  is shown in the inset. (B) Orientation of the  $\text{C}_\beta$ -methylene protons with respect to the phenol ring, as extracted from the observed hyperfine couplings. (C)  $a_{\text{iso}}$  as a function of the dihedral angle for each  $\text{C}_\beta$ -methylene proton, calculated from a DFT model for  $\text{NH}_2\text{Y}^*$  (Fig. S6†).

Table 1 Summary of  $g$  values and large hf couplings ( $>8$  MHz) of  $\text{ND}_2\text{Y}_{731}^*$  in the double and single mutants<sup>a</sup>

Sample	$g$ values			$a_{\text{iso}}$ (MHz)		
	$g_x$	$g_y$	$g_z$	$\text{H}\beta_1$	$\text{H}\beta_2$	$^{14}\text{N}$
$\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A}-\alpha$	2.0051	2.0041	2.0022	10	10	12
$\text{ND}_2\text{Y}_{731}^*-\alpha^b$	2.0051	2.0041	2.0022	22	9	12

<sup>a</sup> Uncertainties in the  $g$  values and hf couplings are about 0.00005 and up to 10%, respectively, as obtained from the spectral simulations. <sup>b</sup>  $g$  values and hf couplings were reported in ref. 12 and 34, respectively.

$\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A}-\alpha 2$  is estimated to be  $\approx 90^\circ$  by using  $B_1$  of 162 MHz (ref. 59) for tyrosyl radicals, an electron spin density  $\rho_{\text{C}1}$  of 0.214,<sup>12</sup> and an isotropic  $\text{C}_\beta$ -methylene proton hf coupling  $a_{\text{iso}} = 10 \pm 1$  MHz (Table 1). This dihedral angle is indeed consistent with the hf couplings of the two  $\text{C}_\beta$ -methylene  $^1\text{H}$  resonances being indistinguishable, as reported in Table 1 and seen in Fig. 2B and C. This result was confirmed by DFT calculations on the observed hf couplings of  $\text{NH}_2\text{Y}^*$ , in which the ring orientation was modeled with respect to the backbone and showed a symmetric orientation relative to the  $p_z$  orbital

axis of  $\text{C}_1$  (Fig. 2B). In this calculation, a  $\theta_{\text{C}\beta}$  angle of  $90^\circ$  corresponds to  $a_{\text{iso}} = 9 \pm 3$  MHz (grey area in Fig. 2C) for both  $\text{C}_\beta$ -methylene protons,  $\text{H}_{\beta 2/1}$ .

### ENDOR for detection of hydrogen bonds to $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A}-\alpha 2$

Given that the  $\text{R}_{411}\text{A}$  mutation had little effect on  $g_x$ ,  $^2\text{H}$  ENDOR spectroscopy was used to further examine a possible correlation of the observed  $g_x$  value ( $g_x = 2.0051$ ) with the hydrogen bonding environment. Fig. 3 illustrates the  $^2\text{H}$  Mims ENDOR spectra of  $\text{ND}_2\text{Y}_{731}^*-\alpha 2$  and  $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A}-\alpha 2$ . Both spectra contain a broad signal that extends over  $\pm 2$  MHz, arising from the strongly coupled amino deuterons, which is a common feature of  $\text{ND}_2\text{Y}^*$  Mims ENDOR spectra.<sup>12,33</sup> However, we observe that the  $^2\text{H}$  hf tensor previously assigned to the moderately strong hydrogen bond between  $\text{Y}_{730}$  and  $\text{Y}_{731}$  in  $\text{ND}_2\text{Y}_{731}^*-\alpha 2$ , which is almost perpendicular to the tyrosine ring plane,<sup>12</sup> is absent in the  $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A}-\alpha 2$  spectrum. Therefore, the hydrogen bonding environment of  $\text{NH}_2\text{Y}_{731}^*/\text{R}_{411}\text{A}-\alpha 2$  is distinct from that of the single mutant, consistent with the different side chain conformations observed by HF EPR spectroscopy. Note that almost the complete EPR line of  $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A}-\alpha 2$  can be excited at 34 GHz by using very short microwave pulses,






Fig. 3 34 GHz  $^2\text{H}$  ENDOR spectra of  $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A-}\alpha 2$  (solid black line) and  $\text{ND}_2\text{Y}_{731}^*-\alpha 2$  (solid red line). Simulations are shown with dashed lines. The hydrogen bond peaks, "H-bond", assigned to a deuteron located between  $\text{Y}_{730}-\alpha 2$  and  $\text{ND}_2\text{Y}_{731}^*-\alpha 2$  (ref. 12), are not observed for  $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A-}\alpha 2$ . The inset shows the structured and broad matrix line of  $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A-}\alpha 2$ . Exp. conditions: Mims ENDOR with  $\pi/2 = 6$  ns,  $\tau = 200$  ns and 320 ns for the inset, shot repetition time = 15 ms, random RF acquisition<sup>60</sup> with 1 shot/point, acquisition time = 15–20 h,  $T = 70$  K. Excitation in the EPR line is set to  $B_0 \parallel g_y$ . ENDOR spectra are centered at the Larmor frequency of  $^2\text{H}$ ,  $\nu_0 = 7.9$  MHz at 1.2 T.

and thus hf couplings cannot be missed due to orientation selective effects.

Although no exchangeable moderately strong hydrogen bonds ( $r_{\text{O-H}} \sim 1.7\text{--}2$  Å) to  $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A-}\alpha 2$  are observed, the ENDOR spectrum of  $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A-}\alpha 2$  exhibits a broad and structured matrix line, which is associated with weak hf interactions of the radical with distant nuclei<sup>61</sup> (see Fig. 3, inset). The structure in this matrix line suggests the presence of weakly coupled deuterons that cannot be resolved from the matrix ones (matrix line). We note that the ENDOR spectrum of  $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A-}\alpha 2$  is reminiscent of the one previously observed for  $\text{ND}_2\text{Y}_{356}^*-\beta 2$ , also located at the subunit interface and likely surrounded by a defined hydrogen bonded network of water molecules.<sup>12</sup> The similarity between the ENDOR spectra of  $\text{ND}_2\text{Y}_{356}^*-\beta 2$  and  $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A-}\alpha 2$  suggests a similar origin for the  $g_x$  values in these two mutants, which is distinct from that in  $\text{ND}_2\text{Y}_{731}^*-\alpha 2$ . As noted above, in the case of  $\text{ND}_2\text{Y}_{356}^*-\beta 2$  the  $g_x$  value was also strongly shifted ( $\text{NH}_2\text{Y}_{356}^*$ :  $g_x = 2.0049$  vs. free  $\text{NH}_2\text{Y}^*$ :  $g_x = 2.0061$  (ref. 33)). Therefore, we propose that the  $g_x$ -shift in  $\text{NH}_2\text{Y}_{731}^*/\text{R}_{411}\text{A-}\alpha 2$ , as well as in  $\text{ND}_2\text{Y}_{356}^*-\beta 2$ , arises from weakly coupled hydrogen bonds observed in the 0.3 MHz region of the ENDOR spectrum. The complexity of the  $g$  tensor interpretation was underlined by our recent DFT calculations, in which three distinct models for  $\text{NH}_2\text{Y}_{731}^*-\alpha 2$  resulted in similar  $g$ -shifts.<sup>12</sup> Overall, these data clearly indicate that the molecular orientation of  $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A-}\alpha 2$  is different to that of  $\text{ND}_2\text{Y}_{731}^*-\alpha 2$  and is affected by  $\text{R}_{411}\text{A-}\alpha 2$  substitution.

### PELDOR gives evidence for a conformational change in $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A-}\alpha 2$

Our previous PELDOR studies<sup>26</sup> have demonstrated that half sites reactivity of *E. coli* RNR allows for the detection of the diagonal inter-spin distance between  $\text{Y}_{122}^*$  in one  $\alpha\beta$  pair and any radical trapped in the second  $\alpha\beta$  pair (Fig. 4A).<sup>25,62</sup> To gain insight into the location of  $\text{NH}_2\text{Y}_{731}^*/\text{R}_{411}\text{A-}\alpha 2$ , three sets of PELDOR experiments were recorded using broadband excitation with a high-power Q-band set up at different excitation positions in the EPR line<sup>63–66</sup> (see Fig. 4B and S8†). The recorded time traces are displayed in Fig. 4C and show substantial differences in modulation depth (10 to 50%), which is typical for orientation selection effects. Trace D1 also shows a higher frequency component that arises from the parallel component of a dipolar Pake pattern (Fig. S8†). For this reason, the background corrected PELDOR time traces from the three sets of experiments were summed and the resulting trace was analyzed as shown in Fig. 4C and D. Additional comparison of the Fourier-transformed traces (Fig. S8†) shows that the sum trace leads to an almost complete Pake pattern. Distance distribution analysis revealed a clear dominant peak at 35 Å with a distance distribution of  $\Delta r = \pm 2.7$  Å. We note that the error in the peak distance is much less than the distribution and is estimated to be  $\leq \pm 0.5$  Å. The width of the distance distribution is slightly larger than in previous measurements within the *E. coli* RNR  $\alpha 2\beta 2$  complex,<sup>25,26,62</sup> suggesting more conformational heterogeneity for  $\text{ND}_2\text{Y}_{731}^*/\text{R}_{411}\text{A-}\alpha 2$ , consistent with the observed flexibility of this residue. Nevertheless, the results clearly indicate that the  $\text{R}_{411}$  mutation induces a conformational change of  $\text{ND}_2\text{Y}_{731}^*$  into a new well-defined conformation.

The peak distance of 35.0 Å has never been observed between any radicals formed in this pathway before, and it is 3 Å shorter than that previously measured for  $\text{ND}_2\text{Y}_{731}^*-\alpha 2$ .<sup>26</sup> This distance might appear to be rather close to the initial distance (prior to turnover) between the two stable  $\text{Y}_{122}^*$ s, that is  $33.1 \pm 0.2$  Å.<sup>62</sup> To confirm our assignment, we recorded PELDOR experiments at higher temperature (50 K), in which the  $\text{Y}_{122}^*-\beta 2$  contribution to the re-focused echo is filtered and  $\text{ND}_2\text{Y}_{731}^*-\alpha 2$  is the only radical species detected (Fig. S9†). However,  $\text{Y}_{122}^*-\beta 2$  can still be excited by the pump pulse and contributes to the PELDOR signal. Under these conditions, any distance observed in the PELDOR experiments at 50 K is related to  $\text{Y}_{122}^*-\text{ND}_2\text{Y}_{731}^*$  and cannot be associated with the  $\text{Y}_{122}^*-\text{Y}_{122}^*$  distance, as the latter radical is not detected. The distance distribution analysis of the 50 K measurements yielded a peak distance of 35.3 Å with a distribution of  $\Delta r = \pm 2.0$  Å, and thus validated our assignment (see Fig. S9†).

To gain more insight into the conformation of  $\text{NH}_2\text{Y}_{731}^*/\text{R}_{411}\text{A-}\alpha 2$  and the role of  $\text{R}_{411}$ , we examined the available X-ray structures of *E. coli*  $\alpha 2$ s in the  $\text{R}_{411}$  region. In the structure of *E. coli*  $\text{NH}_2\text{Y}_{730}-\alpha 2$  (2XO4),<sup>30</sup>  $\text{Y}_{731}$  is flipped away from  $\text{NH}_2\text{Y}_{730}$ , as shown in Fig. 5. This altered conformation is compared with a second  $\alpha$  in the unit cell, in which the  $\text{Y}_{731}$  is not flipped. To match the 35 Å distance observed by PELDOR spectroscopy, the aromatic ring of  $\text{NH}_2\text{Y}_{731}^*$  must rotate away from  $\text{Y}_{730}$  toward the  $\beta 2$  subunit, as observed for  $\text{Y}_{731}$  in the *E. coli*  $\text{Y}_{730}\text{NH}_2\text{Y-}\alpha 2$  structure (Fig. 5).






Fig. 4 (A) Diagonal distance between  $Y_{122}^*$  in  $\beta 2$  and  $ND_2Y_{731}^*/R_{411}A-\alpha 2$ . (B) The ESE spectrum of  $ND_2Y_{731}^*/R_{411}A-\alpha 2$  is composed of the  $Y_{122}^*-\beta 2$  spectrum (dashed and dotted black line) and the  $ND_2Y_{731}^*-\alpha 2$  spectrum (dotted black line). Detect (D) and pump (P) frequency positions for each PELDOR measurement are displayed by red and black arrows, respectively. Exp. conditions (EPR):  $\pi/2 = 6$  ns,  $\tau = 280$  ns, shot repetition time = 120 ms, shots/point = 10, 4 scans,  $T = 20$  K. (C) Background- and phase-corrected, normalized 34 GHz PELDOR time traces of three experimental setups ( $D_1$ ,  $D_2$ ,  $D_3$ ). The sum of the three traces ( $D_1 + D_2 + D_3$ ) was analyzed by DeerAnalysis 2015 (ref. 67) and is shown in black with the fitting overlaid in a solid red line. Detect/pump  $\pi$  pulse lengths for  $D_1$ ,  $D_2$  and  $D_3$  were 30 ns/12 ns, 30 ns/12 ns and 20 ns/14 ns, respectively. The frequency separation between detect and pump pulses was 80 MHz for all data sets. (D) Distance distribution obtained from the analysis in (C). Asterisks indicate artifacts attributed to the analysis procedure.




Fig. 5 The *E. coli*  $Y_{730}NH_2Y-\alpha 2$  structure (2XO4)<sup>30</sup> in green shows the reoriented  $Y_{731}$  overlaid with the stacked  $Y_{731}$  in a different monomer (blue) of the unit cell. The diagonal distances between the “flipped” and non-flipped  $Y_{731}$  and  $Y_{122}$  are 3.5 nm and 3.9 nm, respectively. These distances, which are between two phenolic oxygen atoms of the tyrosine residues, are based on the alignment with the *E. coli*  $\alpha 2\beta 2$  docking model. Residue  $Y_{356}$  is shown in grey with a “?” because its position is unknown.

This reorientation is also supported by the ENDOR data, which indicate that the stacked conformation between  $NH_2Y_{731}^*$  and  $Y_{730}$  with a shared, perpendicular hydrogen bond

is absent in  $NH_2Y_{731}^*/R_{411}A-\alpha 2$ , and that the radical is instead surrounded by weakly coupled hydrogen bonds, likely water molecules at the  $\alpha 2\beta 2$  subunit interface. The exposure of  $NH_2Y_{731}^*/R_{411}A-\alpha 2$  to the interface and the buffer in this new conformation might be the origin of the instability of the radical as compared to the single mutant (Table S1†).

We have also examined another possible conformation, in which the amino group of  $NH_2Y_{731}^*-\alpha 2$  moves to occupy the vacancy created by the mutation of arginine to alanine. This conformation is displayed in Fig. S10.† However, in this case the expected distance between the oxygen atoms of  $NH_2Y_{731}^*$  and  $Y_{122}$  exceeds the observed distance by  $\geq 2$  Å. We note that the “flipped” conformation has not been observed in the single mutant  $NH_2Y_{731}^*-\alpha 2$  or in the double mutant  $NH_2Y_{731}^*/Y_{730}F-\alpha 2$ , in which  $Y_{731}$  lacks its hydrogen bonding partner,<sup>12</sup> suggesting the importance of  $R_{411}$  in stabilizing the stacked conformation. This change between a flipped and non-flipped conformation of the interface Y might play an active role in the PCET process between  $Y_{731}$  and  $Y_{356}$  in wt RNR, the mechanism of which is still not understood. With the wt enzyme, this conformational change is kinetically masked by physical gating, which rate-limits RNR, and is too fast to be detected based on the recently



measured rate constants for electron transfer (ET) ( $10^4$  to  $10^5$  s $^{-1}$ ) at the interface by photo-RNRs that unmask this gating.<sup>68,69</sup> Thus, the R<sub>411</sub>A mutation might have fortuitously allowed detection of this movement at the subunit interface.

While the lack of structural information at the subunit interface poses a challenge for a mechanistic understanding of interfacial PCET, the detection of the NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub> provides us with a spectroscopic probe of this interface. Mutagenesis and site-specific isotopic labeling of interface residues could provide us with additional insight into how this step is controlled. Finally, the mechanism of PCET across the subunit interface observed with the *E. coli* RNR is likely to be conserved in all class I RNRs based on their subunit structures and the conserved weak subunit associations dictated by the C-terminal tail of  $\beta 2$ .<sup>70,71</sup> The pathway for oxidation is conserved between RNR classes Ia, Ib and Ic, as is the regulation of the pathway by NDP/effector binding.<sup>72</sup> Thus, while the “details” of the radical transfer mechanism might be different in the individual class I RNRs, general principles will likely emerge from the studies on *E. coli* RNR, given all of the evolutionarily conserved features.

## Conclusions

This study has revealed that the *E. coli* RNR double mutant NH<sub>2</sub>Y<sub>731</sub>/R<sub>411</sub>A- $\alpha 2$  unmasks a new conformation of pathway residue 731 in the  $\alpha 2\beta 2$  complex. This is the first experimental evidence for the flexibility of this pathway or any pathway residue in the active enzyme. The results have provided insight into the mechanisms of PCET within  $\alpha 2$ , as well as through the  $\alpha 2\beta 2$  interface. First, R<sub>411</sub> appears to play a role in the stabilization of the stacked conformation of Y<sub>731</sub> and Y<sub>730</sub>, and thus in the facilitation of collinear PCET within the  $\alpha 2$  subunit. Second, the new conformation is consistent with Y<sub>731</sub> pointing toward the subunit interface, in the direction of the adjacent pathway residue Y<sub>356</sub>, located in the flexible C-terminal tail of subunit  $\beta 2$ . The flexibility of these two contiguous pathway residues, which have been suggested to communicate during PCET,<sup>69</sup> might be the key to driving the RT chemistry at the subunit interface through water clusters.<sup>67</sup> This opens up a new hypothesis for the PCET mechanism between residues Y<sub>731</sub>- $\alpha 2$  and Y<sub>356</sub>- $\beta 2$ , which could involve a gated conformational change in Y<sub>731</sub>- $\alpha 2$  in wt RNR on a fast time scale, not observable without the R<sub>411</sub>A mutation. While this hypothesis remains to be proven, the present results will serve as a basis to design new experiments aimed at detecting a possible combined role of Y<sub>731</sub>- $\alpha 2$  and Y<sub>356</sub>- $\beta 2$  in PCET through the subunit surface.

## Acknowledgements

We acknowledge Igor Tkach for the help with technical aspects of the HF EPR spectrometers. MK thanks Karin Halbmaier for the assistance with PELDOR measurements. We gratefully acknowledge financial support for this work from Deutsche Forschungsgemeinschaft DFG-IRTG 1422 (GRK 1422 to MK and MB) and DFG-SPP 1601, the Max Planck Society and NIH (GM29595 to JS).

## References

- 1 A. Migliore, N. F. Polizzi, M. J. Therien and D. N. Beratan, *Chem. Rev.*, 2014, **114**, 3381–3465.
- 2 D. N. Beratan, C. Liu, A. Migliore, N. F. Polizzi, S. S. Skourtis, P. Zhang and Y. Zhang, *Acc. Chem. Res.*, 2015, **48**, 474–481.
- 3 D. R. Weinberg, C. J. Gagliardi, J. F. Hull, C. F. Murphy, C. A. Kent, B. C. Westlake, A. Paul, D. H. Ess, D. G. McCafferty and T. J. Meyer, *Chem. Rev.*, 2012, **112**, 4016–4093.
- 4 S. Y. Reece, J. M. Hodgkiss, J. Stubbe and D. G. Nocera, *Philos. Trans. R. Soc., B*, 2006, **361**, 1351–1364.
- 5 S. Y. Reece and D. G. Nocera, *Annu. Rev. Biochem.*, 2009, **78**, 673–699.
- 6 J. Bonin, C. Costentin, C. Louault, M. Robert and J. M. Savéant, *J. Am. Chem. Soc.*, 2011, **133**, 6668–6674.
- 7 J. M. Savéant, *Annu. Rev. Anal. Chem.*, 2014, **7**, 537–560.
- 8 R. I. Cukier and D. G. Nocera, *Annu. Rev. Phys. Chem.*, 1998, **49**, 337–369.
- 9 J. M. Mayer, *Annu. Rev. Phys. Chem.*, 2004, **55**, 363–390.
- 10 A. Jordan and P. Reichard, *Annu. Rev. Biochem.*, 1998, **67**, 71–98.
- 11 J. Stubbe and W. A. van der Donk, *Chem. Rev.*, 1998, **98**, 705–762.
- 12 T. U. Nick, W. Lee, S. Kofmann, F. Neese, J. Stubbe and M. Bennati, *J. Am. Chem. Soc.*, 2015, **137**, 289–298.
- 13 L. Thelander, *J. Biol. Chem.*, 1973, **248**, 4591–4601.
- 14 E. C. Minnihan, D. G. Nocera and J. Stubbe, *Acc. Chem. Res.*, 2013, **46**, 2524–2535.
- 15 N. C. Brown and P. Reichard, *J. Mol. Biol.*, 1969, **46**, 25–38.
- 16 N. C. Brown and P. Reichard, *J. Mol. Biol.*, 1969, **46**, 39–55.
- 17 A. Hofer, M. Crona, D. T. Logan and B. M. Sjöberg, *Crit. Rev. Biochem. Mol. Biol.*, 2012, **47**, 50–63.
- 18 J. A. Stubbe, *J. Biol. Chem.*, 1990, **265**, 5329–5332.
- 19 M. Eriksson, U. Uhlin, S. Ramaswamy, M. Ekberg, K. Regnström, B. M. Sjöberg and H. Eklund, *Structure*, 1997, **5**, 1077–1092.
- 20 A. Ehrenberg and P. Reichard, *J. Biol. Chem.*, 1972, **247**, 3485–3488.
- 21 B. M. Sjöberg, P. Reichard, A. Graslund and A. Ehrenberg, *J. Biol. Chem.*, 1978, **253**, 6863–6865.
- 22 P. Nordlund, B. M. Sjöberg and H. Eklund, *Nature*, 1990, **345**, 593–598.
- 23 U. Uhlin and H. Eklund, *Nature*, 1994, **370**, 533–539.
- 24 J. A. Stubbe, D. G. Nocera, C. S. Yee and M. C. Y. Chang, *Chem. Rev.*, 2003, **103**, 2167–2201.
- 25 M. Bennati, J. H. Robblee, V. Mugnaini, J. Stubbe, J. H. Freed and P. Borbat, *J. Am. Chem. Soc.*, 2005, **127**, 15014–15015.
- 26 M. R. Seyedsayamdost, C. T. Y. Chan, V. Mugnaini, J. Stubbe and M. Bennati, *J. Am. Chem. Soc.*, 2007, **129**, 15748–15749.
- 27 N. Ando, E. J. Brignole, C. M. Zimanyi, M. A. Funk, K. Yokoyama, F. J. Asturias, J. Stubbe and C. L. Drennan, *Proc. Natl. Acad. Sci. U. S. A.*, 2011, **108**, 21046–21051.
- 28 E. C. Minnihan, N. Ando, E. J. Brignole, L. Olshansky, J. Chittuluru, F. J. Asturias, C. L. Drennan, D. G. Nocera



- and J. Stubbe, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, 3835–3840.
- 29 J. Ge, G. Yu, M. A. Ator and J. Stubbe, *Biochemistry*, 2003, **42**, 10071–10083.
- 30 E. C. Minnihhan, M. R. Seyedsayamdost, U. Uhlin and J. Stubbe, *J. Am. Chem. Soc.*, 2011, **133**, 9430–9440.
- 31 K. R. Ravichandran, E. C. Minnihhan, Y. Wei, D. G. Nocera and J. Stubbe, *J. Am. Chem. Soc.*, 2015, **137**, 14387–14395.
- 32 M. R. Seyedsayamdost, J. Xie, C. T. Y. Chan, P. G. Schultz and J. Stubbe, *J. Am. Chem. Soc.*, 2007, **129**, 15060–15071.
- 33 T. Argirević, C. Riplinger, J. Stubbe, F. Neese and M. Bennati, *J. Am. Chem. Soc.*, 2012, **134**, 17661–17670.
- 34 M. R. Seyedsayamdost, T. Argirević, E. C. Minnihhan, J. Stubbe and M. Bennati, *J. Am. Chem. Soc.*, 2009, **131**, 15729–15738.
- 35 B. Wörsdörfer, D. A. Conner, K. Yokoyama, J. Livada, M. Seyedsayamdost, W. Jiang, A. Silakov, J. Stubbe, J. M. Bollinger and C. Krebs, *J. Am. Chem. Soc.*, 2013, **135**, 8585–8593.
- 36 K. Yokoyama, U. Uhlin and J. Stubbe, *J. Am. Chem. Soc.*, 2010, **132**, 8385–8397.
- 37 S. P. Salowe, M. A. Ator and J. Stubbe, *Biochemistry*, 1987, **26**, 3408–3416.
- 38 S. P. Salowe and J. Stubbe, *J. Bacteriol.*, 1986, **165**, 363–366.
- 39 P. T. Chivers, K. E. Prehoda, B. F. Volkman, B. M. Kim, J. L. Markley and R. T. Raines, *Biochemistry*, 1997, **36**, 14985–14991.
- 40 M. Russel and P. Model, *J. Bacteriol.*, 1985, **163**, 238–242.
- 41 J. R. Steeper and C. D. Steuart, *Anal. Biochem.*, 1970, **34**, 123–130.
- 42 S. Climent, B. M. Sjöberg and C. Y. Huang, *Biochemistry*, 1991, **30**, 5164–5171.
- 43 A. Savitzky and M. J. E. Golay, *Anal. Chem.*, 1964, **36**, 1627–1639.
- 44 S. Stoll and A. Schweiger, *J. Magn. Reson.*, 2006, **178**, 42–55.
- 45 F. Neese, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2012, **2**, 73–78.
- 46 A. D. Becke, *Phys. Rev. A: At., Mol., Opt. Phys.*, 1988, **38**, 3098–3100.
- 47 A. D. Becke, *J. Chem. Phys.*, 1993, **98**, 5648–5652.
- 48 C. Lee, W. Yang and R. G. Parr, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 1988, **37**, 785–789.
- 49 A. Schäfer, C. Huber and R. Ahlrichs, *J. Chem. Phys.*, 1994, **100**, 5829–5835.
- 50 F. Weigend and R. Ahlrichs, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297–3305.
- 51 A. Klamt and G. Schüürmann, *J. Chem. Soc., Perkin Trans. 2*, 1993, 799–805.
- 52 S. Grimme, J. Antony, S. Ehrlich and H. Krieg, *J. Chem. Phys.*, 2010, **132**, 154104.
- 53 S. Grimme, S. Ehrlich and L. Goerigk, *J. Comput. Chem.*, 2011, **32**, 1456–1465.
- 54 F. Neese, F. Wennmohs, A. Hansen and U. Becker, *Chem. Phys.*, 2009, **356**, 98–109.
- 55 S. Kossmann, B. Kirchner and F. Neese, *Mol. Phys.*, 2007, **105**, 2049–2071.
- 56 F. Neese, in *Multifrequency Electron Paramagnetic Resonance*, ed. S. K. Misra, Wiley-VCH Verlag GmbH & Co. KGaA, 2011, ch. 6, pp. 295–326.
- 57 S. Un, M. Atta, M. Fontecave and A. W. Rutherford, *J. Am. Chem. Soc.*, 1995, **117**, 10713–10719.
- 58 H. M. McConnell, *J. Chem. Phys.*, 1956, **24**, 764–766.
- 59 R. W. Fessenden and R. H. Schuler, *J. Chem. Phys.*, 1963, **39**, 2147–2195.
- 60 B. Epel, D. Arieli, D. Baute and D. Goldfarb, *J. Magn. Reson.*, 2003, **164**, 78–83.
- 61 A. V. Astashkin and A. Kawamori, *J. Magn. Reson.*, 1998, **135**, 406–417.
- 62 M. Bennati, A. Weber, J. Antonic, D. L. Perlstein, J. Robblee and J. Stubbe, *J. Am. Chem. Soc.*, 2003, **125**, 14988–14989.
- 63 V. P. Denysenkov, T. F. Prisner, J. Stubbe and M. Bennati, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 13386–13390.
- 64 A. D. Milov, A. G. Maryasov and Y. D. Tsvetkov, *Appl. Magn. Reson.*, 1998, **15**, 107–143.
- 65 Y. Polyhach, A. Godt, C. Bauer and G. Jeschke, *J. Magn. Reson.*, 2007, **185**, 118–129.
- 66 G. Sicoli, T. Argirević, J. Stubbe, I. Tkach and M. Bennati, *Appl. Magn. Reson.*, 2010, **37**, 539–548.
- 67 G. Jeschke, V. Chechik, P. Ionita, A. Godt, H. Zimmermann, J. Banham, C. R. Timmel, D. Hilger and H. Jung, *Appl. Magn. Reson.*, 2006, **30**, 473–498.
- 68 L. Olshansky, A. A. Pizano, Y. Wei, J. Stubbe and D. G. Nocera, *J. Am. Chem. Soc.*, 2014, **136**, 16210–16216.
- 69 D. Y. Song, A. A. Pizano, P. G. Holder, J. Stubbe and D. G. Nocera, *Chem. Sci.*, 2015, **6**, 4519–4524.
- 70 M. Kolberg, K. R. Strand, P. Graff and K. K. Andersson, *Biochim. Biophys. Acta*, 2004, **1699**, 1–34.
- 71 Y. Zhang, X. An, J. Stubbe and M. Huang, *J. Biol. Chem.*, 2013, **288**, 13951–13959.
- 72 P. Nordlund and P. Reichard, *Annu. Rev. Biochem.*, 2006, **75**, 681–706.

