Cite this: *Chem. Sci.*, 2016, 7, 1051

A highly selective near-infrared fluorescent probe for imaging H₂Se in living cells and *in vivo*†

Fanpeng Kong, Lihong Ge, Xiaohong Pan, Kehua Xu,* Xiaojun Liu and Bo Tang*

Hydrogen selenide (H₂Se), a highly reactive Se species, is an important selenium metabolism intermediate involved in many physiological and pathological processes. This compound is of scientific interest with regard to the real-time monitoring of H₂Se in living cells and *in vivo* to understand the anti-cancer mechanism of selenium. However, monitoring H₂Se in living cells is still challenging due to the lack of straight forward, highly selective and rapid methods. Here, we developed a novel small-molecule fluorescent probe, **NIR-H₂Se**, for imaging endogenous H₂Se. **NIR-H₂Se** exhibited high selectivity toward H₂Se over selenocysteine (Sec), H₂S and small molecule thiols and was successfully used to image the H₂Se content in HepG2 cells during Na₂SeO₃-induced apoptosis. Increased H₂Se content and reduced ROS levels were observed under hypoxic conditions compared to normoxic conditions, which indicated that the cell apoptosis induced by Na₂SeO₃ under a hypoxic environment is *via* a non-oxidative stress mechanism. Thus, this probe should serve as a powerful tool for exploring the physiological function of H₂Se and Se anticancer mechanisms in a variety of physiological and pathological contexts.

Received 15th September 2015

Accepted 28th October 2015

DOI: 10.1039/c5sc03471j

www.rsc.org/chemicalscience

Introduction

Selenium (Se) is an essential trace element for various physiological functions in the human body. Insufficient or excessive Se intake has been associated with a number of diseases.^{1,2} Importantly, selenium plays a role in cancer prevention and treatment.³ However, the anticancer mechanism for Se is still not fully understood. The effectiveness of Se compounds as anticancer agents is correlated to their chemical form and dose.⁴ H₂Se, a reduced form of selenium, is an important metabolite of dietary Se compounds.⁵ Endogenous H₂Se is generated by reducing selenite *via* GSH and other reduction systems⁶ and is involved in many physiological and pathological processes.⁷ A reliable and rapid method to determine H₂Se *in vivo* must be developed to investigate the physiological function of H₂Se and the anticancer mechanism of Se.

Fluorescent probes with high sensitivity, good selectivity, and short response times *via* direct observation are powerful tools for the *in vivo* detection of biomolecules.⁸ Several fluorescent probes for detecting Se species were recently reported. Maeda *et al.* developed the first fluorescent probe, BESThio, to

discriminate selenols from their Cys counterparts based on their different nucleophilicity at pH = 5.8.⁹ Using the recognition group Maeda reported, Fang and Lin exploited probes by adjusting the fluorophore to detect selenols with high selectivity under physiological conditions.¹⁰ In addition, Fang *et al.* developed another novel fluorescent probe (TRFS-green) using a 1,2-dithiolane reporter group to selectively image thioredoxin reductase (TrxR) at pH = 7.4.¹¹ Wu reported a cadmium sulfide (CdS) quantum dot (QD) probe for HSe[−] ions in aqueous solution.¹² However, a small-molecule fluorescent probe for imaging the highly reactive H₂Se in living cells and *in vivo* has not been reported.

Previously, our group discovered that the Se–N bond can be cleaved using sulfhydryl groups and we developed a series of fluorescence methods to monitor thiols and their redox state in living cells.^{13–15} We subsequently found that the Se–N bond in 2,1,3-benzoselenadiazole (BS) can be specifically broken by selenol *via* direct nucleophilic addition and we designed a new fluorescent probe (HB) to image endogenous Sec in living cells and *in vivo*.¹⁶ These results inspired us to develop a H₂Se fluorescent probe based on the BS group by modifying the probe molecule structure.

Here, we integrated the BS moiety into a near infrared (NIR) merocyanine dye¹⁷ to develop a novel small-molecule fluorescent probe for detecting H₂Se (**NIR-H₂Se**). **NIR-H₂Se** responds to H₂Se rapidly, with a high selectivity over H₂S, Sec and biological thiols and was used to successfully image endogenous H₂Se in living cells and *in vivo*. Moreover, we used a radical oxygen species (ROS) probe and found that the H₂Se contents increased during HepG2 cell apoptosis induced by Na₂SeO₃ in a time- and

College of Chemistry, Chemical Engineering and Materials Science, Collaborative Innovation Center of Functionalized Probes for Chemical Imaging in Universities of Shandong, Key Laboratory of Molecular and Nano Probes, Ministry of Education, Shandong Provincial Key Laboratory of Clean Production of Fine Chemicals, Shandong Normal University, Jinan 250014, P. R. China. E-mail: tangb@sdnu.edu.cn; xukehua@sdnu.edu.cn

† Electronic supplementary information (ESI) available: The NMR and HR MS spectra of **NIR-H₂Se**, HPLC analysis of the reaction of the probe and H₂Se, and other materials. See DOI: 10.1039/c5sc03471j

Results and discussion

To evaluate the selectivity of **NIR-H₂Se** for H₂Se, we measured the fluorescence spectra for the probe with H₂Se, H₂S and thiols. High concentrations of thiols such as cysteine (Cys), homocysteine (Hcy) and glutathione (GSH) (1 mM for each) produced low fluorescence responses, and **NIR-H₂Se** also gave a negative response for H₂S, Sec, Na₂SeO₃, *N*-acetyl-L-cysteine (NAC), thioredoxin reductase (TrxR) and vitamin C (Vc). The reactivity of **NIR-H₂Se** towards ROS was then tested. Fig. 2a shows that biologically relevant ROS, including H₂O₂, ⁻OCl, O₂^{-•}, and ROO[•], did not trigger any fluorescence changes in the probe. Moreover, considering that nitric oxide (NO) may be a potential interfering species,¹⁸ the fluorescence response of **NIR-H₂Se** to differing amounts of NO were studied both with and without H₂Se. The results indicated that even high NO concentrations (10 equiv.) did not induce an obvious fluorescence change in the detection system. The interference from

We found that the apparent discrepancies above were related to the aerobic metabolism of H_2Se . Under normoxic conditions (20% pO_2), H_2Se rapidly oxidized to generate a mass of superoxide anion radicals ($\text{O}_2^{\cdot-}$) and an accumulation of reactive

(a) Selenite 12 h

	Control	2 μM	5 μM	10 μM
20% pO_2	[Image]	[Image]	[Image]	[Image]
1% pO_2	[Image]	[Image]	[Image]	[Image]

(b) Selenite 5 μM

	Control	2 h	6 h	12 h
20% pO_2	[Image]	[Image]	[Image]	[Image]
1% pO_2	[Image]	[Image]	[Image]	[Image]

(c)

Na_2SeO_3 Concentration (μM)	20% pO_2	1% pO_2
Control	1.0	1.0
2	1.5	1.6
5	1.8	2.7
10	2.3	3.4

(d)

Time (h)	20% pO_2	1% pO_2
Control	1.0	1.0
2	1.5	1.6
6	1.8	2.5
12	2.4	3.5

oxygen species during the tumor apoptosis process induced by Na_2SeO_3 .²⁰ However, under a hypoxic environment (1% pO_2), H_2Se cannot quickly metabolize due to the lack of oxygen, which allows H_2Se to effectively accumulate. To further confirm our

The above results indicate that **NIR-H₂Se** has potential for detecting endogenous H₂Se *in vivo*. To evaluate this proposal, mice bearing subcutaneously implanted tumors grown from murine hepatoma cell line H22 were subcutaneously injected with buffer solutions containing **NIR-H₂Se** (10 μM) and sodium selenite (10 μM), and fluorescence images were then obtained at

Chem. Sci., 2016, 7, 1051–1056 | 1053

different times using an *in vivo* imaging system (IVIS). Fig. 5 shows that the fluorescence signal of the probe was exclusively observed in the tumor region without a background signal, and that the fluorescence intensity increased from 3 h to 12 h post injection. The results indicated that the hypoxic solid tumor generated H_2Se from sodium selenite and that the H_2Se gradually accumulated.

Conclusions

In summary, we developed a novel small-molecule fluorescent probe, **NIR- H_2Se** , to detect H_2Se . The probe rapidly responded to H_2Se , exhibited high sensitivity and good selectivity for H_2Se over Sec , H_2S , ROS and other sulfur-containing species and was successfully used to image endogenous H_2Se in living cells and *in vivo*. Furthermore, we found that normoxic conditions increased the H_2O_2 content, while the H_2Se level rose only slightly during HepG2 cell apoptosis induced by Na_2SeO_3 . However, H_2Se accumulated gradually and the ROS remained low in a hypoxic environment. These findings indicate that the anticancer mechanism of Se for hypoxic solid tumors is *via* non-oxidative stress. We anticipate that the current probe will provide an ideal tool for further studies into the biological functions of H_2Se and Se anticancer mechanisms.

Experimental

Materials and instruments

All chemicals were available commercially and the solvents were purified by conventional methods before use. Cysteine (Cys), homocysteine (Hcy), glutathione (GSH) and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) were purchased from Sigma Chemical Company. Na_2SeO_3 , *N*-acetyl-L-cysteine (NAC), thioredoxin reductase (TrxR) and vitamin C (Vc) were purchased from Sigma-Aldrich Co. Ltd. The silica gel (100–200 mesh) was used for the flash chromatography. Cy7.Cl was synthesized in our laboratory. Sartorius ultrapure water (18.2 M Ω cm) was used throughout the analytical experiments. H_2Se was prepared by the reaction of Al_2Se_3 with H_2O in an N_2 atmosphere for 30 min at room temperature before use every time.²⁴ H_2O_2 , *tert*-butylhydroperoxide (TBHP), and hypochlorite (NaOCl) were delivered from 30%, 70%, and

10% aqueous solutions respectively. Nitric oxide (NO) was used from a stock solution prepared by sodium nitroprusside. Superoxide ($\text{O}_2^{\cdot-}$) was delivered from KO_2 in DMSO solution or from xanthine oxidase.

^1H NMR and ^{13}C NMR spectra were determined using 300 MHz and 400 MHz Bruker NMR spectrometers. The mass spectra were obtained using a Bruker maXis ultra-high resolution-TOF MS system. The melting points were measured using a SGW X-4 Melting Point Tester. The fluorescence spectra measurements were performed using an FLS-920 Fluorescence Spectrometer (Edinburgh Instruments, UK). All pH measurements were performed with a pH-3c digital pH-meter (Shanghai Lei Ci Device Works, Shanghai, China) with a combined glass/calomel electrode. The fluorescence images of cells were taken using a TCS SP5 confocal laser scanning microscope with an objective lens ($\times 40$). The absorbance was measured using a TRITURUS microplate reader in the MTT assay.

Synthesis and characterization of NIR- H_2Se

Synthesis of compound 1. 4-Amino-3-nitrophenol (39 mg, 0.25 mmol) and NaH (60% in mineral oil) (10 mg, 0.25 mmol) were dissolved in anhydrous *N,N*-dimethylformamide (DMF) (7 mL). The mixture was stirred at room temperature for 10 min under an argon atmosphere. Then a solution of heptamethine cyanine chlorine (Cy7.Cl) (64 mg, 0.10 mmol) in anhydrous DMF (2 mL) was added to the mixture *via* a syringe. The reaction mixture was further stirred for 4 h at room temperature. The solvent was removed under reduced pressure, then the crude product was purified by silica gel chromatography with 6% MeOH in dichloromethane (DCM) to afford the desired product as a dark green solid (55 mg, 73%). mp = 149–150 °C. ^1H NMR (400 MHz, CDCl_3): δ 1.26–1.40 (m, 18H), 2.03 (s, 4H), 2.17 (s, 2H), 2.68 (s, 4H), 4.08 (s, 4H), 5.97 (d, J = 12 Hz, 2H), 7.09 (d, J = 8 Hz, 2H), 7.21–7.29 (m, 5H), 7.54 (s, 3H), 7.80 (d, J = 8 Hz, 1H), 7.92 (d, J = 12 Hz, 2H). ^{13}C NMR (75 MHz, CDCl_3): δ 12.2, 21.0, 24.3, 27.8, 39.4, 49.1, 99.3, 108.3, 110.2, 122.1, 122.4, 125.3, 128.7, 141.1, 141.5, 142.2, 143.4, 164.4, 171.7. HR MS [$\text{M} - \text{I}$] $^+$: m/z calcd 629.3486, found 629.3455.

Synthesis of compound 2. SnCl_2 (900 mg, 4 mmol) and concentrated HCl (0.8 mL) were added to a solution of compound 1 (151 mg, 0.20 mmol) in MeOH (6 mL), stirred, and maintained at 70 °C overnight under an argon atmosphere. The mixture was neutralized with 2 N NaOH, and the precipitate was removed by filtration and washed with DCM. The filtrate and washings were washed with water. The organic layer was dried over Na_2SO_4 , and the solvent was evaporated to yield a dark green solid (49 mg, 45%). mp = 225–228 °C. ^1H NMR (400 MHz, CDCl_3): δ 1.31 (s, 3H), 1.69 (s, 6H), 1.84 (s, 2H), 2.57 (s, 2H), 2.70 (s, 2H), 3.89 (s, 2H), 4.63 (s, 1H), 5.68 (s, 1H), 6.75 (s, 1H), 6.86–6.92 (m, 2H), 7.03 (s, 1H), 7.28 (s, 2H), 7.39 (s, 2H), 8.10 (d, J = 12 Hz, 1H). ^{13}C NMR (75 MHz, CDCl_3): δ 11.6, 21.0, 24.5, 28.3, 28.7, 48.0, 97.5, 107.2, 108.4, 115.7, 121.5, 128.2, 137.1, 139.3, 148.5, 152.6, 160.8, 166.8. HR MS [$\text{M} - \text{I}$] $^+$: m/z calcd 412.2383, found 412.2378.

Synthesis of NIR- H_2Se . To a porcelain mortar was added compound 2 (108 mg, 0.20 mmol) and SeO_2 (22 mg, 0.2 mmol).

Fig. 5 (a) *In vivo* fluorescence imaging of H22 tumor-bearing mice injected with the probe NIR- H_2Se (10 μM) in response to sodium selenite (10 μM) at different times. (b) The fluorescence intensity for (a). The data were normalized to the control.

The mixture was fully grinded for 30 min, and TLC showed full conversion of compound 2 to the probe. The resulting mixture was extracted by MeOH (20 mL) and the solvent was removed under reduced pressure, then the crude product was purified by silica gel chromatography with 6% MeOH in DCM to afford the desired product as a dark purple solid (87 mg, 71%). mp = 252–253 °C. ^1H NMR (400 MHz, CDCl_3): δ 1.61 (t, J = 6 Hz, 3H), 1.86 (s, 6H), 2.01 (t, J = 3 Hz, 2H), 2.41 (s, 2H), 2.75 (t, J = 3 Hz, 2H), 3.00 (t, J = 3 Hz, 2H), 5.00 (q, J = 6 Hz, 2H), 7.38 (d, J = 16 Hz, 1H), 7.49 (s, 1H), 7.54–7.59 (m, 4H), 7.68 (s, 1H), 7.74 (d, J = 16 Hz, 1H). ^{13}C NMR (75 MHz, CDCl_3): δ 13.7, 14.1, 20.6, 24.5, 29.3, 30.9, 42.9, 51.7, 104.7, 110.7, 114.3, 119.0, 119.6, 122.5, 124.7, 126.5, 128.0, 128.8, 129.7, 132.6, 140.7, 142.9, 146.6, 152.7, 155.5, 179.6. HR MS: $[\text{M} - \text{I}]^+$: m/z calcd 488.1218, found 488.1237.

Fluorescence analysis

Fluorescence spectra were obtained with an FLS-920 Fluorescence Spectrometer (Edinburgh Instruments, UK). After dilution to 10 μM of the probe with 10 mM PBS, various amounts of H_2Se were added. The fluorescence intensity was measured at $\lambda_{\text{ex}}/\lambda_{\text{em}}$ = 688/735 nm.

Cell culture

HepG2 cells were maintained following the protocols provided by the American Type Tissue Culture Collection. Cells were first grown in a circular Petri dish (60 mm) using high glucose Dulbecco's Modified Eagle Medium (DMEM, 4.5 g of glucose per L) supplemented with 10% fetal bovine serum (FBS), NaHCO_3 (2 g L^{-1}) and 1% antibiotics (penicillin/streptomycin, 100 U mL^{-1}). Cultures were maintained in a humidified incubator at 37 °C, in 5% CO_2 /95% air. One day before imaging, cells were passed and plated on 18 mm glass coverslips in a culture dish. The culture medium was refreshed every 24 h. All cells used were in the exponential growth phase.

Confocal imaging

Fluorescence imaging studies were performed with a TCS SP5 confocal laser scanning microscope (Germany Leica Co., Ltd) with an objective lens ($\times 40$). Excitation of the H_2Se probe-loaded cells at 633 nm was carried out with an argon laser, and emission was collected using a META detector between 650 and 750 nm. Excitation of the H_2O_2 probe-loaded cells at 532 nm was carried out with an argon laser, and emission was collected using a META detector between 650 and 750 nm. Prior to imaging, the medium was removed. Cell imaging was carried out after washing cells with PBS (pH = 7.4, 10 mM) three times.

Animal and tumor models

All animal experiments were carried out according to the Principles of Laboratory Animal Care (People's Republic of China) and the Guidelines of the Animal Investigation Committee, and approved by the local Animal Care and Use Committee. Six- to eight-week-old female Kunmin male mice were purchased from the Shanghai SLAC Laboratory Animal Co., Ltd. During

procedures, the mice were anesthetized with inhaled isoflurane. 1×10^6 H22 cells were injected into the enterocoelia of each Kunmin mouse, ascites were formed after 5 or 7 days, which were further used after three passages. Four- to six-week-old nude mice received a subcutaneous injection of 1×10^6 H22 ascite tumor cells into the axillary lateral subcutaneous of their right forelimbs. Tumors were then allowed to grow over a period of 15 to 20 days until reaching 0.5–1.5 cm in diameter.

Acknowledgements

This work was supported by 973 Program (2013CB933800), National Natural Science Foundation of China (21535004, 21227005, 21390411, 21275092, 21575081 and 21405098).

Notes and references

- (a) C. M. Weekley and H. H. Harris, *Chem. Soc. Rev.*, 2013, **42**, 8870–8894; (b) M. P. Rayman, *Lancet*, 2012, **379**, 1256–1268.
- (a) M. A. Reeves and P. R. Hoffmann, *Cell. Mol. Life Sci.*, 2009, **66**, 2457–2478; (b) S. Misra, D. Peak, N. Chen, C. Hamilton and S. Niyogi, *Comp. Biochem. Physiol., Part C: Pharmacol., Toxicol. Endocrinol.*, 2012, **155**, 560–565; (c) S. J. Fairweather-Tait, Y. Bao, M. R. Broadley, R. Collings, D. Ford, J. E. Hesketh and R. Hurst, *Antioxid. Redox Signaling*, 2011, **14**, 1337–1383.
- (a) C. M. Weekley, J. B. Aitken, S. Vogt, L. A. Finney, D. J. Paterson, M. D. de Jonge, D. L. Howard, I. F. Musgrave and H. H. Harris, *Biochemistry*, 2011, **50**, 1641–1650; (b) A. J. Duffield-Lillico, M. E. Reid, B. W. Turnbull, G. F. Combs Jr, E. H. Slate, L. A. Fischbach, J. R. Marshall and L. C. Clark, *Cancer Epidemiol., Biomarkers Prev.*, 2002, **11**, 630–639; (c) C. M. Weekley, J. B. Aitken, S. Vogt, L. A. Finney, D. J. Paterson, M. D. de Jonge, D. L. Howard, P. K. Witting, I. F. Musgrave and H. H. Harris, *J. Am. Chem. Soc.*, 2011, **133**, 18272–18279.
- (a) C. M. Weekley and H. H. Harris, *Chem. Soc. Rev.*, 2013, **42**, 8870–8894; (b) L. C. Clark, G. F. Combs, B. W. Turnbull, E. H. Slate, D. K. Chalker, J. Chow, L. S. Davis, R. A. Glover, G. F. Graham and E. G. Gross, *J. Am. Med. Assoc.*, 1996, **276**, 1957–1963; (c) S. M. Lippman, E. A. Klein, P. J. Goodman, M. S. Lucia, I. M. Thompson, L. G. Ford, H. L. Parnes, L. M. Minasian, J. M. Gaziano and J. A. Hartline, *J. Am. Med. Assoc.*, 2009, **301**, 39–51.
- G. Combs and W. Gray, *Pharmacol. Ther.*, 1998, **79**, 179–192.
- M. Wallenberg, E. Olm, C. Hebert, M. Björnstedt and A. P. Fernandes, *Biochem. J.*, 2010, **429**, 85–93.
- (a) Z. Veres, L. Tsai, T. D. Scholz, M. Politino, R. S. Balaban and T. C. Stadtman, *Proc. Natl. Acad. Sci. U. S. A.*, 1992, **89**, 2975–2979; (b) R. S. Glass, W. P. Singh, W. Jung, Z. Veres, T. D. Scholz and T. C. Stadtman, *Biochemistry*, 1993, **32**, 12555–12559; (c) Y. Kobayashi, Y. Ogra, K. Ishiwata, H. Takayama, N. Aimi and K. T. Suzuki, *Proc. Natl. Acad. Sci. U. S. A.*, 2002, **99**, 15932–15936.
- (a) L. Yuan, W. Y. Lin, K. B. Zheng, L. W. He and W. M. Huang, *Chem. Soc. Rev.*, 2013, **42**, 622–661; (b)

- Z. Q. Guo, S. Park, J. Y. Yoon and I. Shin, *Chem. Soc. Rev.*, 2014, **43**, 16–29; (c) R. Weissleder and V. Ntziachristos, *Nat. Med.*, 2003, **9**, 123–128.
- 9 H. Maeda, K. Katayama, H. Matsuno and T. Uno, *Angew. Chem., Int. Ed.*, 2006, **45**, 1810–1813.
- 10 (a) B. Zhang, C. Ge, J. Yao, Y. Liu, H. Xie and J. Fang, *J. Am. Chem. Soc.*, 2015, **137**, 757; (b) H. Chen, B. Dong, Y. Tang and W. Lin, *Chem.-Eur. J.*, 2015, **21**, 11696–11700.
- 11 L. Zhang, D. Duan, Y. Liu, C. Ge, X. Cui, J. Sun and J. Fang, *J. Am. Chem. Soc.*, 2014, **136**, 226–233.
- 12 C. L. Wu and Y. B. Zhao, *Anal. Bioanal. Chem.*, 2007, **388**, 717–722.
- 13 B. Tang, Y. Xing, P. Li, N. Zhang, F. Yu and G. Yang, *J. Am. Chem. Soc.*, 2007, **129**, 11666–11667.
- 14 K. Xu, M. Qiang, W. Gao, R. Su, N. Li, Y. Gao, Y. Xie, F. Kong and B. Tang, *Chem. Sci.*, 2013, **4**, 1079–1086.
- 15 B. Tang, L. Yin, X. Wang, Z. Chen, L. Tong and K. Xu, *Chem. Commun.*, 2009, **45**, 5293–5295.
- 16 F. Kong, B. Hu, Y. Gao, X. H. Pan, F. Huang, Q. Zheng, H. Chen and B. Tang, *Chem. Commun.*, 2015, **51**, 3102–3105.
- 17 (a) L. Yuan, W. Lin, S. Zhao, W. Gao, B. Chen, L. He and S. Zhu, *J. Am. Chem. Soc.*, 2012, **134**, 13510–13523; (b) H. Chen, W. Lin and L. Yuan, *Org. Biomol. Chem.*, 2013, **11**, 1938–1941; (c) J. A. Richard, *Org. Biomol. Chem.*, 2015, **13**, 8169–8172; (d) H. Chen, W. Lin, H. Cui and W. Jiang, *Chem.-Eur. J.*, 2015, **21**, 733–745.
- 18 (a) H. Kojima, N. Nakatsubo, K. Kikuchi, S. Kawahara, Y. Kirino, H. Nagoshi, Y. Hirata and T. Nagano, *Anal. Chem.*, 1998, **70**, 2446–2453; (b) Y. Gabe, Y. Urano, K. Kikuchi, H. Kojima and T. Nagano, *J. Am. Chem. Soc.*, 2004, **126**, 3357–3367; (c) E. Sasaki, H. Kojima, H. Nishimatsu, Y. Urano, K. Kikuchi, Y. Hirata and T. Nagano, *J. Am. Chem. Soc.*, 2005, **127**, 3684–3685.
- 19 (a) H. E. Ganther, *Biochemistry*, 1971, **10**, 4089–4098; (b) M. Wallenberg, E. Olm, C. Hebert, M. Björnstedt and A. P. Fernandes, *Biochem. J.*, 2010, **429**, 85–93; (c) S. J. Fairweather-Tait, Y. Bao, M. R. Broadley, R. Collings, D. Ford, J. E. Hesketh and R. Hurst, *Antioxid. Redox Signaling*, 2011, **14**, 1337–1383; (d) M. Björnstedt, S. Kumar and A. Holmgren, *J. Biol. Chem.*, 1992, **267**, 8030–8034; (e) S. Kumar, M. Björnstedt and A. Holmgren, *Eur. J. Biochem.*, 1992, **207**, 435–439; (f) M. Björnstedt, M. Hamberg, S. Kumar, J. Xue and A. Holmgren, *J. Biol. Chem.*, 1995, **270**, 11761–11764.
- 20 L. Yan and J. E. Spallholz, *Biochem. Pharmacol.*, 1993, **45**, 429–437.
- 21 (a) N. Karton-Lifshin, E. Segal, L. Omer, M. Portnoy, R. Satchi-Fainaro and D. Shabat, *J. Am. Chem. Soc.*, 2011, **133**, 10960–10965; (b) X. Pan, X. Wang, L. Wang, K. Xu, F. Kong and B. Tang, *Anal. Chem.*, 2015, **87**, 7092–7097.
- 22 (a) P. Vaupel, F. Kallinowski and P. Okunieff, *Cancer Res.*, 1989, **49**, 6449–6465; (b) H. Harada, *Nat. Commun.*, 2012, **3**, 783; (c) D. Hanahan and R. A. Weinberg, *Cell*, 2000, **100**, 57–70; (d) M. C. Hung, G. B. Mills and D. Yu, *Nat. Med.*, 2009, **15**, 246–247.
- 23 (a) N. S. Rajasekaran, P. Connell, E. S. Christians, L.-J. Yan, R. P. Taylor, A. Orosz, X.-Q. Zhang, T. J. Stevenson, R. M. Peshock, J. A. Leopold, W. H. Barry, J. Loscalzo, S. J. Odelberg and I. J. Benjamin, *Cell*, 2007, **130**, 427–439; (b) V. M. Labunsky, B. C. Lee, D. E. Handy, J. Loscalzo, D. L. Hatfield and V. N. Gladyshev, *Antioxid. Redox Signaling*, 2011, **14**, 2327–2336.
- 24 (a) C. Mealli, S. Midollini and L. Sacconi, *Inorg. Chem.*, 1978, **17**, 632–637; (b) V. V. Matyilitsky, A. Shavel, N. Gaponik, A. Eychmüller and J. Wachtveitl, *J. Phys. Chem. C*, 2008, **112**, 2703–2710.