

Cite this: *Chem. Sci.*, 2016, 7, 1611

## Correction: Base pairing involving artificial bases *in vitro* and *in vivo*

Omprakash Bande,<sup>b</sup> Darren Braddick,<sup>a</sup> Stefano Agnello,<sup>b</sup> Miyeon Jang,<sup>b</sup> Valérie Pezo,<sup>a</sup> Guy Schepers,<sup>b</sup> Jef Rozenski,<sup>b</sup> Eveline Lescrinier,<sup>b</sup> Philippe Marlière<sup>a</sup> and Piet Herdewijn<sup>\*ab</sup>

DOI: 10.1039/c5sc90070k

Correction for 'Base pairing involving artificial bases *in vitro* and *in vivo*' by Omprakash Bande *et al.*, *Chem. Sci.*, 2016, DOI: 10.1039/c5sc03474d.[www.rsc.org/chemicalscience](http://www.rsc.org/chemicalscience)

In this manuscript, the captions of Fig. 10 and Fig. 11 were mistakenly swapped. The correct legends should read:

**Fig. 10** The *in vivo* interpretation of the 8-Oxo-dI nucleotide within the essential thyA gene. The normalized ratio is the experimentally derived average number of thymidine-prototrophic colonies ( $\text{bla}^+ \text{thyA}^+$ ) from the average total number of colonies ( $\text{bla}^+ \text{thyA}^-$  and  $\text{bla}^+ \text{thyA}^+$ ). The modified section of the oligomer sequence indicates the position of the 8-Oxo-dI nucleotide/s, for single (left) and double (right) codons.

**Fig. 11** The *in vivo* interpretation of the 8-Oxo-dG nucleotide (top) and 8-Oxo-dA nucleotide (bottom) within the essential thyA gene. The normalized ratio is the experimentally derived average number of thymidine-prototrophic colonies ( $\text{bla}^+ \text{thyA}^+$ ) from the average total number of colonies ( $\text{bla}^+ \text{thyA}^-$  and  $\text{bla}^+ \text{thyA}^+$ ). The modified section of the oligomer sequence indicates the position of the 8-Oxo-dG and 8-Oxo-dA nucleotide/s.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup>iSSB – CNRS FRE3561, University of Evry-Val-d'Essonne, 5 rue Henri Desbruères, Genopole Campus 1, Bât. 6, F-91030 Évry Cedex, France. E-mail: [piet.herdewijn@rega.kuleuven.be](mailto:piet.herdewijn@rega.kuleuven.be); Tel: +32 16 337387

<sup>b</sup>Medicinal Chemistry, Rega Institute for Medical Research, KU Leuven, Minderbroedersstraat 10, 3000 Leuven, Belgium

