



CrossMark  
click for updates

Cite this: *Chem. Sci.*, 2016, 7, 5989

Received 18th May 2016

Accepted 23rd May 2016

DOI: 10.1039/c6sc02191c

www.rsc.org/chemicalscience

# Diastereodivergent combined carbometalation/ zinc homologation/C–C fragmentation reaction as an efficient tool to prepare acyclic allylic quaternary carbon stereocenters†

Sudipta Raha Roy, Dorian Didier, Amir Kleiner and Ilan Marek\*


A new strategy has been developed to construct enantiomerically enriched acyclic allylic quaternary carbon stereocenters in a single-pot operation through a combined carbometalation/zinc homologation/fragmentation sequence. Proper tuning of the reaction conditions enables the synthesis of the two enantiomers starting from a single enantiomer of the starting material.

## Introduction


In the last few decades, numerous approaches to integrate multiple chemical steps in a single-pot operation<sup>1</sup> have been described, offering reliable and powerful strategies for the synthesis of fine chemicals.<sup>2</sup> In this context, the construction of several carbon–carbon (C–C) bonds with simultaneous control of newly formed asymmetric centers, including the formation of challenging quaternary carbon stereocenters,<sup>3</sup> is of paramount importance for the development of complex molecular frameworks.<sup>4</sup> Particularly interesting would be the formation of such stereocenters adjacent to allylic motifs<sup>4</sup> as these sub-structures are abundant in biomolecules and natural products.<sup>5</sup> Although several excellent and highly selective approaches have been reported for the construction of acyclic allylic carbon quaternary stereocenters,<sup>6–10</sup> it becomes more intricate when the synthesis has to be performed in a single-pot operation.<sup>11</sup> For instance, although copper catalysed asymmetric conjugate addition is probably the most well established transformation (Fig. 1, path A),<sup>12</sup> asymmetric 1,4-addition to an extended conjugated system is more challenging as the 1,6-addition product is usually preferred over 1,4-addition (Fig. 1, path B).<sup>13</sup> A more successful alternative approach to reach the same products is the asymmetric 1,4-addition of vinyl metal species to Michael acceptors.<sup>14</sup> Although methods coupling asymmetric catalysis with fragmentation of strained building blocks have been reported recently, none could be used for the preparation of acyclic allylic carbon quaternary

stereocenters.<sup>15</sup> Herein, we would like to report our efforts to address this issue by performing a new tandem approach leading to the formation of two new carbon–carbon bonds in a single-pot operation, including the formation of the desired acyclic allylic quaternary carbon stereocenter (Fig. 1, Path C) from easily accessible starting materials.


path A: *Asymmetric conjugate addition*



path B: *Asymmetric conjugate addition on extended system*



path C: *This work*



The Mallat Family Laboratory of Organic Chemistry, Schulich Faculty of Chemistry and Lise Meitner-Minerva Center for Computational Quantum Chemistry, Technion-Israel Institute of Technology, Technion City, Haifa 32000, Israel. E-mail: chilann@tx.technion.ac.il; Fax: +972-4-829-37-09; Tel: +972-4-829-37-09

† Electronic supplementary information (ESI) available: Experimental procedures, spectroscopic data and copies of <sup>1</sup>H and <sup>13</sup>C NMR spectra. See DOI: 10.1039/c6sc02191c

Fig. 1 Approaches to construct allylic quaternary stereocenters.



This combined reaction consists of the diastereoselective carbometalation reaction<sup>16</sup> of enantiomerically enriched substituted cyclopropenyl esters **1**, followed by a homologation reaction<sup>17</sup> of the resulting cyclopropyl metal species **2** leading to a cyclopropylmethyl metal **3**, which subsequently undergoes a carbon–carbon bond cleavage<sup>18</sup> to give the enantiomerically enriched acyclic allylic quaternary carbon stereocenter **4** as described in Fig. 1, path C. Although this proposed sequence is very appealing, the unique formation of the linear product **4** directly from **1**, promoted by a combination of organometallic species in high chemical yield and with a high enantiomeric ratio, requires a cascade of high-yielding events with perfect control of all the elementary steps. It should be noted that the enantiomerically enriched cyclopropenyl ester starting materials **1** are readily accessible through the asymmetric metal-catalyzed decomposition of diazoesters with alkynes.<sup>19</sup> The enantiomeric ratio of the acyclic allylic quaternary stereocenter in **4** results from the diastereoselectivity of the carbometalation reaction.

## Results and discussion


We started our research by investigating the diastereoselectivity of the carbometalation reaction of cyclopropenyl ester **1**. In our synthetic plans, the presence of the ester was not only needed to achieve good diastereoselectivity for the carbometalation reaction, but was also a key element for successful ring-fragmentation of the newly formed donor–acceptor cyclopropylmethyl metal species **3**.<sup>20</sup>

Hence, to address the initial step of this sequence (transformation of **1** into **2**), our model cyclopropenyl ester substrate (racemic **1a**) was treated with various organocopper species in different solvents as described in Table 1. When **1a** was treated at  $-45\text{ }^{\circ}\text{C}$  in THF with 1.05 equivalents of MeCu, easily prepared

by mixing MeLi and CuI in a 1 : 1 ratio, the carbometalated product **2a<sub>anti</sub>** was rapidly obtained after hydrolysis (less than 30 minutes) with a very high *anti*-diastereoselectivity in good yield (Table 1, entry 1).<sup>21</sup> The configuration of **2a<sub>anti</sub>** was established by NOE experiments (see ESI†). The preferential formation of diastereomer **2a<sub>anti</sub>** can only be possible if the carbometalation reaction is sterically driven. A coordinating solvent such as THF makes the organocopper species less prone to chelation by the ester and therefore it prefers to react on the *re*-face of the cyclopropene derivative **1a**. If the assumption that solvent plays a crucial role in the control of the diastereoselectivity of the carbometalation step is correct, then a less Lewis basic solvent should favour addition on the *si*-face through coordination of the organometallic species with the ester. Indeed, when a slightly less coordinating solvent such as 2-methyl THF was used, the selectivity towards the formation of **2a<sub>anti</sub>** dropped slightly (Table 1, entry 2). On further decreasing the Lewis basicity of the solvent (Table 1, entries 3 to 5, Et<sub>2</sub>O, hexane and toluene respectively), the tendency of the organometallic species to be coordinated by the ester group increases and therefore the isomer **2a<sub>syn</sub>** could be prepared as the unique diastereoisomer (Table 1, entry 5). The relative configuration of **2a<sub>syn</sub>** was established by comparing the NOE and <sup>13</sup>C NMR experiments with **2a<sub>anti</sub>** (see ESI†). The same trend was also found for the addition of organocopper species originating from Grignard reagent (prepared by mixing MeMgBr and CuI in a 1 : 1 ratio), and although the formation of the *anti*-addition product **2a<sub>anti</sub>** in THF was less diastereoselective (Table 1, compare entries 1 and 6), it was still excellent for the formation of the *syn*-diastereoisomer **2a<sub>syn</sub>** (Table 1, entry 8). It should be noted that diastereoselective carbometalation of cyclopropenyl esters such as **1a** has already been described, but it was always achieved through variation of the organometallic species and never by simple variation of the solvent for a given organocopper entity.<sup>16</sup>

Having access to both *syn* and *anti* diastereoisomers of the carbometalated products at will according to the nature of the solvent, we then turned our attention to the zinc homologation reaction of the racemic cyclopropyl copper species **2Cu<sub>anti</sub>** originating from the carbometalation reaction of MeCu in THF (MeCu was formed from MeLi and CuI as described in Table 1, entry 1). For this purpose, CH<sub>2</sub>I<sub>2</sub> and Et<sub>2</sub>Zn in a 1 : 1 ratio were added to the reaction mixture at  $-45\text{ }^{\circ}\text{C}$ , and upon warming to  $-20\text{ }^{\circ}\text{C}$ , within 2 hours, the homologated cyclopropylmethyl zinc derivative<sup>22</sup> **3** gave **4a** as described in Fig. 2. Indeed, as cyclopropylcopper **2Cu<sub>anti</sub>** does not react with CH<sub>2</sub>I<sub>2</sub>, the reaction between Et<sub>2</sub>Zn and CH<sub>2</sub>I<sub>2</sub> occurs first leading to the *in situ* formation of the zinc carbenoid ICH<sub>2</sub>ZnEt.<sup>23</sup> Then, **2Cu<sub>anti</sub>** is homologated by the zinc carbenoid to generate the *in situ* reactive cyclopropylmethyl zinc (or copper) derivative **3** (see Fig. 1), which instantaneously undergoes a C–C bond fragmentation.<sup>24</sup> Although the combined carbometalation/zinc homologation/fragmentation reaction proceeded smoothly according to our original plan to give **4a**, the conversion of **2Cu<sub>anti</sub>** into **4a** was only moderate (**2a<sub>anti</sub>**/**4a** = 27/73 after hydrolysis).

Table 1 Controlling the diastereoselectivity for the initial carbocupration step



Entry	Me[M]	Solvent	<b>2a<sub>syn</sub></b> / <b>2a<sub>anti</sub></b> <sup>a</sup>
1	MeLi	THF	1 : 99 (75)
2	MeLi	2-MeTHF	6 : 94
3	MeLi	Et <sub>2</sub> O	89 : 11
4	MeLi	Hexane	97 : 3
5	MeLi	C <sub>6</sub> H <sub>5</sub> Me	99 : 1 (73)
6	MeMgBr	THF	28 : 72
7	MeMgBr	Et <sub>2</sub> O	99 : 1
8	MeMgBr	C <sub>6</sub> H <sub>5</sub> Me	99 : 1 (72)

<sup>a</sup> **1a** was consumed completely and the ratio between products **2a<sub>syn</sub>** and **2a<sub>anti</sub>** was determined by GC-MS of the crude reaction mixture. Numbers in parentheses represent the isolated yields after purification by column chromatography.






Fig. 2 Combined carbometalation/zinc homologation/fragmentation reaction of cyclopropyl ester **1**.

To improve the conversion, it was then anticipated that increasing the nucleophilicity of the carbon atom bearing the organocopper species should increase its reactivity towards the ambiphilic zinc carbenoid species EtZnCH<sub>2</sub>I.




Fig. 3 Diastereodivergent combined carbometalation/zinc homologation/fragmentation en route to enantioenriched allylic quaternary carbon stereocenters.




Fig. 4 Combined carbometalation/zinc homologation/fragmentation reactions for the preparation of the opposite enantiomer.




Fig. 5 Diastereodivergent strategy for the preparation of enantioenriched **4**.

To this end, various donor ligands were added to the reaction mixture to improve the reactivity of the cyclopropyl copper species (see ligands in Fig. 2).<sup>25</sup> Indeed, we were pleased to observe that the addition of TMEDA (**L<sub>1</sub>**) or 2-2'-bipyridine (**L<sub>2</sub>**) as ligand improved the conversion of the desired product **4a** (**2a<sub>anti</sub>**/**4a** = 12/88 and 13/87 respectively after hydrolysis).



The best conversion could be reached when phenanthroline ( $L_3$ ) was added to  $2Cu_{anti}$ , as the ratio became excellent ( $2a_{anti}/4a = 6/94$  after hydrolysis) and the final product **4a** could be isolated in 72% yield. This sequence of carbocupration/zinc homologation/C–C bond cleavage has been generalized to different functionalized substrates (see scope in Fig. 2) and in all cases, the reactions proceed smoothly for the one-pot transformation of **1** into **4**.

Both cyclopropenyl esters (–OEt and –OBn) undergo this combined transformation (compare **4b** to **4c**, **4d** to **4e** and **4f** to **4g** in Fig. 2), but it should be noted that the products resulting from the reaction with the benzyl ester (**4b**, **4d**, **4g**) are easier to purify by column chromatography from the remaining carbometalated products after hydrolysis ( $2_{anti}$ ) than the products possessing the ethyl ester (**4c**, **4e**, **4f**). These combined reactions are not restricted to the introduction of a methyl group, as various other alkyl groups could be added in good yields by simply changing the nature of the starting organocopper reagent (compare **4b** to **4d**, **4f** to **4h** and **4g** to **4i**, Fig. 2). The reaction of cyclopropenyl ester **1d** with phenylcopper also proceeds well, but we were unable to separate the desired product **4m** from the carbometalated product  $2m_{anti}$  after hydrolysis. Similarly, when we treated a vinylcopper derivative (prepared from ethenylmagnesium bromide and CuI in a 1 : 1 ratio), as a representative example of an  $sp^2$  organometallic species, with cyclopropenyl ester **1j**, the reaction proceeds smoothly but again, we were unable to separate the desired product **4n** from the hydrolysed carbometalated product  $2n_{anti}$ . It is worth mentioning that this sequence of vinyl cupration/zinc homologation/fragmentation opens new avenues for the preparation of skipped dienes possessing a quaternary carbon stereocenter. Notably, the presence of other functional groups in the original alkyl chain of the cyclopropenyl ester is well tolerated (formation of **4j–l**, Fig. 2). Having established an easy protocol for the preparation of racemic **4** from these very simple starting materials, the synthesis of enantiomerically enriched cyclopropenyl esters **1b** ( $R^1 = Ph(CH_2)_2$ ), **1e** ( $R^1 = PhCH_2$ ) and **1f** ( $R^1 = Hex$ ) was easily achieved in good enantiomeric ratio (**1b** *er* 96 : 4, **1e** *er* 93 : 7 and **1f** *er* 91 : 9) through cyclopropenylation of the terminal alkyne with benzyl diazoacetate and  $Rh_2(OAc)(R,R-DPTI)_3$  (DPTI = diphenyltriflylimidazolidinone) as catalyst.<sup>26</sup> Interestingly, the enantiomeric ratios of these cyclopropenyl benzyl esters are lower than the ones obtained for cyclopropenyl ethyl esters (ethyl diazoacetate generally leads to enantiomeric ratios in the range of 97 : 3), but for the sake of simplicity, we decided to illustrate our concept with starting materials that would require a minimum number of chemical steps. When the sequence of carbocupration, zinc homologation and C–C bond fragmentation was performed on these enantiomerically enriched cyclopropenyl esters, enantioenriched acyclic allylic molecular architectures (**4b,g,h,i**) possessing a quaternary carbon stereocenter were easily obtained through the creation of two C–C bonds in a single-pot operation. HPLC analyses (see ESI†) show that the chiral information at the carbon atom connected to the ester moiety in **1b,e** and **f** was quantitatively transferred to the final products **4b** (*er* 96 : 4), **4g** and **4h** (*er* 93 : 7) and **4i** (*er* 91 : 1), Fig. 2. It was therefore clear that the

quaternary stereocenter was at no risk of epimerization through the whole process, and one could prepare the expected products with the same enantiomeric ratio as in the starting materials.

In contrast to any typical enantioselective synthesis where the independent formation of the two enantiomers of a product can only be achieved by changing the chirality of the ligand (or substrate) associated with the enantioselective transformation,<sup>27</sup> the diastereodivergent carbometalation of **1**, controlled by the nature of the solvent as described in Table 1, should therefore lead to the enantiodivergent synthesis of **4** from the same enantiomerically enriched starting material **1** (as summarized in Fig. 3).

Although the best ligand to promote the zinc homologation of  $2Cu_{anti}$  was phenanthroline  $L_3$ , it was unclear that the same ligand would be optimal when the first step was performed in a non-polar solvent. To reach this new goal, we had to again optimise the reaction sequence for the formation of the desired products **4** from the *syn*-diastereomer  $2Cu_{syn}$  in the presence of a non-polar solvent (such as toluene, see Table 1, entries 5 and 8). It was first rapidly found that the zinc homologation for the transformation of  $2Cu_{syn}$  into **4a** was more efficient when the initial organocopper species was prepared from an organolithium ( $R^2Cu \cdot LiI$ , Table 1, entry 5) instead of an organomagnesium species ( $R^2Cu \cdot MgX_2$ , Table 1, entry 7). After extensive experiments, we were pleased to find that the zinc homologation could now be best performed in the presence of a non-nucleophilic potassium *tert*-butoxide ligand (for all other ligands tested, see ESI†) with addition of THF as co-solvent. It is important to note that the reaction performed under identical conditions but in the absence of the potassium *tert*-butoxide ligand results in very poor conversion, which underlines the effect of this particular ligand on the nucleophilicity of the metalated carbon center stabilized by the chelating ester. Once the best experimental conditions were in hand, various allylic esters **4** possessing quaternary carbon stereocenters were prepared in a single-pot operation from cyclopropenyl esters **1** as illustrated in Fig. 4.

The scope of this new reaction sequence of *syn*-diastereoselective carbometalation, zinc homologation and selective C–C bond cleavage is equally good albeit with slightly lower yields for the final products **4**. To illustrate the potential of this diastereodivergent approach, both enantiomers of **4i** were prepared from the same cyclopropenyl ester enantiomer **1f** ( $R^1 = Hex$ , *er* 91 : 9). When the carbocupration of (*S*)-**1f** was performed with  $MeCu \cdot LiI$  in THF followed by the subsequent addition of  $CH_2I_2$ ,  $Et_2Zn$  and phenanthroline ( $L_3$ ) as ligand, the selective C–C bond cleavage led to the corresponding allylic (*S*)-**4i** with an enantiomeric ratio of 91 : 9 in 65% yield (Fig. 5).

On the other hand, if the same cyclopropenyl ester (*S*)-**1f** is added to the organocopper  $MeCu \cdot LiI$  in toluene followed by subsequent zinc homologation in the presence of potassium *tert*-butoxide as ligand, the selective C–C bond cleavage provides the allylic (*R*)-**4i** in similar yield (63%) with almost the same enantiomeric ratio (*er* 90 : 10) (Fig. 5).

## Conclusions

A new sequence of diastereoselective carbometalation/zinc homologation/C–C bond cleavage allows the easy transformation



of enantiomerically enriched cyclopropenyl esters into acyclic allylic moieties bearing challenging quaternary carbon stereocenters in a single-pot reaction through the formation of two new C–C bonds. As the carbometalation reaction may lead to two different diastereoisomers according to the nature of the solvent, this strategy paves the way to the diastereodivergent synthesis of both enantiomers of **4** at will.

## Acknowledgements

This research is supported by the European Research Council under the European Community's Seventh Framework Program (ERC grant agreement no. 338912) and by the Israel Science Foundation administered by the Israel Academy of Sciences and Humanities (140/12). S. R. R. acknowledges the Planning and Budgeting Committee (PBC) for financial assistance. I. M. is holder of the Sir Michael and Lady Sobell Academic Chair.

## Notes and references

- For recent reviews, see: (a) H. Pellissier, *Tetrahedron*, 2013, **69**, 7171; (b) H. Pellissier, *Chem. Rev.*, 2013, **113**, 442; (c) A. E. Allen and D. W. C. MacMillan, *Chem. Sci.*, 2012, **3**, 633; (d) N. T. Patil, V. S. Shinde and B. Gajula, *Org. Biomol. Chem.*, 2012, **10**, 211; (e) H. Clavier and H. Pellissier, *Adv. Synth. Catal.*, 2012, **354**, 3347; (f) C. De Graaff, E. Ruijter and R. V. A. Orru, *Chem. Soc. Rev.*, 2012, **41**, 3969; (g) H. Pellissier, *Adv. Synth. Catal.*, 2012, **354**, 237; (h) Z. Zhang and J. C. Antilla, *Angew. Chem., Int. Ed.*, 2012, **51**, 11778; (i) S. Piovesana, D. M. Scarpino Schietroma and M. Bella, *Angew. Chem., Int. Ed.*, 2011, **50**, 6216; (j) L. Albrecht, H. Jiang and K. A. Jørgensen, *Angew. Chem., Int. Ed.*, 2011, **50**, 8492; (k) M. Ruiz, P. Lopez-Alvarado, G. Giorgi and J. C. Menendez, *Chem. Soc. Rev.*, 2011, **40**, 3445; (l) C. Vaxelaire, P. Winter and M. Christmann, *Angew. Chem., Int. Ed.*, 2011, **50**, 3605.
- R. W. Dugger, J. A. Ragan and D. H. B. Ripin, *Org. Process Res. Dev.*, 2005, **9**, 253.
- For recent reviews on the formation of quaternary carbon stereogenic centers, see: (a) K. W. Quasdorf and L. E. Overman, *Nature*, 2014, **516**, 181; (b) A. Y. Hong and B. M. Stoltz, *Eur. J. Org. Chem.*, 2013, 2745; (c) J. P. Das and I. Marek, *Chem. Commun.*, 2011, **47**, 4593; (d) C. Hawner and A. Alexakis, *Chem. Commun.*, 2010, **46**, 7295; (e) M. Bella and T. Casperi, *Synthesis*, 2009, 1583; (f) P. G. Cozzi, R. Hilgraf and N. Zimmermann, *Eur. J. Org. Chem.*, 2007, 5969; (g) I. Marek and G. Sklute, *Chem. Commun.*, 2007, **43**, 1683; (h) B. M. Trost and C. Jiang, *Synthesis*, 2006, 369; (i) J. Christoffers and A. Baro, *Adv. Synth. Catal.*, 2005, **347**, 1473; (j) C. J. Douglas and L. E. Overman, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 5363; (k) I. Denissova and L. Barriault, *Tetrahedron*, 2003, **59**, 10105; (l) J. Christoffers and A. Mann, *Angew. Chem., Int. Ed.*, 2001, **40**, 4591; (m) E. J. Corey and A. Guzman-Perez, *Angew. Chem., Int. Ed.*, 1998, **37**, 388.
- (a) G. Eppe, D. Didier and I. Marek, *Chem. Rev.*, 2015, **115**, 9175; (b) I. Marek, Y. Minko, M. Pasco, T. Mejuch, N. Gilboa, H. Chechik and J. P. Das, *J. Am. Chem. Soc.*, 2014, **136**, 2682; (c) T. Newhouse, P. S. Baran and R. W. Hoffmann, *Chem. Soc. Rev.*, 2009, **38**, 3010; (d) P. S. Baran, T. J. Maimone and J. M. Richter, *Nature*, 2007, **446**, 404; (e) L. F. Tietze, *Chem. Rev.*, 1996, **96**, 115; (f) B. M. Trost, *Science*, 1991, **254**, 1471; (g) P. A. Wender and B. L. Miller, *Nature*, 2009, **460**, 197; (h) I. Marek, *Chem.–Eur. J.*, 2008, **14**, 7460.
- (a) K.-i. Takao, K. Tsunoda, T. Kurisu, A. Sakama, Y. Nishimura, K. Yoshida and K.-i. Tadano, *Org. Lett.*, 2015, **17**, 756; (b) L.-Q. Wang, Y.-G. Chen, J.-J. Xu, Y. Liu, X.-M. Li and Y. Zhao, *Chem. Biodiversity*, 2008, **5**, 1879; (c) B. Biswas and R. V. Venkateswaran, *Synth. Commun.*, 2005, **35**, 1769; (d) Y. Kita, A. Furukawa, J. Futamura, K. Ueda, Y. Sawama, H. Hamamoto and H. Fujioka, *J. Org. Chem.*, 2001, **66**, 8779; (e) M. Takahashi, Y. Shioura, T. Murakami and K. Ogasawara, *Tetrahedron: Asymmetry*, 1997, **8**, 1235.
- (a) M. Raducan, R. Alam and K. J. Szabó, *Angew. Chem., Int. Ed.*, 2012, **51**, 7263; (b) Y. Yatsumonji, T. Nishimura, A. Tsubouchi, K. Noguchi and T. Takeda, *Chem.–Eur. J.*, 2009, **15**, 2680; (c) G. Dunet, P. Meyer and P. Knochel, *Org. Lett.*, 2008, **10**, 117; (d) H. Ren, G. Dunet, P. Mayer and P. Knochel, *J. Am. Chem. Soc.*, 2007, **129**, 5376; (e) L. F. Tietze, T. Knizel and S. Schmatz, *J. Am. Chem. Soc.*, 2006, **128**, 11483; (f) M. Yasuda, K. Hirata, A. Nishino, A. Yamamoto and A. Baba, *J. Am. Chem. Soc.*, 2002, **124**, 13442.
- (a) J.-N. Heo, G. C. Micalizio and W. R. Roush, *Org. Lett.*, 2003, **5**, 1693; (b) J. W. J. Kennedy and D. G. Hall, *J. Am. Chem. Soc.*, 2002, **124**, 898; (c) S. E. Denmark and J. Fu, *Org. Lett.*, 2002, **4**, 1951; (d) S. E. Denmark and J. Fu, *J. Am. Chem. Soc.*, 2001, **123**, 9488.
- (a) R. M. Maksymowicz, A. J. Bissette and S. P. Fletcher, *Chem.–Eur. J.*, 2015, **21**, 5668; (b) K. Hojoh, Y. Shido, H. Ohmiya and M. Sawamura, *Angew. Chem., Int. Ed.*, 2014, **53**, 4954; (c) F. Gao, K. P. McGrath, Y. Lee and A. H. Hoveyda, *J. Am. Chem. Soc.*, 2010, **132**, 14315; (d) C. A. Luchaco-Cullis, H. Mizutani, K. E. Murphy and A. H. Hoveyda, *Angew. Chem., Int. Ed.*, 2001, **40**, 1456.
- (a) F. Nowrouzi, A. N. Thadani and R. A. Batey, *Org. Lett.*, 2009, **11**, 2631; (b) U. Schneider, M. Ueno and S. Kobayashi, *J. Am. Chem. Soc.*, 2008, **130**, 13284; (c) L. Carosi and D. G. Hall, *Angew. Chem., Int. Ed.*, 2007, **46**, 5913; (d) S. R. Denmark, J. Fu and M. J. Lawler, *J. Org. Chem.*, 2006, **71**, 1523; (e) S. R. Denmark, J. Fu, D. M. Coe, X. Su, N. E. Pratt and B. D. Griedel, *J. Org. Chem.*, 2006, **71**, 1513; (f) R. Wada, K. Oisaki, M. Kanai and M. Shibasaki, *J. Am. Chem. Soc.*, 2004, **126**, 8910.
- (a) R. Alam, T. Vollgraff, L. Eriksson and K. J. Szabó, *J. Am. Chem. Soc.*, 2015, **137**, 11262; (b) J. L.-Y. Chen and V. K. Aggarwal, *Angew. Chem., Int. Ed.*, 2014, **53**, 10992.
- M. Yus, J. C. González-Gómez and F. Foubelo, *Chem. Rev.*, 2013, **113**, 5595.
- (a) D. Müller, M. Tissot and A. Alexakis, *Org. Lett.*, 2011, **13**, 3040; (b) Q. Naeemi, T. Robert, D. Kranz, J. Velder and H.-G. Schmalz, *Tetrahedron: Asymmetry*, 2011, **22**, 887; (c) T. Robert, J. Velder and H.-G. Schmalz, *Angew. Chem., Int.*



- Ed.*, 2008, **47**, 7718; (d) M. Vuagnoux-d'Augustin and A. Alexakis, *Chem.-Eur. J.*, 2007, **13**, 9647; (e) K. H. Ahn, R. B. Klassen and S. J. Lippard, *Organometallics*, 1990, **9**, 3178.
- 13 (a) T. den Hartog, Y. Huang, M. Fananas-Mastral, A. Meuwese, A. Rudolph, M. Perez, A. J. Minnaard and B. L. Feringa, *ACS Catal.*, 2015, **5**, 560; (b) N. Yoshikai, T. Yamashita and E. Nakamura, *Angew. Chem., Int. Ed.*, 2005, **44**, 4721; (c) S. Mori, M. Uerdingen, N. Kraus and K. Morokuma, *Angew. Chem., Int. Ed.*, 2005, **44**, 4715; (d) Y. Yamamoto, S. Yamamoto, H. Yatagai, Y. Ishihara and K. Maruyama, *J. Org. Chem.*, 1982, **47**, 119.
- 14 (a) K. P. McGrath and A. H. Hoveyda, *Angew. Chem., Int. Ed.*, 2014, **53**, 1910; (b) D. Müller and A. Alexakis, *Chem. Commun.*, 2012, **48**, 12037.
- 15 (a) L. Soullart and N. Cramer, *Chem. Sci.*, 2014, **5**, 837; (b) T. Seiser and N. Cramer, *J. Am. Chem. Soc.*, 2010, **132**, 5340; (c) T. Seiser and N. Cramer, *Angew. Chem., Int. Ed.*, 2008, **47**, 9294; (d) T. Matsuda, M. Shigeno and M. Murakami, *J. Am. Chem. Soc.*, 2007, **129**, 12086.
- 16 D. Didier, P.-O. Delaye, M. Simaan, B. Island, G. Eppe, H. Eijlsberg, A. Kleiner, P. Knochel and I. Marek, *Chem.-Eur. J.*, 2014, **20**, 1038.
- 17 (a) R. Vabre, B. Island, C. J. Diehl, P. R. Schreiner and I. Marek, *Angew. Chem., Int. Ed.*, 2015, **54**, 9996; (b) T. Mejuch, M. Botoshansky and I. Marek, *Org. Lett.*, 2011, **13**, 3604; (c) B. Dutta, N. Gilboa and I. Marek, *J. Am. Chem. Soc.*, 2010, **132**, 5588; (d) G. Sklute and I. Marek, *J. Am. Chem. Soc.*, 2006, **128**, 4642; (e) A. Abramovitch, J. P. Varghese and I. Marek, *Org. Lett.*, 2004, **6**, 621; (f) G. Sklute, D. Amsallem, A. Shabli, J. P. Varghese and I. Marek, *J. Am. Chem. Soc.*, 2003, **125**, 11776.
- 18 For recent reviews on C-C cleavage, see: (a) I. Marek, A. Masarwa, P.-O. Delaye and M. Leibelng, *Angew. Chem., Int. Ed.*, 2015, **54**, 414; (b) K. Ruhland, *Eur. J. Org. Chem.*, 2012, 2683; (c) T. Seiser, T. Saget, D. N. Tran and N. Cramer, *Angew. Chem., Int. Ed.*, 2011, **50**, 7740; (d) T. Seiser and N. Cramer, *Org. Biomol. Chem.*, 2009, **7**, 2835; (e) Y. J. Park, J.-W. Park and C.-H. Jun, *Acc. Chem. Res.*, 2008, **41**, 222; (f) T. Satoh and M. Miura, *Top. Organomet. Chem.*, 2005, **14**, 1; (g) C.-H. Jun, *Chem. Soc. Rev.*, 2004, **33**, 610; (h) M. E. Van Der Boom and D. Milstein, *Chem. Rev.*, 2003, **103**, 1759; (i) B. Rybtchinski and D. Milstein, *Angew. Chem., Int. Ed.*, 1999, **38**, 870.
- 19 For recent examples of enantioselective cyclopropenylation of alkynes, see: (a) J. F. Briones and H. M. L. Davies, *J. Am. Chem. Soc.*, 2012, **134**, 11916; (b) M. Uehara, H. Suematsu, Y. Yasutomi and T. Katsuki, *J. Am. Chem. Soc.*, 2011, **133**, 170; (c) X. Cui, X. Xu, H. Lu, S. Zhu, L. Wojitas and X. P. Zhang, *J. Am. Chem. Soc.*, 2011, **133**, 3304; (d) K. Goto, K. Takeda, N. Shiamda, H. Nambu, M. Anada, M. Shiro, K. Ando and S. Hashimoto, *Angew. Chem., Int. Ed.*, 2011, **50**, 6803; (e) J. F. Briones and H. M. L. Davies, *Tetrahedron*, 2011, **67**, 4313; (f) J. F. Briones, J. H. Hansen, K. I. Hardcastle, J. Autschbach and H. M. L. Davies, *J. Am. Chem. Soc.*, 2010, **132**, 17211; (g) M. P. Doyle, W. R. Winchester, J. A. A. Hoorn, V. Lynch, S. H. Simonsen and R. Ghosh, *J. Am. Chem. Soc.*, 1993, **115**, 9968.
- 20 (a) T. F. Schneider, J. Kaschel and D. Werz, *Angew. Chem., Int. Ed.*, 2014, **53**, 5504; (b) M. A. Cavitt, L. H. Phun and S. France, *Chem. Soc. Rev.*, 2014, **43**, 804; (c) M. Yu and B. L. Pagenkopf, *Tetrahedron*, 2005, **61**, 321; (d) H.-U. Reissig and R. Zimmer, *Chem. Rev.*, 2003, **103**, 1151.
- 21 D. Müller and I. Marek, *Chem. Soc. Rev.*, 2016, DOI: 10.1039/c5cs00897b.
- 22 (a) M. Pasco, N. Gilboa, T. Mejuch and I. Marek, *Organometallics*, 2013, **32**, 942; (b) I. Marek, *Tetrahedron*, 2002, **58**, 9463; (c) P. Knochel, N. Jeong, M. J. Rozema and M. C. P. Yeh, *J. Am. Chem. Soc.*, 1989, **111**, 6474; (d) P. Knochel, T.-S. Chou, H. G. Chen, M. C. P. Yeh and M. J. Rozema, *J. Org. Chem.*, 1989, **54**, 5202; (e) A. R. Sidduri and P. Knochel, *J. Am. Chem. Soc.*, 1992, **114**, 7579; (f) A. R. Sidduri, M. J. Rozema and P. Knochel, *J. Org. Chem.*, 1993, **58**, 2694.
- 23 (a) A. Voituriez, L. E. Zimmer and A. B. Charette, *J. Org. Chem.*, 2010, **75**, 1244; (b) J. Furukawa, N. Kawabata and J. Nishimura, *Tetrahedron Lett.*, 1966, **7**, 3353.
- 24 (a) M. Simaan, P.-O. Delaye, M. Shi and I. Marek, *Angew. Chem., Int. Ed.*, 2015, **54**, 12345; (b) P. O. Delaye, D. Didier and I. Marek, *Angew. Chem., Int. Ed.*, 2013, **52**, 5333.
- 25 (a) E. Nakamura and S. Mori, *Angew. Chem., Int. Ed.*, 2000, **39**, 3750; (b) M. D. Janssen, M. A. Corsten, A. L. Spek, D. M. Grove and G. van Koten, *Organometallics*, 1996, **15**, 2810.
- 26 Y. Lou, M. Horikawa, R. A. Kloster, N. A. Hawryluk and E. J. Corey, *J. Am. Chem. Soc.*, 2004, **126**, 8916.
- 27 *Comprehensive Asymmetric Catalysis*, ed. E. N. Jacobsen, A. Pfaltz and H. Yamamoto, Springer, New York, 1999, vol. I-III, Suppl. I-II.

