

Cite this: *Chem. Sci.*, 2016, 7, 5596

DOI: 10.1039/c6sc90039a

[www.rsc.org/chemicalscience](http://www.rsc.org/chemicalscience)

## Correction: Essential role of hydride ion in ruthenium-based ammonia synthesis catalysts

Masaaki Kitano,<sup>a</sup> Yasunori Inoue,<sup>b</sup> Hiroki Ishikawa,<sup>b</sup> Kyosuke Yamagata,<sup>b</sup>  
Takuya Nakao,<sup>b</sup> Tomofumi Tada,<sup>a</sup> Satoru Matsuishi,<sup>a</sup> Toshiharu Yokoyama,<sup>ac</sup>  
Michikazu Hara<sup>\*bcd</sup> and Hideo Hosono<sup>\*abcd</sup>Correction for 'Essential role of hydride ion in ruthenium-based ammonia synthesis catalysts' by Masaaki Kitano *et al.*, *Chem. Sci.*, 2016, 7, 4036-4043.

The authors regret that in the original article incorrect units were used to define both the number of surface Ru atoms ( $N_s$ ) and the ammonia synthesis rate ( $r_{\text{NH}_3}$ ) in columns 6 and 7 of Table 1. In both cases, 'μmol' should have been used instead of 'mmol'. Therefore, the correct units for  $N_s$  and  $r_{\text{NH}_3}$  are 'μmol g<sup>-1</sup>' and 'μmol g<sup>-1</sup> h<sup>-1</sup>', respectively.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup>Materials Research Center for Element Strategy, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, Yokohama 226-8503, Japan. E-mail: hosono@mssl.titech.ac.jp

<sup>b</sup>Laboratory for Materials and Structures, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, Yokohama 226-8503, Japan. E-mail: mhara@mssl.titech.ac.jp

<sup>c</sup>ACCEL, Japan Science and Technology Agency, 4-1-8 Honcho, Kawaguchi, Saitama 332-0012, Japan

<sup>d</sup>Frontier Research Center, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, Yokohama 226-8503, Japan

