Chemical Science

EDGE ARTICLE



View Article Online View Journal | View Issue

Open Access Article. Published on 25 March 2024. Downloaded on 7/29/2025 2:25:52 AM.

Check for updates

Cite this: Chem. Sci., 2024, 15, 6028

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 2nd December 2023 Accepted 11th March 2024

DOI: 10.1039/d3sc06459j rsc.li/chemical-science

Introduction

Chemotherapy, the main method for treating malignant tumours, usually produces significant drug resistance during the later stages of treatment.¹ This reduces therapeutic effects and even results in resistance to alternative drugs that use different mechanisms, which is termed multidrug resistance (MDR).² MDR can interfere with the intake and accumulation of drugs in tumour cells, reducing the ability of anticancer drugs

‡ These authors contributed equally to this paper.

Near-infrared imaging for visualizing the synergistic relationship between autophagy and NFS1 protein during multidrug resistance using an ICT-TICT integrated platform⁺

Wei Hu, ^[b] ‡^{abc} Yifan He,‡^a Haixian Ren, ^{[b] *b} Li Chai,^b Haiyan Li,^a Jianbin Chen, ^{[b] d} Chunya Li, ^[b] *^a Yanying Wang^{*a} and Tony D. James ^[b] *^{ce}

Drug resistance is a major challenge for cancer treatment, and its identification is crucial for medical research. However, since drug resistance is a multi-faceted phenomenon, it is important to simultaneously evaluate multiple target fluctuations. Recently developed fluorescence-based probes that can simultaneously respond to multiple targets offer many advantages for real-time and *in situ* monitoring of cellular metabolism, including ease of operation, rapid reporting, and their non-invasive nature. As such we developed a dual-response platform (Vis-H₂S) with integrated ICT–TICT to image H₂S and viscosity in mitochondria, which could simultaneously track fluctuations in cysteine desulfurase (NFS1 protein and H₂S inducer) and autophagy during chemotherapy-induced multidrug resistance. This platform could monitor multiple endogenous metabolites and the synergistic relationship between autophagy and NFS1 protein during multidrug resistance induced by chemotherapy. The results indicated that chemotherapeutic drugs simultaneously up-regulate the levels of NFS1 protein and autophagy. It was also found that the NFS1 protein was linked with autophagy, which eventually led to multidrug resistance. As such, this platform could serve as an effective tool for the in-depth exploration of drug resistance mechanisms.

to induce cell death, resulting in drug inactivation and degradation, and the activation of anti-apoptosis and antioxidant defence, as well as DNA repair and replication. These factors are the main causes of the failure of chemotherapy.³ Common tumour MDR detection methods include MTT assays, drug sensitivity tests, MDR gene and pathway detection, and highthroughput screening technologies.⁴ However, using these methods makes it impossible to achieve *in situ*, real-time measurements and accurate detection of MDR in tumour cells. The methods are also expensive and time-consuming, which significantly limits their application for the evaluation of MDR mechanisms.

Confocal fluorescence imaging requires simple operation and exhibits high sensitivity and temporal and spatial resolution and can provide *in situ* and real-time analysis.⁵ However, fluorescence-based sensors based on traditional sensing mechanisms have only one recognition site and can only recognize one analyte due to their single mode sensing mechanism.⁶ These are unsuitable for evaluating multiple interconnected biomolecules in complex living systems. To overcome this deficiency, fluorescence-based sensors have emerged that use two or more sensing mechanisms.^{7,8}

Chemotherapeutic drugs which increase the amounts of reactive oxygen species (ROS) (such as doxorubicin and

^aKey Laboratory of Catalysis and Energy Materials Chemistry of Ministry of Education, Key Laboratory of Analytical Chemistry of the State Ethnic Affairs Commission, College of Chemistry and Materials Science, South-Central Minzu University, Wuhan 430074, China. E-mail: lichychem@mail.scuec.edu.cn; wangyychem@mail.scuec.edu.cn

^bDepartment of Chemistry, Xinzhou Normal University, Xinzhou, Shanxi, 034000, China. E-mail: hxren326@163.com

^cDepartment of Chemistry, University of Bath, Bath, BA27AY, UK. E-mail: t.d.james@ bath.ac.uk

^dSchool of Chemistry and Chemical Engineering, Qilu University of Technology (Shandong Academy of Sciences), Jinan, Shandong 250353, China

^eSchool of Chemistry and Chemical Engineering, Henan Normal University, Xinxiang 453007, China

[†] Electronic supplementary information (ESI) available: Experimental details and characterization probe Vis-H₂S supplementary spectral and cell experiments results. See DOI: https://doi.org/10.1039/d3sc06459j

cisplatin) can kill tumour cells. However, in addition they induce many factors that can result in MDR, including the activation of autophagy⁹ (caused by excessive ROS, which then prevents cancer cell apoptosis by degrading the ROS) and upregulating the levels of cysteine desulfurase¹⁰ (NFS1 protein, which can catalyse cysteine to produce endogenous hydrogen sulfide and consume ROS, which prevents cancer cell apoptosis). However, the synergistic relationship between autophagy and NFS1 protein overexpression *via* ROS-based chemotherapeutic drugs has not yet been reported.

To achieve the *in situ* imaging of the synergistic relationship between autophagy and reductive protein overexpression, a probe with mitochondrial anchoring ability with a dualresponse toward viscosity (changes in mitochondrial viscosity can indicate autophagy¹¹) and H₂S (mitochondrial H₂S fluctuations may indicate the NFS1 protein level¹²) is required. With this research, a cyanine dye **QCy7** with mitochondrial anchoring ability was used as the parent fluorophore¹³ and 2-iodobenzoate was used as the specific recognition site for H₂S (**Vis-H₂S**). When **Vis-H₂S** reacts with H₂S, intramolecular charge transfer (ICT) is activated and the fluorescence emission wavelength red-shifts from 492 nm to 687 nm, allowing it to be used to monitor changes in the H₂S concentration, while the double bond contained in **QCy7** can undergo twisted intramolecular charge transfer (TICT), resulting in a significant decrease in the fluorescence intensity, but this can be inhibited by a high-viscosity environment. Therefore, when the viscosity increased from 0.903 cp to 965 cp, the fluorescence intensity at 492 nm increased 110-fold. This enabled the monitoring of environmental viscosity. Using the ICT–TICT dual-response strategy, **Vis-H₂S** could simultaneously respond to mitochondrial viscosity and H₂S fluctuations. Thus, it can be used to explore the synergistic relationship between autophagy and the NFS1 protein.

Results and discussion

Design and synthesis of Vis-H₂S

As mentioned above, the "AND" logic gate-based "ICT–TICT dual-response mechanism" probe (**Vis-H**₂**S**) requires the coexistence of two analytes (viscosity and H₂S) and as such can monitor the synergistic relationship between autophagy and NFS1 protein. As shown in Scheme 1a and b, **Vis-H**₂**S** generated NIR fluorophores under the simultaneous activation by two agents, *i.e.*, H₂S and viscosity. In a low viscosity environment, weak NIR fluorescence of **Vis-H**₂**S** was observed in the presence



Scheme 1 Schematic illustration (a) and "AND" logic gate (b) of the dual-response platform; (c) the response mechanism of $Vis-H_2S$ toward H_2S and viscosity; (d) the sensing mechanism of H_2S using $Vis-H_2S$.

of H_2S , while in the presence of high viscosity (autophagyregulates mitochondrial viscosity) and the absence of H_2S , **Vis-H₂S** emits remarkable green fluorescence.

If, however, both high viscosity and H₂S are present, it emits strong near-infrared fluorescence. For normal cells, H₂S is present at a low level, and mitochondria have a high viscosity due to the presence of many proteins (resulting in strong green fluorescence), but H₂S and mitochondrial viscosity are easily destroyed by hypoxia and autophagy, respectively. Increased cell hypoxia can lead to the overexpression of reducing proteins such as NFS1 protein which results in a strong NIR fluorescence response. On the other hand, autophagy reduces the viscosity of the mitochondria, resulting in minimal fluorescence from either the red or green channel. As such, MDR produced by chemotherapy can lead to a simultaneous increase in autophagy and hypoxia, resulting in only weak NIR fluorescence from the system. The response mechanism is provided in Scheme 1c and d. Vis-H₂S, with a molecular rotor structure (two symmetric vinyl-coupled indolium moieties), enhanced the probe fluorescence in a high-viscosity environment due to restricted intramolecular rotation inhibiting nonradiative twisted intramolecular charge transfer (TICT¹⁴⁻¹⁶). The nucleophilic reaction of the ester with HS⁻ releases QCy7 in an anionic form, and the negative charge is delocalized to form quinone QCy7a or QCy7-b.

Vis-H₂S was synthesized according to Scheme S1,[†] and characterization data are provided in Fig. S1-S4.[†] The reaction mechanism between Vis-H₂S and H₂S was verified by measuring the relative molecular mass before and after the reaction using HR-MS. As shown in Fig. S5,† when Vis-H₂S was reacted with H_2S , a peak appeared at m/z = 461.2098, representing the reaction product and the peak at m/z = 346.1212 (representing the reactant) disappeared. We concurrently employed ¹H-NMR spectroscopy to conduct a mechanistic analysis of Vis-H₂S's response to H₂S. The results are illustrated in Fig. S6,[†] and exhibit significant chemical shift variations in the methyl hydrogens (H_a, H_b, H_c and H_d) on the indole moiety of the probe. Simultaneously, the signals corresponding to the hydrogens from the iodinated benzene disappear, substantiating the validity of the predicted mechanism underlying the response of Vis-H₂S's to H₂S. To further investigate the response of Vis-H₂S and QCy7 to viscosity, a PBS-glycerol solvent system was employed to monitor the spectral response of Vis-H₂S to viscosity changes (Fig. S7 and S8[†]). Vis-H₂S and QCy7 exhibited a nearly 110-fold and 22-fold increase in fluorescence, respectively, in response to the increase in solution viscosity from 0.903 cP to 965 cP. The relative fluorescence intensity of Vis-H₂S and QCy7 exhibited a linear relationship with viscosity ($R^2 =$ 0.985 for Vis-H₂S and $R^2 = 0.984$ for QCy7). The above results confirmed that Vis-H₂S and QCy7 exhibit excellent responses to viscosity and can be used to detect viscosity changes during in vivo and in vitro experiments. Additionally, we delved into the polarity responsiveness of QCy7 by examining its behavior in PBS-THF mixtures with different volume fractions (f_w) of PBS, spanning the range from 10% to 100%. As shown in Fig. S9,† there was minimal change in fluorescence intensity, indicating that QCv7 is not sensitive to polarity.

2-Iodobenzoate was used to regulate the "push-pull" electronic effects (intramolecular charge transfer, ICT) of Vis-H₂S so that it could specifically respond to changes in H₂S. We initially examined the absorption and fluorescence spectra of Vis-H₂S in response to H₂S. The results revealed that Vis-H₂S initially exhibited low levels of absorption and fluorescence intensity, but a significant increase was observed upon the addition of H₂S (Fig. S10a and b[†]). We also investigated the response of Vis- H_2S to H_2S in environments with different viscosities (PBS, pH = 7.4, containing 0%, 50%, and 70% glycerol). The ability of Vis- H_2S to simultaneously respond to H_2S and viscosity was also evaluated. As shown in Fig. S10a[†] and 1, upon increasing the H_2S concentration in the absence of glycerol, Vis- H_2S vs. H_2S exhibited a "turn-on" response, with an approximately 20-fold increase in the fluorescence intensity at 687 nm. Upon increasing the glycerol concentration, a ratiometric sign was found with increasing concentrations of H2S (Vis-H2S exhibited a decreasing intensity at $\lambda_{em} = 492$ nm and an increasing intensity at $\lambda_{\rm em}=$ 687 nm). Upon increasing the glycerol concentration, Igreen and Ired both increased significantly. The ratio signal also exhibited a good linear relationship at 50% and 70% glycerol when the H_2S concentration was lower than 75 μ M. From Fig. 1e and f the correlation coefficients were 0.982 and 0.989, respectively, and the limit of detection (LOD) values were calculated to be 45 nM and 50 nM, respectively, according to $3\sigma/$ k (where σ is the standard deviation of the blank (n = 11) and k is the slope for the range of linearity). The above experiments confirmed the dual-response of Vis-H₂S to viscosity and H₂S. Kinetic experiments indicated that when using 70% glycerol, the reaction between Vis-H₂S and H₂S (100 µM) reached a plateau within 60 min, indicating that Vis-H₂S could rapidly recognize H₂S (Fig. S10f[†]). Eighteen different interferents were selected to evaluate the selectivity of Vis-H₂S. As shown in Fig. S10g,† these interferents did not change the fluorescence intensity of Vis-H₂S in either a high-viscosity (glycerol) or lowviscosity (PBS) environment. These results clearly indicate that Vis-H₂S exhibits minimal interference and can be used for in situ imaging of cellular viscosity. Finally, we evaluated the effect of pH on the response of Vis-H₂S to H₂S. As shown in Fig. S10c,† for pH over a range from 4-8, there was no obvious change in the fluorescence signal before or after the simultaneous action of Vis-H₂S with H₂S and glycerol. In addition, the interference of cell viscosity stimulators (temperature and monensin17) on the fluorescence intensity of Vis-H₂S was investigated. The results in Fig. S10d[†] show that neither temperature nor monensin interfered with the fluorescence intensity of Vis-H₂S, which provides an excellent basis for subsequent cell experiments. According to the literature,18 acylphenol can serve as a recognition moiety for carboxylesterase-2, which is structurally similar to our recognition domain. To exclude the potential interference of carboxylesterase-2 in this study, we conducted photostability experiments of Vis-H2S in response to carboxylesterase-2, as shown in Fig. S10f.[†] The results indicate that over time, there was no change in fluorescence intensity, suggesting that carboxylesterase-2 has no interference with H₂S.

With the above results in hand, we evaluated the *in situ* imaging capability of **Vis-H₂S** for H₂S in cells. First, different



Fig. 1 Fluorescence intensity changes of Vis-H₂S (10 μ M) in response to H₂S and viscosity. (a) 0% glycerol, (b) 50% glycerol, and (c) 70% glycerol; fluorescence titration curves of the probes Vis-H₂S and H₂S (0–500 μ M); (d)–(f) the fluorescence intensity responses of Vis-H₂S for (a)–(c) to different concentrations of H₂S (0–500 μ M) at the ratio signal ($I_{687 nm}/I_{492 nm}$), respectively. The inset shows the linear relationship of the Vis-H₂S response to a low concentration of H₂S in different glycerol systems. The excitation wavelength was 400 nm, and the test medium was phosphate buffer (10 mM; pH = 7.4).

concentrations of Vis-H₂S were incubated with HepG2 cells for 24 h. The MTT assay results indicated that the cell viability remained above 80% and the IC_{50} was 63 $\mu\text{M},$ which indicates that Vis-H₂S was not cytotoxic and could be used for cell imaging (Fig. S11[†]). To accurately image autophagy and NFS1 protein simultaneously, the probe needs to exhibit mitochondrial-anchoring ability. According to the literature, as an organelle with a double-membrane structure, the negative charge of the inner mitochondrial membrane enables positively charged molecules to be specifically enriched and anchored in the mitochondria.¹⁹ Since the indole group of the Vis-H₂S probe is positively charged, the intracellular distribution should be targeted to the mitochondria. To this end, Vis-H₂S was mixed with a commercial mitochondrial localization dye, 100 nM Mito-Tracker Red, a commercial lysosome localization dye, 100 nM Lyso-Tracker Red, and a commercial endoplasmic reticulum localization dye, 100 nM ER-Tracker Red. These were then co-incubated with HepG2 cells for 30 min and imaged using confocal microscopy. As shown in Fig. S12[†], Vis-H₂S was mainly distributed in the mitochondria (Pearson colocalization coefficient was 0.94), while the Pearson colocalization coefficients in the lysosome and endoplasmic reticulum were 0.41 and 0.33, respectively, indicating that Vis-H₂S exhibits good mitochondrial localization and could detect the response to viscosity and H2S in the mitochondria during cellular experiments.

We then investigated the ability of **Vis-H₂S** (10 μ M) to respond to exogenous and endogenous H₂S and cell viscosity. The results are given in Fig. S13.† First cells were treated with sulfhydryl scavenger *N*-ethylmaleimide²⁰ (NEM, 0.5 mM) and then incubated with different concentrations of H₂S for 40 min. The experimental results indicated that upon increasing the NaHS (H₂S donor) concentration (0, 5, 10, or 20 μ M), I_{green} gradually decreased and I_{red} gradually increased, indicating that **Vis-H₂S** could detect exogenous H₂S in cells. Compared with the control group, the fluorescence intensity was significantly reduced for cells in the presence of NEM (0.5 mM). This was because of the removal of endogenous H₂S by NEM. Then, in the absence of H₂S due to the addition of NEM (reduced interference by H₂S), intracellular viscosity levels were regulated using monensin and different cell incubation temperatures (25 °C and 4 °C; a lower temperature increases the intracellular viscosity). As shown in Fig. S14,† upon decreasing the temperature and adding monensin, I_{green} increased significantly, while I_{red} was almost negligible. These experiments confirm that **Vis-H₂S** can readily respond to cell viscosity.

Next, we evaluated the dual-response of **Vis-H**₂**S** to H₂S and autophagy at the cellular level. Previous studies have shown that during autophagy, mitochondria are wrapped by vesicles to form autophagosomes that then fuse with lysosomes to form autolysosomes.²¹ This allows mitochondria and their contents to be degraded, resulting in reduced mitochondrial viscosity.²² As shown in Fig. S15,† compared with the control group and NEM group, after cells were treated with autophagy inducer rapamycin (100 nM, mTORC1 complex inhibitor²³), *I*_{green} was negligible, indicating that autophagy significantly reduced the mitochondrial viscosity. In addition, different concentrations of NaHS (5, 10, or 20 μ M) did not affect the *I*_{green}, but *I*_{red} gradually increased with an increase in H₂S concentration, indicating that **Vis-H₂S** could generate a dual response to autophagy and H₂S.

To further evaluate the ability of $Vis-H_2S$ to monitor autophagy and H_2S in real-time, confocal real-time imaging analysis was performed on HepG2 cells stained with $Vis-H_2S$ at different times and under cell culture conditions. The results are shown in Fig. 2a-e. The fluorescence of the red and green channels of the control group (HepG2 cells were untreated) remained almost constant for HepG2 cells loaded with Vis-H₂S with a 5 min interval for a total of 60 min, indicating that Vis-H₂S possesses good photostability. The cells were then starved to induce autophagy,²⁴ and the results indicated that both I_{green} and I_{red} were significantly reduced. From the above experimental results, a decrease in I_{green} occurred due to a decrease in mitochondrial viscosity during autophagy. In addition, the gradual decrease in I_{red} is attributed to the role of NFS1 protein as a reductive protein that facilitates the release of H₂S through desulfuration of L-cysteine. During the autophagy process, lysosomes degrade NFS1 protein within the mitochondria, resulting in a reduction of endogenous H₂S and subsequently weakening of $I_{\rm red}$. After the cells were starved and the autophagy inhibitor 3-methyladenine²⁵ (3-MA, 5 mM) was added, both Igreen and Ired were significantly enhanced and the fluorescence intensity remained almost unchanged for the next 60 min. This again shows that autophagy can simultaneously reduce mitochondrial viscosity and H2S levels. When cells were preincubated with NEM, regardless of whether the cells were incubated with 3-MA, changes in I_{green} were the same as those without NEM, while Ired was almost negligible. Due to the ability of flow cytometry to rapidly analyze a large number of individual cells, achieving high-throughput data collection is crucial for comprehensive assessment of fluorescence signal changes. In this regard, our study employed flow cytometry to investigate fluorescence intensity alterations under the aforementioned conditions, as illustrated in Fig. S16.[†] In comparison to the control group, cells subjected to starvation exhibited a significant decrease in fluorescence intensity in both the green and red channels, with the addition of 3-MA inhibiting this process. Concurrent starvation and NEM incubation resulted in a notable reduction in fluorescence intensity in both channels, with 3-MA incubation during this process leading to an increase in green channel fluorescence intensity and minimal interference with red channel fluorescence intensity. These findings align with real-time imaging results, confirming the dualresponsive capability of Vis-H₂S to simultaneously monitor cellular viscosity and H₂S levels.

Finally, we evaluated the source of MDR during tumor chemotherapy. Adriamycin, a widely used chemotherapeutic drug, works by consuming oxygen to produce ROS, which in turn triggers autophagy (ROS has been confirmed as the main cause of upregulating cellular autophagy) and overexpression of NFS1 protein due to increased hypoxia in tumor cells.²⁶ This



Fig. 2 Real-time fluorescence imaging images of the HepG2 cells in: (a) control group, (b) starvation treatment, (c) starvation treatment + 5 mM 3-MA, (d) starvation treatment + 0.5 mM NEM, and (e) starvation treatment + 5 mM 3-MA + 0.5 mM NEM incubated with Vis-H₂S (10 μ M). (f) Fluorescence trend graph of intracellular I_{green} as a function of time. (g) Fluorescence trend graph of intracellular I_{green} as a function of time. (g) Fluorescence trend graph of intracellular I_{green} as a function of time. Light collection range: green channel 450–520 nm, red channel 650–720 nm, the excitation light source is 552 nm, and the scale bar is 20 μ m.

Edge Article

combination of events leads to MDR but the synergistic effect between cellular autophagy and NFS1 protein overexpression in causing MDR has not been previously reported. To investigate the combined impact of autophagy and NFS1 protein on multidrug resistance (MDR) during tumor chemotherapy, we employed **Vis-H₂S** for imaging. Before conducting imaging experiments, we initially employed western blot analysis to assess the NFS1 protein levels. This step aimed to validate our hypothesis that the upregulation of NFS1 protein, induced by Adriamycin, leads to a substantial increase in H₂S production. The mechanism suggests that elevated NFS1 levels act as a protective response in tumor cells, preventing the cytotoxic effects of Adriamycin and thereby contributing to the development of multidrug resistance. The results are depicted in Fig. S18,† where the addition of Adriamycin led to a significant increase in NFS1 protein content, while NFS1KD rendered the NFS1 protein content negligible. As shown in Fig. 3a–f, 3h and i, the control group experiment indicates that **Vis-H₂S** retained excellent photostability (Fig. 3a). After adding Adriamycin, I_{green} was significantly reduced, while I_{red} increased first and then decreased (Fig. 3b). This occurred because the addition of Adriamycin significantly increased the overexpression of NFS1 protein. Then upon prolonged incubation time, autophagy decreased the mitochondrial viscosity resulting in decreased I_{red} . To validate this result, cells incubated with Adriamycin were also treated with the autophagy inhibitor 3-MA (10 μ M) or the NFS1 gene of HepG2 cells was eliminated (NFS1KD, operational approach according to the reported method²⁷). The results indicated that I_{green} gradually decreased and I_{red} increased gradually in cells pretreated with 3-MA (Fig. 3c). This



Fig. 3 (a–f) Real-time imaging fluorescence image of the HepG2 cells in (a) control group, (b) Adriamycin (5 μ M), (c) Adriamycin (5 μ M) + 3-MA (5 mM), (d) Adriamycin (5 μ M) after cysteine desulfurase (NFS1) gene knockout, (e) negative control of NFS1 and (f) Adriamycin (5 μ M) + 3-MA (5 mM) after NFS1 gene knockout. (g) Apoptosis assay of cells in (a–f); viable cells (FITC⁻/PI⁻), early apoptosis (FITC⁺/PI⁻), late apoptosis (FITC⁻/PI⁺), and necrosis (FITC⁺/PI⁺). The time-dependence of the relative fluorescence intensity of (h) I_{green} and (i) I_{red} . (j) MTT test of the HepG2 cells corresponding to (a–f). (k) Quantification data of the apoptosis assay results corresponding to (g).

confirms that the fluorescence intensity was only regulated by NFS1-induced H_2S . In the Adriamycin + NFS1KD group, I_{green} gradually decreased, and I_{red} was negligible, indicating that the fluorescence intensity was only regulated by autophagy. Compared with the Adriamycin group, the Adriamycin + NFS1KD group exhibited a more pronounced decrease in Igreen. We believe that the knockout of the NFS1 gene led to an increase in the cellular ROS level, which increased the autophagy level. In addition, the results of the negative control (performed according to the previous report²⁸) group were similar to those of the Adriamycin group, indicating that the NFS1KD group results were caused by the reduction of H₂S levels via the knockout of the NFS1 protein. Finally, when autophagy and NFS1 protein were simultaneously inhibited (as shown in Fig. 3f), unsurprisingly, Igreen remained almost unchanged, while Ired was negligible. We then further validated the accuracy of the results using flow cytometric analysis, as illustrated in Fig. S17.† The addition of Adriamycin resulted in a significant decrease in fluorescence intensity in both the green and red channels. Under these conditions, inhibiting autophagy (3-MA) led to a significant increase in red channel fluorescence intensity, while inhibiting NFS1 protein levels (NFS1KD) resulted in a significant decrease in red channel fluorescence intensity. This aligns with the real-time imaging results, demonstrating the probe's capability to detect cellular autophagy and NFS1 level fluctuations during the development of Adriamycin-induced drug resistance.

Having ascertained the autophagy and NFS1 protein levels under the given conditions, this study also employed Annexin V-FITC/PI apoptosis detection agent assay and the CCK8 assay to comprehensively evaluate the Adriamycin drug resistance under different conditions. As shown in Fig. 3g, 3j and k, the cells in the control group maintained a high survival rate, while the survival rate of the cells in the Adriamycin group decreased to 60.5%. Interestingly, when autophagy was inhibited or when the NFS1 protein was knocked out, the apoptosis rate was significantly increased to 49.6% and 31.1%, respectively. In addition, the lowest cell survival rate (9.7%) was achieved by the simultaneous inhibition of autophagy and knockdown of the NFS1 protein. The above experiments indicate that autophagy and the NFS1 protein are responsible for Adriamycin drug resistance.

Conclusions

A near-infrared fluorescence-based probe, Vis-H₂S, which could simultaneously respond to H₂S and viscosity, was constructed based on the QCy7 parent fluorophore. The probe could simultaneously monitor autophagy and H₂S during the treatment of cells with Adriamycin and further confirmed the direct relationship between autophagy and NFS1 protein overexpression with ROS-based chemotherapeutic drugs. The results indicated that Adriamycin up-regulated the degree of autophagy and the NFS1 protein levels during tumour treatment, which resulted in MDR. Furthermore, reducing the level of NFS1 protein results in a small increase in autophagy levels, which may be because the NFS1 protein is unable to fully clear the ROS. Hence, using probe Vis- H_2S we could confirm that both autophagy and NFS1 protein are closely related to the MDR of tumour cells. Therefore, Vis- H_2S is a powerful analytical tool for the in-depth exploration of the drug resistance mechanism of Adriamycin.

Data availability

All data supporting this study are provided as ESI[†] accompanying this paper.

Author contributions

Wei Hu: conceptualization, investigation, methodology, data curation, writing – original draft. Yifan He: conceptualization, data curation. Haixian Ren: conceptualization, supervision, funding acquisition. Li Chai: validation, data curation. Haiyan Li: methodology, data curation. Jianbin Chen: validation, funding acquisition. Chunya Li: supervision, validation, funding acquisition. Yanying Wang: supervision, validation. Tony D. James: conceptualization, validation, supervision, project administration, writing – review and editing.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

CYL wishes to thank the National Natural Science Foundation of China (No. 22074160 and 21874157), Fundamental Research Funds for the Central Universities of South-Central Minzu University (No. CZZ22003), Open Project Funding of Ministry of Education Key Laboratory for the Synthesis and Application of Organic Functional Molecules, Hubei University (No. KLSAOFM2305); HXR wishes to thank the National Natural Science Foundation of China (No. 22277104); JBC wishes to thank the National Natural Science Foundation of China (No. 22171154 and 22201152), the Special Funds of Taishan Scholar Project (No. tsqn202211206), the Youth Innovative Talents Recruitment and Cultivation Program of Shandong Higher Education, Jinan Science & Technology Bureau (No. 2021GXRC080). TDJ wishes to thank the University of Bath and the Open Research Fund of the School of Chemistry and Chemical Engineering, Henan Normal University (2020ZD01) for support.

References

 M. Tang, H. A. Abbas, M. V. Negrao, M. Ramineni, X. Hu, S. M. Hubert, J. Fujimoto, A. Reuben, S. Varghese, J. Zhang, J. Li, C. W. Chow, X. Mao, X. Song, W. C. Lee, J. Wu, L. Little, C. Gumbs, C. Behrens, C. Moran, A. Weissferdt, J. J. Lee, B. Sepesi, S. Swisher, C. Cheng, J. Kurie, D. Gibbons, J. V. Heymach, I. I. Wistuba, P. A. Futreal, N. Kalhor and J. Zhang, *Nat. Commun.*, 2021, 12, 7081.

- 2 R. Wang, X. Shu, H. Zhao, Q. Xue, C. Liu, A. Wu, F. Cheng, L. Wang, Y. Zhang, J. Feng, N. Wu and M. Li, *Nat. Commun.*, 2023, **14**, 2078.
- 3 K. Xu, L. Jin, L. Xu, Y. Zhu, L. Hong, C. Pan, Y. Li, J. Yao, R. Zou, W. Tang, J. Wang, A. Wu and W. Ren, *J. Nanobiotechnol.*, 2022, 20, 315.
- 4 M. Shah, S. Paradis, J. Betz, N. Beylis, R. Bharadwaj, T. Caceres, E. Gotuzzo, M. Joloba, V. Mave, L. Nakiyingi, M. P. Nicol, N. Pradhan, B. King, D. Armstrong, D. Knecht, C. E. Maus, C. K. Cooper, S. E. Dorman and Y. C. Manabe, *Clin. Infect. Dis.*, 2020, **71**, 1161–1167.
- 5 R. Chen, K. Qiu, G. Han, B. K. Kundu, G. Ding, Y. Sun and J. Diao, *Chem. Sci.*, 2023, **14**, 10236–10248.
- 6 V. Schroeder, S. Savagatrup, M. He, S. Lin and T. M. Swager, *Chem. Rev.*, 2019, **119**, 599–663.
- 7 K. Chanda and B. Mm, Chem. Soc. Rev., 2021, 50, 3706-3719.
- 8 P. R. Cheng, S. S. He, C. Z. Zhang, J. Liu and K. Y. Pu, *Angew*, 2023, **62**, e202301625.
- 9 Q. Jiang, M. Pan, J. Hu, J. Sun, L. Fan, Z. Zou, J. Wei, X. Yang and X. Liu, *Chem. Sci.*, 2020, **12**, 148–157.
- 10 J. Zhang, Y. Zhou, N. Li, W. T. Liu, J. Z. Liang, Y. Sun, W. X. Zhang, R. D. Fang, S. L. Huang, Z. H. Sun, Y. Wang and Q. Y. He, *Adv. Sci.*, 2020, 7, 2002306.
- 11 L. Hao, Z. W. Li, D. Y. Zhang, L. He, W. Liu, J. Yang, C. P. Tan, L. N. Ji and Z. W. Mao, *Chem. Sci.*, 2019, **10**, 1285–1293.
- K. M. Fujihara, B. Z. Zhang, T. D. Jackson, M. O. Ogunkola, B. Nijagal, J. V. Milne, D. A. Sallman, C. S. Ang, I. Nikolic, C. J. Kearney, S. J. Hogg, C. S. Cabalag, V. R. Sutton, S. Watt, A. T. Fujihara, J. A. Trapani, K. J. Simpson, D. Stojanovski, S. Leimkühler, S. Haupt, W. A. Phillips and N. J. Clemons, *Sci. Adv.*, 2022, 8, eabm9427.
- 13 H. D. Li, H. J. Kim, F. Xu, J. J. Han, Q. C. Yao, J. Y. Wang,
 K. Y. Pu, X. J. Peng and J. J. Yoon, *Chem. Soc. Rev.*, 2021,
 51, 1795–1835.
- 14 B. X. Shen, L. H. Liu, Y. B. Huang, J. C. Wu, H. Feng, Y. Liu, H. Huang and X. Zhang, *Aggregate*, 2023, 5, e421.
- 15 J. J. Chen, L. Q. Chen, Z. P. She, F. Zeng and S. Z. Wu, *Aggregate*, 2023, 5, e419.

- 16 B. X. Shen, K. H. Jung, S. T. Ye, C. A. Hoelzel, C. H. Wolstenholme, H. Huang, Y. Liu and X. Zhang, *Aggregate*, 2022, 4, e301.
- 17 L. Deng, L. Wojciech, C. W. Png, D. Y. Q. Kioh, Y. Gu, T. T. Aung, B. Malleret, E. C. Y. Chan, G. Peng, Y. Zhang, N. R. J. Gascoigne and K. S. W. Tan, *Theranostics*, 2023, 13, 1165–1179.
- 18 B. Yang, X. D. Ding, Z. M. Zhang, J. K. Li, S. Y. Fan, J. Y. Lai, R. Su, X. H. Wang and B. Wang, *Talanta*, 2024, 269, 125418.
- 19 Y. Su, H. Lin, Y. Tu, M. M. Wang, G. D. Zhang, J. Yang, H. K. Liu and Z. Su, *Chem. Sci.*, 2022, **13**, 1428–1439.
- 20 H. Lin, Y. Yu, L. Zhu, N. Lai, L. Zhang, Y. Guo, X. Lin, D. Yang, N. Ren, Z. Zhu and Q. Dong, *Redox Biol.*, 2023, 59, 102601.
- 21 H. J. Hwang, H. Ha, B. S. Lee, B. H. Kim, H. K. Song and Y. K. Kim, *Nat. Commun.*, 2022, **13**, 1436.
- 22 M. Tu, V. P. Tan, J. D. Yu, R. Tripathi, Z. Bigham, M. Barlow, J. M. Smith, J. H. Brown and S. Miyamoto, *Cell Death Differ.*, 2022, 29, 2472–2486.
- 23 C. Alquezar, K. M. Schoch, E. G. Geier, E. M. Ramos,
 A. Scrivo, K. H. Li, A. R. Argouarch, E. E. Mlynarski,
 B. Dombroski, M. DeTure, D. W. Dickson, J. S. Yokoyama,
 A. M. Cuervo, A. L. Burlingame, G. D. Schellenberg,
 T. M. Miller, B. L. Miller and A. W. Kao, *Sci. Adv.*, 2021, 7,
 eabg3897.
- 24 M. Kinnebrew, R. E. Woolley, T. B. Ansell, E. F. X. Byrne, S. Frigui, G. Luchetti, R. Sircar, S. Nachtergaele, L. Mydock-McGrane, K. Krishnan, S. Newstead, M. S. P. Sansom, D. F. Covey, C. Siebold and R. Rohatgi, *Sci. Adv.*, 2022, 8, eabm5563.
- 25 L. L. Zhou, Q. Guan, W. Zhou, J. L. Kan and Y. B. Dong, *Chem. Sci.*, 2022, **13**, 7846–7854.
- 26 Y. He, W. Hu, L. Chai, Y. Wang, X. Wang, T. Liang, H. Li and C. Li, *Chem. Commun.*, 2022, **58**, 7301–7304.
- 27 Y. Hu, J. You, M. Tong, X. Zheng, Z. Xu, X. Cheng and T. Jiang, *Adv. Sci.*, 2020, 7, 2000799.
- 28 D. Wu, F. Fei, Q. Zhang, X. Wang, Y. Gong, X. Chen, Y. Zheng, B. Tan, C. Xu, H. Xie, W. Fang, Z. Chen and Y. Wang, *Sci. Adv.*, 2022, 8, eabm3381.