ChemComm



View Article Online

COMMUNICATION



Anthony F. Hill* and Richard Y. Kong

Cite this: Chem. Commun., 2017, 53, 759

Received 9th November 2016, Accepted 10th December 2016

DOI: 10.1039/c6cc08946a

CrossMark

www.rsc.org/chemcomm

Bromination of the carbyne complexes $[W(\equiv CR)Br(CO)_2(dcpe)]$ (R = Ph, SiPh₃; dcpe = 1,2-bis(dicyclohexylphosphino)ethane) provides high oxidation state derivatives $[W(\equiv CPh)Br_3(dcpe)]$ and $[W(\equiv CBr)Br_3(dcpe)]$, the latter *via* an unprecedented bromodesilylation process.

In 1983 Lalor reported the first halocarbyne complexes $[M(\equiv CX)(CO)_2(Tp^*)]$ (M = Mo, W; X = Cl, Br; Tp* = hydrotris-(dimethylpyrazolyl)borate).¹ In the interim these have been proven to be remarkably versatile building blocks for the construction of elaborate and otherwise inaccessible carbyne derivatives via either nucleophilic substitution (spontaneous or Pd⁰-mediated)² or alternatively, lithium–halogen exchange to afford nucleophilic lithiocarbynes [M(=CLi)(CO)₂(Tp*)] (Scheme 1).³ Such protocols are however denied wider generality in that some 33 years later Lalor's complexes still remain the only known halocarbynes with the exception of Hughes' fluorocarbynes $[M(\equiv CF)(CO)_2(L)]$ $(M = Cr, Mo, W; L = \eta - C_5 R_5, R = H, Me)^4$ for which no subsequent reactivity has been reported. Extension of Lalor's synthetic approach involving the reaction of the carbonylates $[M(CO)_3(Tp^*)]^-$ with diazonium salts in haloform solvents, to any other ligand sets or metals has not met with success despite considerable effort.^{1b} We have therefore considered alternative approaches to the synthesis of $M \equiv C_1(\pm)$ synthons so as to broaden the applicability of the methodologies outlined in Scheme 1. In this pursuit, we have recently shown that fluoride-mediated protodesilyation of the silylcarbyne complex $[W(\equiv CSiPh_3)Br(CO)_2(dcpe)][(1a, \ddagger dcpe) =$ 1,2-bis(dicyclohexylphosphino)ethane] affords a rare example of a remarkably stable 'parent' methylidyne complex [W(=CH)Br- $(CO)_{2}(dcpe)$] (1b, Scheme 2).⁵

Roper first showed that it was possible to oxidise carbyne complexes without compromising the metal-carbon triple bond⁵ and this was further developed by Mayr who demonstrated the bromination of Fischer's carbyne $[W(\equiv CPh)Br(CO)_4]$ in dimethoxyethane (dme) to afford the Schrock-type carbyne $[W(\equiv CPh)Br_3(dme)]$.⁶ We have therefore explored the possibility of preparing a high-oxidation state§ silylcarbyne *fac*- $[W(\equiv CSiPh_3)Br_3(dcpe)]$ (2a) by bromination of 1a.

Prior to the pursuit of **2a**, we first validated the approach by brominating the new but more conventional benzylidyne complex $[W(\equiv CPh)Br(CO)_2(dcpe)]$ (**1c**, Scheme 2), obtained by a classical Fischer-Mayr oxide abstraction route.⁷ Thus treating $W(CO)_6$ sequentially with phenyl lithium, oxalyl bromide, 4-picoline and dcpe provides **1c** (see Fig. 1 and ESI† for characterisational data). Treating **1c** with bromine at low temperature results in the formation of the high-valent tungsten benzylidyne complex *fac*-[W(\equiv CPh)Br₃(dcpe)] (**2c**, 98%, Fig. 2). The ¹³C resonance for the benzylidyne appears at $\delta_C = 309.7$ (²*J*_{CP} = 13.4 Hz), moved some 46 ppm to lower frequency from that for **1c** ($\delta_C = 263.7$, ²*J*_{CP} = 9.3 Hz). The ³¹P resonance at 53.1 ppm shows satellite



Scheme 1 Umpolung of bromocarbyne reactivity (Nu = nucleophile, E = electrophile, M = Mo, W).

Research School of Chemistry, Australian National University, Canberra, Australian Capital Territory, ACT 2601, Australia. E-mail: a.hill@anu.edu.au

[†] Electronic supplementary information (ESI) available: Synthetic procedures, spectroscopic and crystallographic data, the latter in .cif format. CCDC 1515752– 1515754. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c6cc08946a





Fig. 1 Molecular structure of **1c** in a crystal (60% displacement ellipsoids, H-atoms, cyclohexyl groups simplified). Selected bond lengths (Å) and angles (°): W1–P1 2.5318(6), W1–P2 2.5407(6), W1–Br 2.6929(3), W1–C1 1.821(3), P1–W1–P2 80.38(2), C1–W1–P1 93.97(8), C1–W1–P2 97.55(8), C1–W1–Br1 171.46(8), C2–C1–W1 171.6(2).

resonances arising from coupling to 183 W ($^{1}J_{WP}$ = 212 Hz) which is comparable in magnitude to that observed for **1c** (225 Hz).

With the oxidative dibromination of **1c** to provide **2c** having proceeded without issue, the corresponding reaction with **1a** was explored. Addition of bromine to a solution of **1a** at -78 °C followed by slow warming to room temperature resulted in the formation of a khaki complex, the yield of which is optimised by employing two equivalents of Br₂. The product was formulated on the basis of spectroscopic (see ESI†) and structural data (Fig. 3) as the bromocarbyne complex [W(\equiv CBr)Br₃(dcpe)] (**2d**) resulting from both oxidation of the metal centre and electrophilic bromodesilylation of the silylcarbyne ligand (Scheme 2).



Fig. 2 Molecular structure of **2c** in a crystal (60% displacement ellipsoids, H-atoms and solvent omitted, cyclohexyl groups simplified, one of two crystallographically independent molecules shown). Selected bond lengths (Å) and angles (°): W1–Br3 2.7549(5), W1–Br4 2.4914(5), W1–Br6 2.4782(5), W1–P9 2.5786(11), W1–C23 1.789(5), C34–C23 W1 174.2(4).



Fig. 3 Molecular structure of 2d in a crystal of 2d·CH₂Cl₂ (60% displacement ellipsoids, H-atoms and solvent omitted, cyclohexyl groups simplified, one of two crystallographically independent molecules shown). Selected bond lengths (Å) and angles (°): W2–Br3 2.6918(6), W2–Br5 2.5016(7), W2–Br6 2.4800(7), W2–P3 2.5862(16), W2–P4 2.5734(16), W2–C1 1.793(6), Br1–C1 1.822(6), P4–W2–P3 76.35(5), W2–C1–Br1 171.5(4).

The key spectroscopic datum of interest is the triplet ¹³C resonance observed at $\delta_{\rm C}$ = 229.0 which shows coupling $(^{2}I_{CP} = 16.4 \text{ Hz})$ approximately twice the magnitude of those found for lower valent dcpe tungsten carbyne complexes. Within the low-valent regime, complexes of the form $[M(\equiv CR)(CO)_2(L)]$ $(M = Cr, Mo, W; L = HB(pzR_2-3,5)_3 R = H, Me)^8$ provide the most extensive range of comparative spectroscopic data. Of these, the bromocarbyne complexes have characteristically high frequency resonances (e.g., $[M(\equiv CBr)(CO)_2(Tp^*)]$: $\delta_C =$ 198.0 M = W; 202.5 M = Mo), exceeded only by the iodocarbyne $[W(\equiv CI)(CO)_2(Tp^*)]$ ($\delta_C = 183.2$).⁹ Thus the *ca.* 81 ppm shift to high frequency of the carbyne resonance for 2d relative to 2c is to be expected. The infrared spectrum included a comparatively strong absorption at 1260 cm⁻¹ which on the basis of simulation (DFT: M06-LACVP) for the simpler model $[W(\equiv CBr)Br_3(dmpe)]$ $(\nu_{\rm WC} = 1203 \text{ cm}^{-1})$ we assign to the primarily $\nu_{\rm WC}$ mode. No such

ChemComm

'pure' mode could be identified for the corresponding benzylidyne complex due to extensive coupling with phenyl ring modes. Infrared data for carbyne ligands are somewhat sparse in the literature and this may be traced, with good reason, to a number of factors. Firstly, they appear in the fingerprint region amongst co-ligand absorptions. The polarity of $M \equiv C$ bonds (and thus changes in dipole moments associated with IR intensities) can vary dramatically, hence the Fischer–Schrock dichotomy. Finally, and most importantly, in contrast to metal oxo and nitride ligands, for which IR data are invaluable and unambiguous, the $M \equiv C$ oscillator is invariably coupled to modes due to the substituent, a phenomenon that has been discussed in detail by Dao *et al.*¹⁰ These factors taken together mean that the value of ν_{MC} covers a wide range of both frequency and intensity such that reliable assignments are best made with recourse to simulation.

The molecular geometry of **2d** (Fig. 3) involves pseudooctahedral tungsten with a typically short W2–C1 bond length of 1.793(6) Å that is not however significantly different (1 e.s.d.) to that found in **2c**. Carbyne ligands typically exert a significant *trans* influence and this is reflected in the W2–Br3 bond length (2.6918(6) Å) being significantly longer than those *trans* to phosphorus (2.5016(7), 2.4800(7) Å), but less elongated relative to the corresponding bromide in **2c** (126 e.s.d.), suggesting that the bromocarbyne ligand, for which no previous structural data are available, has a pronounced *trans* influence, though less than that for more conventional alkylidynes. The C1–Br1 bond length of 1.822(6) Å is somewhat longer than any previously reported C(sp)–Br bonds which span the range 1.77–1.80 Å. The acute dcpe bite angle of 76.35(5)° allows the *cis*-bromo ligands to adopt a significantly obtuse angle (109.35(2)°).

The availability of both 2c and 2d provides an opportunity to compare these disparate carbyne substituents. Computational interrogation of the models $[W(\equiv CR)Br_3(dmpe)]$ (R = Ph, Br, DFT: MO6 LACVP) indicates that replacing the phenyl substituent in 2c with a bromide in 2d results in an increase in the negative (natural atomic) charge on the carbyne carbon (-0.29 to -0.51). The positive charge on tungsten is essentially unchanged for the benzylidyne (+0.75) and bromocarbyne (+0.78) models.

The stepwise mechanism by which 1a is converted to 2d would appear to involve initial oxidative decarbonylation of the tungsten followed by slower bromo-desilylation of the silylcarbyne. We assert this sequence based on the following observations. (i) CO evolution is immediately observed at low temperature; (ii) bromination of $[W(\equiv CPh)Br(CO)_2(dcpe)]$ (1c) cleanly affords the benzylidyne complex $[W(\equiv CPh)Br_3(dcpe)]$ (2c). As to the intimate mechanism for C-Si bond cleavage, in an isolobal context Isobe provided the protocol of choice for the electrophilic bromination of alkynylsilanes by N-bromosuccinimide to afford alkynyl bromides,¹¹ however a silver co-catalyst (AgNO₃, AgF) is required, implicating silver alkynyl intermediates. In the conversion of 1a to 2d however direct attack by Br₂ presumably affords transient [W(=CBrSiPh₃)Br₃(dcpe)]Br with subsequent attack at silicon by bromide. Electrophilic C-halogenation of carbyne ligands has been described previously, but only in one instance (chlorination of $[Os(\equiv CR)Cl(CO)(PPh_3)_2]$ to afford $[Os(=CClR)Cl_2(CO)(PPh_3)_2]$, R = C₆H₄Me-4)¹² and whilst bromocarbene complexes^{13,14} are rare and highly reactive due to the exceptional halide nucleofugacity, the complex $[Re_2(=CBrSiPh_3)(CO)_9]^{15}$ provides limited precedent.

In conclusion, a considerable number of silylcarbyne complexes are available *via* classical Fischer alkoxide/oxide abstraction,^{5,15,16} Schrock α -hydrogen elimination/abstraction¹⁷ or alkynylsilane metathesis¹⁸ protocols. The demonstration that halodesilylation may be achieved without rupture of the metal-carbon multiple bond augurs well for the elaboration of further synthetically valuable halocarbyne complexes to open the 33 year hiatus in their wider exploration.

Notes and references

[‡] Hereafter in compound numbers **a** = SiPh₃, **b** = H, **c** = Ph, **d** = Br. § Throughout, we employ the $[CR]^{3-}$ formalism for electron counting and the assignment of oxidations states and associated d-configurations. This is an arbitrary distinction in such highly covalent compounds and the alternative $[CR]^+$ formalism also has advocates.

- (a) T. Desmond, F. J. Lalor, G. Ferguson and M. Parvez, *Chem. Commun.*, 1983, 457–458; (b) T. J. Desmond, F. J. Lalor, G. Ferguson, M. Parvez and T. Wieckowski, *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.*, 1990, 46, 59–61; (c) F. J. Lalor, T. J. Desmond, G. M. Cotter, C. A. Shanahan, G. Ferguson, M. Parvez and B. Ruhl, *J. Chem. Soc., Dalton Trans.*, 1995, 1709–1726.
- 2 (a) T. Desmond, F. J. Lalor, G. Ferguson and M. Parvez, J. Chem. Soc., Chem. Commun., 1984, 75-77; (b) F. J. Lalor and S. A. O'Neill, J. Organomet. Chem., 2003, 684, 249-265; (c) S. Chaona, F. J. Lalor, G. Ferguson and M. M. Hunt, J. Chem. Soc., Chem. Commun., 1988, 1606-1608; (d) L. Weber, G. Dembeck, R. Boese and D. Blaeser, Organometallics, 1999, 18, 4603-4607; (e) L. Weber, G. Dembeck, Boese and D. Blaser, Chem. Ber., 1997, 130, 1305-1308; R. (f) B. E. Woodworth and J. L. Templeton, J. Am. Chem. Soc., 1996, 118, 7418-7419; (g) G. M. Jamison, P. S. White and J. L. Templeton, Organometallics, 1991, 10, 1954–1959; (h) M. Etienne, P. S. White and J. L. Templeton, J. Am. Chem. Soc., 1991, 113, 2324-2325; (i) M. I. Bruce, M. L. Cole, M. Gaudio, B. W. Skelton and A. H. White, J. Organomet. Chem., 2006, 691, 4601-4614; (j) D. J. Armitt, M. I. Bruce, M. Gaudio, N. N. Zaitseva, B. W. Skelton, A. H. White, B. Le Guennic, J.-F. Halet, M. A. Fox, R. L. Roberts, F. Hartl and P. J. Low, Dalton Trans., 2008, 6763-6775; (k) R. L. Cordiner, P. A. Gugger, A. F. Hill and A. C. Willis, Organometallics, 2009, 28, 6632-6635; (1) R. L. Cordiner, A. F. Hill and J. Wagler, Organometallics, 2008, 27, 4532-4540; (m) L. M. Caldwell, A. F. Hill, A. D. Rae and A. C. Willis, Organometallics, 2008, 27, 341-345; (n) L. M. Caldwell, A. F. Hill, R. Stranger, R. N. L. Terrett, K. M. von Nessi, J. S. Ward and A. C. Willis, Organometallics, 2015, 34, 328-334; (o) A. L. Colebatch and A. F. Hill, J. Am. Chem. Soc., 2014, 136, 17442-17445.
- 3 (a) R. L. Cordiner, A. F. Hill and J. Wagler, Organometallics, 2008, 27, 5177–5179; (b) A. L. Colebatch and A. F. Hill, Organometallics, 2016, 35, 2249–2255; (c) A. L. Colebatch, A. F. Hill and M. Sharma, Organometallics, 2015, 34, 2165–2182; (d) E. S. Borren, A. F. Hill, R. Shang, M. Sharma and A. C. Willis, J. Am. Chem. Soc., 2013, 135, 4942–4945; (e) A. F. Hill and R. Shang, Organometallics, 2012, 31, 4635–4638; (f) A. F. Hill, M. Sharma and A. C. Willis, Organometallics, 2012, 31, 2538–2542; (g) A. F. Hill, R. Shang and A. C. Willis, Organometallics, 2011, 30, 3237–3241; (h) A. L. Colebatch, A. F. Hill, R. Shang and A. C. Willis, Organometallics, 2010, 29, 6482–6487.
- 4 (a) H. Huang, R. P. Hughes, C. R. Landis and A. L. Rheingold, J. Am. Chem. Soc., 2006, 128, 7454–7455; (b) H. Huang, R. P. Hughes and A. L. Rheingold, Dalton Trans., 2011, 40, 47–55.
- 5 A. F. Hill, J. S. Ward and Y. Xiong, Organometallics, 2015, 34, 5057-5064.
- 6 (a) G. R. Clark, N. R. Edmonds, R. A. Pauptit, W. R. Roper, J. M. Waters and A. H. Wright, J. Organomet. Chem., 1983, 244, C57–C60; (b) W. R. Roper, J. Organomet. Chem., 1986, 300, 167–190.
- 7 A. Mayr and G. A. McDermott, J. Am. Chem. Soc., 1986, 108, 548-549.
- 8 L. M. Caldwell, Adv. Organomet. Chem., 2008, 56, 1-94.
- 9 A. E. Enriquez, P. S. White and J. L. Templeton, J. Am. Chem. Soc., 2001, 123, 4992-5002.

- 10 N. Q. Dao, E. O. Fischer and C. Kappenstein, *Nouv. J. Chim.*, 1980, 4, 85–94.
- 11 T. Nishikawa, S. Shibuya, S. Hosokawa and M. Isobe, *Synlett*, 1994, 485–486.
- 12 G. R. Clark, K. Marsden, W. R. Roper and L. J. Wright, J. Am. Chem. Soc., 1980, **102**, 6570–6571.
- 13 For a review of halomethyl and halomethylidene complexes see P. J. Brothers and W. R. Roper, *Chem. Rev.*, 1998, **88**, 1293-1326.
- 14 (a) T. G. Richmond, A. M. Crespi and D. F. Shriver, Organometallics, 1984, 3, 314–319; (b) A. M. Crespi and D. F. Shriver, Organometallics, 1985, 4, 1830–1835; (c) Y. Jiang, O. Blacque, T. Fox, C. M. Frech and H. Berke, Chem. Eur. J., 2010, 16, 2240–2249; (d) D. Mansuy, Pure Appl. Chem., 1982, 54, 638–639; (e) E. O. Fischer, W. Kleine and F. R. Kreißl, J. Organomet. Chem., 1976, 107, C23–C25.
- 15 E. O. Fischer, P. Rustemeyer and K. Ackermann, *Chem. Ber.*, 1982, 115, 3851–3859.
- 16 (a) E. O. Fischer, H. Hollfelder, P. Friedrich, F. R. Kreißl and G. Huttner, Angew. Chem., Int. Ed. Engl., 1977, 16, 401-402;
 (b) J. C. Jeffery, M. A. Ruiz and F. G. A. Stone, J. Organomet. Chem.,

1988, 355, 231–242; (*c*) G. M. Jamison, A. E. Bruce, P. S. White and J. L. Templeton, *J. Am. Chem. Soc.*, 1991, 113, 5057–5059; (*d*) R. L. Cordiner, A. F. Hill, R. Shang and A. C. Willis, *Organometallics*, 2011, **30**, 139–144.

- 17 (a) P. D. Savage, G. Wilkinson, M. Motevalli and M. B. Hursthouse, *Polyhedron*, 1987, 6, 1599–1601; (b) L. A. Morton, M. Miao, T. M. Callaway, T. Chen, S.-J. Chen, A. A. Tuinman, X. Yu, Z. Lu and Z.-L. Xue, *Chem. Commun.*, 2013, 49, 9555–9557; (c) L. Giannini, E. Solari, S. Dovesi, C. Floriani, N. Re, A. Chiesi-Villa and C. Rizzoli, *J. Am. Chem. Soc.*, 1999, 121, 2784–2796; (d) G. Balazs, M. Sierka and M. Scheer, *Angew. Chem., Int. Ed.*, 2005, 44, 4920–4924; (e) A. S. Hock and R. R. Schrock, *Chem. – Asian J.*, 2007, 2, 867–874; (f) S. W. Seidel, R. R. Schrock and W. M. Davis, *Organometallics*, 1998, 17, 1058–1068.
- 18 (a) M. L. Listemann and R. R. Schrock, Organometallics, 1985, 4, 74–83; (b) A. V. Safronova, L. N. Bochkarev, N. E. Stolyarova, I. K. Grigor'eva, I. P. Malysheva, G. V. Basova, G. K. Fukin, Y. A. Kursky, S. Y. Khorshev and G. A. Abakumov, Russ. Chem. Bull., 2003, 2140–2145.