



Cite this: *Chem. Commun.*, 2017, 53, 5432

DOI: 10.1039/c7cc90160g

rsc.li/chemcomm

Correction: Light up detection of heparin based on aggregation-induced emission and synergistic counter ion displacement

Shiwu Li,^a Meng Gao,^a Shuxia Wang,^b Rongrong Hu,^a Zujin Zhao,^a Anjun Qin^{*a} and Ben Zhong Tang^{*ac}

Correction for 'Light up detection of heparin based on aggregation-induced emission and synergistic counter ion displacement' by Shiwu Li *et al.*, *Chem. Commun.*, 2017, **53**, 4795–4798

The heparin structures shown in the graphical abstract, Fig. 1 and 6 were incorrect. The corrected versions of Fig. 1 and 6 are shown below and the graphical abstract has been updated.




Fig. 1 Illustration of the detection principle of HPQ-TBP-I for heparin.




Fig. 6 (A) Chemical structure of Hep, ChS and HA; (B) PL spectra of HPQ-TBP-I in aqueous solution treated with different biomolecules and ions (10 μ M), respectively; and (C) ratio of the relative PL intensity (I/I_0) at 501 nm with various biomolecules and ions. [PHB] = [FBS] = 0.05 mg mL⁻¹; [HPQ-TBP-I] = 100 μ M; λ_{ex} = 330 nm.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a State Key Laboratory of Luminescent Materials and Devices, South China University of Technology, Guangzhou 510640, China. E-mail: msqinaj@scut.edu.cn

^b Department of Nuclear Medicine, Guangdong General Hospital, Guangdong Academy of Medical Sciences, Guangzhou 510080, China

^c Department of Chemistry and Hong Kong Branch of Chinese National Engineering Research Center for Tissue Restoration and Reconstruction, The Hong Kong University of Science & Technology, Clear Water Bay, Kowloon, Hong Kong, China. E-mail: tangbenz@ust.hk

