

## CORRECTION

[View Article Online](#)  
[View Journal](#) | [View Issue](#)



Cite this: *Phys. Chem. Chem. Phys.*,  
2017, **19**, 9320

DOI: 10.1039/c7cp90057k

rsc.li/pccp

In the originally published version of our paper, the colors in the legend box of Fig. 1B were erroneously inverted. The correct version of Fig. 1 is represented below.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.



**Fig. 1** (A) Absorption spectra of the fully oxidized, ascorbate reduced and fully reduced form of *bc*<sub>1</sub>. (B) Difference spectra of the fully-reduced and ascorbate-reduced forms, corresponding to the reduction of the *b* hemes. Also shown is the 6-coordinate-minus 5-coordinate spectrum of *b*-hemes obtained from subtraction of the spectra of ferrous cytoglobin and ferrous myoglobin. The spectra are normalized at the 560 nm peak.

<sup>a</sup> LOB, Ecole Polytechnique, CNRS, INSERM, 91128 Palaiseau Cedex, France. E-mail: marten.vos@polytechnique.edu

<sup>b</sup> School of Biological Sciences, University of Essex, UK

<sup>c</sup> Department of Biology, University of Pennsylvania, Philadelphia, PA 19104, USA

