Dalton Transactions

COMMUNICATION

Check for updates

Cite this: Dalton Trans., 2017, 46, 4902

Received 1st March 2017, Accepted 7th March 2017 DOI: 10.1039/c7dt00748e

rsc li/dalton

NHC adducts of tantalum amidohalides: the first example of NHC abnormally coordinated to an early transition metal[†][‡]

Pavel A. Petrov, 💿 *^{a,b} Taisiya S. Sukhikh^a and Maxim N. Sokolov 💿 ^{a,b}

rare.10

The reaction of Ta(NMe₂)₅ with NHC·HBF₄ (NHC = IMes and SIMes) leads to new carbene adducts of the mixed Ta^V amidofluoride, [(NHC)TaF₃(NMe₂)₂]. On the contrary, the reaction of Ta(NMe₂)₅ with IMes·HCl gives a complex mixture, where [(aIMes) TaCl₂(NMe₂)₃] is identified as one of the products. This is the first example of abnormal NHC coordination to an early transition metal.

N-Heterocyclic carbenes (NHC) and their complexes have been attracting much attention in the last few decades, mainly due to the ease of their functionalization and, consequently, the tunability of their properties.¹ The complexes of late transition metals (mainly noble metals) with NHC are widely used in homogeneous catalysis and, in addition, NHC themselves serve as organocatalysts. The chemistry of NHC complexes of main group elements is also intensively developing, in particular, because NHC coordination stabilizes homoelement molecules, formally regarded as unstable "allotropes" (for example, the phosphorus clusters P_2 to P_{12})² and otherwise transitory species, such as SiCl2³ and BH,⁴ in the neutral, ionic or radical form.⁵ In contrast, the coordination chemistry of NHC with the early transition metals (TM) remains much in the shadow, particularly where the group 5 metals are concerned. This scarcity of data is clearly due to high oxophilicity of the hard Lewis acids (V, Nb and Ta), which is expected to disfavor the coordination of relatively soft carbene ligands. In addition, the commonly employed starting compounds both (for instance, halides) of the early TM and their NHC adducts are often extremely moisture-sensitive, particularly in higher oxidation states; however, this 'water hunger' can be satisfied

^aNikolaev Institute of Inorganic Chemistry SB RAS, Ak. Lavrentieva Av., 3, 630090 Novosibirsk, Russian Federation. E-mail: panah@niic.nsc.ru;

Fax: +7 (383)330 9489; Tel: +7 (383) 316 5845

^bNovosibirsk State University, Pirogova St., 2, 630090 Novosibirsk, Russian Federation

‡Electronic supplementary information (ESI) available: Experimental details, NMR spectra, and X-ray crystal data. CCDC 1530658–1530661. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7dt00748e



with partial hydrolysis, which leads to more stable oxohalides

the reactions of $[MX_5]_2$ (X = F, Cl, or Br) with free carbene,

either double salts $[M(NHC)_2Cl_4][MCl_6]$ or molecular species $[M(NHC)X_5]$ are formed. The main structural feature of these

Nb and Ta adducts is abnormal elongation of the M-carbene

bond and lack of π -contribution due to the absence of d elec-

trons in these d⁰ complexes. The electron deficiency of the co-

ordinated carbene is believed to be satisfied with weak

donation of electron density from a coordinated halide to the

carbene carbon atom.^{6,7c} In addition to the 'normal' coordi-

nation mode of NHC (A, Fig. 1), the 'abnormal' coordination

mode (*i.e.* via the C4 atom of the carbenazole cycle, \mathbf{B})⁹ is

known, but, to our knowledge, no complexes of early TM with

aNHC have been reported to date. The bimetallic complexes of

anionic dicarbenes (C) containing an early d-metal are also

routes to NHC adducts of early TM, using metal amides in

order to enable further functionalization via easy substitution

of the NR₂ groups. The addition of free IMes to a solution of

Ta(NMe₂)₅ in C₆D₆ does not lead to coordination, as is seen

from the ¹³C NMR shift of the carbene C atom (δ 219.6 ppm)

(ESI, Fig. S2[‡]). This is very likely due to both the high steric

demands and strong π -donor nature of the NMe₂ groups,

which decrease the acceptor strength of the Ta atom.

Therefore, we tried the reaction of Ta(NMe₂)₅ with the proto-

nated carbene precursor IMes·HBF₄, which was supposed to

Given this scarcity of results we set out to explore new

The first examples of monodentate NHC adducts with Nb and Ta halides⁷ and oxohalides⁸ were very recently reported. In

that sometimes can be isolated as NHC adducts.6

Fig. 1 Bonding modes of N-heterocyclic carbenes.



View Article Online View Journal | View Issue

[†]Dedicated to Prof. Peter W. Roesky on the occasion of his 50th birthday.

lead to the elimination of one equivalent of HNMe2 and simultaneous coordination of an NHC molecule. Much to our surprise, the reaction resulted in the formation of a mixed amidofluoride complex $[(IMes)TaF_3(NMe_2)_2]$ (1a), which was isolated after the appropriate workup in moderate yield as large colorless crystalline blocks (Scheme 1). 1a indeed contains a coordinated IMes, and, in addition, three F atoms (obviously derived from the BF_4^{-} anion) are coordinated to the Ta atom, which illustrates the enormous affinity of tantalum(v) for fluoride. The isostructural complex $[(SIMe_3)TaF_3(NMe_2)_2]$ (1b) with a saturated carbene was prepared in a similar way from $Ta(NMe_2)_5$ and SIMes·HBF₄. Mixed amidofluoride complexes are commonly synthesized by ligand metathesis between a metal amide and fluoride in the desired ratio, and the only known NHC adduct of a transition metal amidofluoride, $[(TiF_2(NEt_2)$ $(NHC)_2(\mu$ -F)₂], was isolated following precisely this pathway.¹¹

Both 1a and 1b adopt the mer, cis ligand arrangement with all the fluoride ligands being in cis positions relative to NHC (Fig. 2). The Ta-C_{carbene} bond length in 1a (2.389(5) Å) is comparable to that in [IPrTaCl₅] $(2.373(5) \text{ Å})^{7c}$ and [IPrTaBr₅] $(2.383(10) \text{ Å})^{7b}$ and is slightly longer than that found in the *trans*-dicarbene cation $[(IMes)_2 TaCl_4]^+$ (2.346(6) Å).^{7c} The Ta–N lengths cis and trans to NHC are almost equal, which suggests that the carbene is not involved in π -bonding with Ta. The fluorides are oriented toward the NHC ligand (the Ccarbene-Ta-F angles fall within the range of 79.77(15)-82.65(15)°), but this is not the case with the *cis*-NMe₂ group where the C(1)-Ta-N(3) angle is 98.65(15)°. The C…F distances between the cis fluorides and the C_{carbene} atom (2.852(6), 2.879(6), and 2.781(6) Å, for F(1), F(2) and F(3), respectively) are less than the sum of the van der Waals radii for carbon and fluorine (3.17 Å),¹² which is a situation that is typical for reported NHC adducts of the early TM halides.^{7,13} The imidazoline cycle is slightly (by 13.5°) turned away from the C(1)F(2)N(3)N(4) plane. The NHC ligand is also slightly tilted away from the cis-NMe₂ ligand (the C(1)-Ta(1)-N(3) angle is 98.65(18)°), clearly to ease steric repulsion.

Complex **1b** retains the major structural features of its congener **1a**. The main difference is that the $Ta-C_{carbene}$ length (2.432(2) Å) is *ca.* 0.04 Å longer than that in **1a**. The reason for this elongation is unclear; in imido alkylidene complexes [(NHC)Mo^{VI}(=CHR)(=NAr)], the difference in the Mo-C



Fig. 2 Molecular structure of 1a. Hydrogen atoms have been omitted for clarity. Thermal ellipsoids are drawn at 30% probability. Selected bond lengths (Å) and angles (°) are given for 1a and the isostructural complex 1b. For 1a: Ta1-C1 2.389(5), Ta1-F1 1.898(3), Ta1-F2 1.941(3), Ta1-F3 1.911(3), Ta1-N3 1.987(4), Ta1-N4 1.991(4), C1-Ta1-N3 98.65(18), and C1-Ta1-N4 167.03(17). For 1b: Ta1-C1 2.432(2), Ta1-F1 1.9127(12), Ta1-F2 1.9398(12), Ta1-F3 1.9114(12), Ta1-N3 2.0017(18), Ta1-N4 1.9861(18), C1-Ta1-N3 99.70(7), and C1-Ta1-N4 167.22(7).

lengths for IMes and SIMes lies within the 0.01 Å range.¹⁴ The Ir–C bond lengths in the $[(NHC)Ir(CO)_2Cl]$ and [(NHC)Ir(cod) Cl] systems (cod = 1,5-cyclooctadiene) also insignificantly depend on the saturation of the NHC backbone.¹⁵

The NMR spectra of **1a** and **1b** in C_6D_6 (ESI, Fig. S3–S8‡) are consistent with the solid state structures. The ¹⁹F NMR spectra show two signals in a 1:2 ratio corresponding to a fluoride *trans* to the NMe₂ group and to a pair of mutually *trans* fluorides. The ¹³C{¹H} NMR spectra show well-resolved quartets at 194.9 (**1a**) and 217.0 (**1b**) ppm due to spin–spin coupling with the three nearly equivalent F atoms *cis* to carbene. The signals of the carbene C atom are shifted *ca*. 25 ppm downfield from the free IMes (219.7)¹⁶ and SIMes (243.8),¹⁷ as is common for the early TM complexes.

In an unexpected contrast to the rather smooth formation of **1a** and **b**, the reaction of $Ta(NMe_2)_5$ with IMes·HCl in a 1:1 molar ratio proceeds in a much more complicated manner. After heating in THF and evaporating all the volatiles, the reaction mixture was consecutively extracted with pentane and toluene. Slow evaporation of the pentane extract afforded a complex mixture of crystals, in which only free IMes and the



Scheme 1

novel complex [IMes–H···IMes][*fac*-TaCl₃(NMe₂)₃] (2) could be identified (ESI, Fig. S1[‡]). The cation in 2 contains an IMes molecule bound with a hydrogen bond to the parent imidazo-lium cation IMesH⁺, which retains the common structural features of such adducts.¹⁸ The isolation of free IMes and 2 shows severe competition between the coordination of the softer NHC (primarily σ -donor) and harder chloride and amide anions (π -donors) to the hard d⁰ Ta^V center, and thus one can deduce that even H-bond formation with a C-H moiety rivals the coordination of NHC to Ta^V.

Slow evaporation of the toluene extract afforded a brown oil with a few yellow needles, which, according to X-ray analysis, turns out to be $[(aIMes)TaCl_2(NMe_2)_3]$ (3), the Ta complex featuring the abnormal coordination of the NHC ligand. To the best of our knowledge, including a search in the Cambridge Structural Database,¹⁹ this is the first example of this coordination mode of a neutral NHC to early transition metals.

The structure of 3 (Fig. 3) represents a distorted octahedral complex with the mer, cis ligand arrangement. The Ta-N length trans to NHC (2.0165(18) Å) is longer than the two Ta-N lengths trans to Cl (1.9760(18) and 1.9710(17) Å). The Ta- $C_{carbene}$ length (2.339(2) Å) is appreciably shorter than that in both 1a and $[TaCl_5(nIPr)]$ (2.373(5) Å),^{7c} even taking into account the much greater steric hindrance caused by the NMe₂ groups. Both the elongated Ta-N and shortened Ta-C lengths might reflect the enhanced donor strength of aNHC compared to the 'normal' coordination mode.⁹ The pairs of complexes that differ exclusively in the coordination manner of NHC are still scarce; however, in complexes that simultaneously contain normally and abnormally coordinated carbene, the M-C_{aNHC} length can be shorter, equal or even slightly longer than the M-C_{nNHC} length.²⁰ Similarly to the nNHC complexes, the cischlorides in 3 form short contacts with the C_{carbene} atom



Fig. 3 Molecular structure of 3, omitting hydrogen atoms except for those at C1 and C3. Thermal ellipsoids are drawn at 30% probability. Selected bond lengths (Å) and angles (°): Ta1–C2 2.339(2), Ta1–Cl1 2.5081(6), Ta1–Cl2 2.5546(5), Ta1–N3 2.0165(18), Ta1–N4 1.9760(18), Ta1–N5 1.9710(17), C2–Ta1–N3 162.50(7), C2–Ta1–N4 97.46(7), and C2–Ta1–N5 90.05(7).

(C(2)-Cl(1) 3.042(4), C(2)-Cl(2) 3.242(2) Å, C(2)-Ta(1)-Cl(1) 77.66(5), and C(2)-Ta(1)-Cl(2) 82.84(5)°), falling well within the sum of the van der Waals radii for C and Cl (3.45 Å).

Elucidation of the structure of 3 enabled us to develop its rational synthesis *via* the cleavage of the dimer $[TaCl_2(NMe_2)_3]_2$ with free NHC. Reaction with IMes (1 : 2 molar ratio) in toluene afforded 3 in good yield as a yellow powder, which was identified by IR spectra and cell parameters. One can conclude that chloride ligands enhance the Lewis acidity of the metal center in TaCl₂(NMe₂)₃ compared to Ta(NMe₂)₅; therefore, it forms a dimer that can be cleaved by NHC.

Recently Dagorne et al. reported a pathway to aNHC adducts of the sterically demanding ItBu carbene and group 13 metals, namely, isomerization of $[(nItBu)MX_3]$ (M = Al, Ga, and In; X = Hal, and alkyl), induced by heating or the addition of a polar solvent (THF).²¹ In contrast, complexes of the less bulky carbenes IPr or IMes did not isomerize. However, with the more crowded precursor, Ga(CH₂SiMe₃)₃, IPr shows fast nNHC to aNHC conversion, whereas ItBu forms only the aNHC complex but not the *n*NHC complex.²² Heating of [(*n*IPr)Fe $(N(SiMe_3)_2)_2$ also leads to isomerization to the *a*IPr complex, whereas $[(nItBu)Fe(N(SiMe_3)_2)_2]$ decomposes upon heating to give [(tBuIm)₂Fe(N(SiMe₃)₂)₂] (tBuIm - N-tert-butylimidazole), $Fe(N(SiMe_3)_2)_2$ and isobutene.²³ In our case, complex 3 is formed under mild conditions (toluene, rt), implying that the steric hindrance around the Ta center is the main driving force for the aNHC coordination. A solution of the less sterically hindered 1a in C₆D₆ remained intact upon heating at 70 °C for 5 days, showing no changes in its NMR spectra.

Unlike 1a and b, the crystals of 3 are almost insoluble in hydrocarbons. The NMR spectrum of a freshly prepared solution in THF-d₈ shows two sets of non-equivalent Mes resonances and two doublets of the imidazoline ring protons at 7.88 and 8.40 ppm, which are characteristic of aNHC complexes.[‡] Two more pairs of such doublets of lower intensity appear immediately after sample preparation, very likely due to the protonation of an NMe₂ group with traces of water. ${}^{13}C{}^{1}H$ NMR shows the C_{carbene} resonance at 186.9 ppm. After several days, however, extensive protonation of the NHC moiety becomes apparent, and complex 3 decomposes almost completely in 2 weeks. This process is accompanied by a gradual increase in the signal at 3.60 ppm, which corresponds to the protons entering the 2,5-positions of THF- d_8 and indicates participation of the solvent in the decomposition. The activation of THF-d₈ with an aNHC complex was reported recently by Goicoechea et al.²⁴ In addition, the HC=CH backbone of imidazoline-base NHC is known to be liable for the H/D exchange.25

In CD_2Cl_2 3 behaves similarly. Immediately after preparation, its ¹H NMR spectrum features characteristic C²H and C⁵H doublets at 7.90 and 7.74 ppm, respectively, along with minor signals belonging to another *a*NHC complex yet to be identified, and at least one additional Mes-containing species. The C_{carbene} atom of 3 resonates at 187.2 ppm. However, both 3 and the other *a*NHC species gradually evolve into a complex mixture of yet unidentified species, demonstrating the above-

mentioned sensitivity of early TM NHC complexes towards solvolysis. Complicated solution dynamics is characteristic of early transition metal amidohalide complexes: the solution of the parent $[Ta(NMe_2)_3Cl_2]_2$ in toluene is unstable towards ligand redistribution and converts into a mixture of $[Ta(NMe_2)_nCl_{5-n}]$ and other unidentified products.²⁶

To conclude, Ta(NMe₂)₅ and [TaCl₂(NMe₂)₃]₂ prove to be excellent precursors for the preparation of Ta^V complexes with NHC. The first complex of an early transition metal with *a*NHC has been prepared taking advantage of the steric constraints caused by coordinated dialkylamido groups. Novel, simple pathway towards Ta mixed amidofluorides through BF₄⁻ salts was found, and the first NHC adducts of Ta^V amidofluoride were isolated. Further studies are underway to investigate the reactivity of this compound.

This study was supported by the Russian Science Foundation (project 14-13-01053).

Notes and references

- 1 M. N. Hopkinson, C. Richter, M. Schedler and F. Glorius, *Nature*, 2014, **510**, 485–496.
- 2 (a) J. D. Masuda, W. W. Schoeller, B. Donnadieu and G. Bertrand, Angew. Chem., Int. Ed., 2007, 46, 7052-7055;
 (b) J. D. Masuda, W. W. Schoeller, B. Donnadieu and G. Bertrand, J. Am. Chem. Soc., 2007, 129, 14180-14181;
 (c) Y. Wang, Y. Xie, P. Wei, R. B. King, H. F. Schaefer, P. v. R. Schleyer and G. H. Robinson, J. Am. Chem. Soc., 2008, 130, 14970-14971; (d) O. Back, G. Kuchenbeiser, B. Donnadieu and G. Bertrand, Angew. Chem., Int. Ed., 2009, 48, 5530-5533; (e) C. L. Dorsey, B. M. Squires and T. W. Hudnall, Angew. Chem., Int. Ed., 2013, 52, 4462-4465;
 (f) C. H. Martin, C. M. Weinstein, C. E. Moore, A. L. Rheingold and G. Bertrand, Chem. Commun., 2013, 49, 4486-4488.
- 3 R. S. Ghadwal, R. Azhakar and H. W. Roesky, *Acc. Chem. Res.*, 2013, **46**, 444–456.
- 4 (a) R. Kinjo, B. Donnadieu, M. A. Celik, G. Frenking and G. Bertrand, *Science*, 2011, **333**, 610–613; (b) D. A. Ruiz, M. Melaimi and G. Bertrand, *Chem. Commun.*, 2014, **50**, 7837–7839.
- 5 (a) T. W. Hudnall, R. A. Ugarte and T. A. Perera, in N-Heterocyclic Carbenes. From Laboratory Curiosities to Efficient Synthetic Tools, ed. S. Díez-González, RSC Publishing, 2nd edn, 2017, ch. 5, pp. 178–237; (b) Y. Wang and G. H. Robinson, Inorg. Chem., 2011, 50, 12326–12337.
- 6 (a) C. D. Abernethy, G. M. Codd, M. D. Spicer and M. K. Taylor, *J. Am. Chem. Soc.*, 2003, 125, 1128–1129;
 (b) C. A. Dodds, M. D. Spicer and T. Tuttle, *Organometallics*, 2011, 30, 6262–6269.
- 7 (a) M. Bortoluzzi, E. Ferretti, F. Marchetti, G. Pampaloni and S. Zacchini, *Chem. Commun.*, 2014, 50, 4472-4474;
 (b) M. Bortoluzzi, E. Ferretti, F. Marchetti, G. Pampaloni and S. Zacchini, *Dalton Trans.*, 2016, 45, 6939-6948;
 (c) Z. Wei, W. Zhang, G. Luo, F. Xu, Y. Mei and H. Cai, *New*

J. Chem., 2016, **40**, 6270–6275; (d) D. Pugh, J. A. Wright, S. Freeman and A. A. Danopoulos, *Dalton Trans.*, 2006, 775–782.

- 8 (a) M. Bortoluzzi, E. Ferretti, F. Marchetti, G. Pampaloni,
 C. Pinzino and S. Zacchini, *Inorg. Chem.*, 2016, 55, 4173-4182; (b) M. Bortoluzzi, E. Ferretti, F. Marchetti,
 G. Pampaloni and S. Zacchini, *J. Coord. Chem.*, 2016, 69, 2766-2774.
- 9 This coordination type is referred to as abnormal or mesoionic: (a) R. H. Crabtree, *Coord. Chem. Rev.*, 2013, 257, 755–766; (b) P. L. Arnold and S. Pearson, *Coord. Chem. Rev.*, 2007, 251, 596–609.
- 10 For heterometallic Y/K and Sm/K complexes see:
 P. L. Arnold and S. T. Liddle, *Organometallics*, 2006, 25, 1485–1491. For heterometallic Mn/Al complex see:
 R. A. Musgrave, R. S. P. Turbervill, M. Irwin and
 J. M. Goicoechea, *Angew. Chem., Int. Ed.*, 2012, 51, 10832–10835.
- 11 G. B. Nikiforov, H. W. Roesky, P. G. Jones, J. Magull, A. Ringe and R. B. Oswald, *Inorg. Chem.*, 2008, 47, 2171– 2179.
- 12 A. Bondi, J. Phys. Chem., 1964, 68, 441-451.
- 13 C. D. Abernethy, G. M. Codd, M. D. Spicer and M. K. Taylor, *J. Am. Chem. Soc.*, 2003, **125**, 1128– 1129.
- S. Sen, R. Schowner, D. A. Imbrich, W. Frey, M. Hunger and M. R. Buchmeiser, *Chem. – Eur. J.*, 2015, 21, 13778– 13788.
- 15 R. A. Kelly, H. Clavier, S. Giudice, N. M. Scott, E. D. Stevens, J. Bordner, I. Samardjiev, C. D. Hoff, L. Cavallo and S. P. Nolan, *Organometallics*, 2008, 27, 202– 210.
- 16 A. J. Arduengo, H. V. R. Dias, R. L. Harlow and M. Kline, J. Am. Chem. Soc., 1992, 114, 5530–5534.
- 17 A. J. Arduengo, R. Krafczyk, R. Schmutzler, H. A. Craig, J. R. Goerlich, W. J. Marshall and M. Unverzagt, *Tetrahedron*, 1999, 55, 14523–14534.
- 18 (a) A. J. Arduengo, S. F. Gamper, M. Tamm, J. C. Calabrese, F. Davidson and H. A. Craig, *J. Am. Chem. Soc.*, 1995, 117, 572–573; (b) C. Jones, D. P. Mills and R. P. Rose, *J. Organomet. Chem.*, 2006, 691, 3060–3064; (c) M. A. Schmidt, P. Müller and M. Movassaghi, *Tetrahedron Lett.*, 2008, 49, 4316–4318.
- 19 CSD v.5.36, updated May 2016.
- 20 (a) H. Lebel, M. K. Janes, A. B. Charette and S. P. Nolan, J. Am. Chem. Soc., 2004, 126, 5046–5047; (b) L.-C. Campeau, P. Thansandote and K. Fagnou, Org. Lett., 2005, 7, 1857– 1860; (c) G. C. Fortman, N. M. Scott, A. Linden, E. D. Stevens, R. Dorta and S. P. Nolan, Chem. Commun., 2010, 46, 1050–1052; (d) M. S. Varonka and T. H. Warren, Organometallics, 2010, 29, 717–720.
- 21 (a) A.-L. Schmitt, G. Schnee, R. Welter and S. Dagorne, *Chem. Commun.*, 2010, 46, 2480–2482; (b) G. Schnee, O. N. Faza, D. Specklin, B. Jacques, L. Karmazin, R. Welter, C. Silva López and S. Dagorne, *Chem. – Eur. J.*, 2015, 21, 17959–17972.

- 22 M. Uzelac, A. Hernán-Gómez, D. R. Armstrong, A. R. Kennedy and E. Hevia, *Chem. Sci.*, 2015, 6, 5719– 5728.
- 23 B. M. Day, T. Pugh, D. Hendriks, C. Fonseca Guerra,
 D. J. Evans, F. M. Bickelhaupt and R. A. Layfield, *J. Am. Chem. Soc.*, 2013, 135, 13338–13341.
- 24 R. S. P. Turbervill and J. M. Goicoechea, Aust. J. Chem., 2013, 66, 1131–1137.
- 25 M. K. Denk and J. M. Rodezno, *J. Organomet. Chem.*, 2001, 617–618, 737–740.
- 26 M. H. Chisholm, J. C. Huffman and L.-S. Tan, *Inorg. Chem.*, 1981, **20**, 1859–1866.