CORRECTION

[View Article Online](#)
[View Journal](#) | [View Issue](#) Cite this: *Dalton Trans.*, 2017, **46**, 11970

Correction: Enhancements in catalytic reactivity and selectivity of homobimetallic complexes containing heteroditopic ligands

Mark R. D. Gatus,^{a,b} Roy T. McBurney,^b Mohan Bhadbhade^c and Barbara A. Messerle^{*a,b}

DOI: 10.1039/c7dt90117h

rsc.li/daltonCorrection for 'Enhancements in catalytic reactivity and selectivity of homobimetallic complexes containing heteroditopic ligands' by Mark R. D. Gatus *et al.*, *Dalton Trans.*, 2017, **46**, 7457–7466.

In the original version of Table 2, the fourth column entitled 'Time/min (% conv.)' under the 'hydroamination step' was mislabelled; the correct column title is 'Time/h (% conv.)'. There is no text referencing the conversion times of the hydroamination step within the discussion that affects this change. This has been corrected in the table below.

Table 2 Comparison of Rh(I) and Ir(I) homobimetallic complexes **12** and **13** with Rh(I) and Ir(I) monometallic complexes **16–18** for the sequential two-step hydroamination–hydrosilylation reactions of **19a** and **20a**

Hydroamination step					Hydrosilylation step ^d	
Entry	Cat.	Substrate	Time/h (% conv.)	TOF ^e (h ^{−1})	Time/min (% conv.)	TOF ^e (h ^{−1})
1 ^a		19a	0.4 (>95)	2118	47.0 (94)	642
2 ^a		20a	4.6 (>95)	173	17.0 (>98)	546
3 ^a		19a	0.1 (>95)	2913	0.2 (>98)	>16 200
4 ^a		20a	1.5 (>95)	432	0.6 (>98)	>5100
5 ^b		19a	0.04 (>95)	3983	1.0 (>98)	>1365
6 ^b		20a	3.6 (>95)	62	1.6 (>98)	>1694
7 ^b		19a	0.4 (>95)	1726	6.0 (>98)	459
8 ^b		20a	4.9 (>95)	53	18.0 (>98)	199
9 ^c		19a	0.6 (>95)	436	3.6 (>98)	365
10 ^c		20a	5.0 (>95)	140	13.2 (>98)	212

^a Conditions – 1.0 mol% cat, 2 mol% NaBAR^F₄, 2 × 10^{−4} mol substrate, toluene-*d*₈, 100 °C. ^b Conditions – 2.0 mol% cat, 2 × 10^{−4} mol substrate, toluene-*d*₈, 100 °C. ^c Conditions – 2.0 mol% cat, 2.0 mol% NaBAR^F₄, 2 × 10^{−4} mol substrate, toluene-*d*₈, 100 °C. ^d 4 × 10^{−4} mol of Ph₂SiH₂ was added to the mixture. ^e TOFs (Turn Over Frequency) were calculated at the point of 50% conversion of substrate to product(s).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aSchool of Chemistry, University of New South Wales, Sydney, NSW 2052, Australia^bDepartment of Chemistry and Biomolecular Sciences, Macquarie University, Sydney 2019, NSW, Australia. E-mail: barbara.messerle@mq.edu.au^cMark Wainwright Analytical Centre, University of New South Wales, Sydney, NSW 2052, Australia