## Energy & Environmental Science



## CORRECTION

**View Article Online** 



Cite this: Energy Environ. Sci., 2017, 10, 2042

DOI: 10.1039/c7ee90049i

rsc.li/ees

## Correction: Mono- and tri-ester hydrogenolysis using tandem catalysis. Scope and mechanism

Tracv L. Lohr, a Zhi Li, a Raieev S. Assarv, b Larry A. Curtiss bc and Tobin J. Marks\*a

Correction for 'Mono- and tri-ester hydrogenolysis using tandem catalysis. Scope and mechanism' by Tracy L. Lohr et al., Energy Environ. Sci., 2016, 9, 550-564.

The authors wish to correct Fig. 4 and associated discussion on p. 555 of the manuscript. The authors discovered a small Excel error in the calculation of Nt. We have therefore replaced Fig. 4 with a new graph that plots % conversion after 1 h. This correction does not affect any other data or change any of the scientific conclusions in the article.

The paragraph on p. 555 should read as follows:

"Substituent effects on the acyl and alkoxy group were also investigated (Fig. 4 and 5). Sequentially replacing the methyl hydrogen atoms of cyclohexyl acetate with methyl groups (R = Me < Et < <sup>i</sup>Pr < <sup>t</sup>Bu; Fig. 4, blue columns) monotonically depresses the conversion after 1 h, under identical reaction conditions, from 89% (Me) to 33% (Bu). Note however that replacing an acyl H with a single electron-withdrawing chloro group increases the conversion to 93% (CH<sub>2</sub>Cl), while including a second chloro group (CHCl<sub>2</sub>) depresses the conversion to 61%. Fully chlorinating the methyl group (CCl<sub>3</sub>) further depresses the conversion to 44%. These results indicate that the most reactive acyl moieties are those with relatively unencumbered electron-withdrawing substituents. Furthermore, appending even more strongly electronwithdrawing substituents such as trifluoro (CF<sub>3</sub>) substantially increases the conversion relative to R = CH<sub>3</sub>. In sum, these results indicate that electron-withdrawing groups which stabilize the RCOO<sup>-</sup> negative charge most enhance the ester hydrogenolysis turnover frequency."

Fig. 4 should appear as follows:




Fig. 4 Influence of acyl substituent on ester hydrogenolysis activity

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

a Department of Chemistry, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208, USA. E-mail: t-marks@northwestern.edu

b Materials Science Division, Argonne National Laboratory, Argonne, IL, 60439, USA

<sup>&</sup>lt;sup>c</sup> Center for Nanoscale Materials, Argonne National Laboratory, Argonne, IL, 60439, USA