Cite this: *Lab Chip*, 2017, 17, 567

DOI: 10.1039/c6lc90129h

www.rsc.org/loc

Correction: Double-emulsion drops with ultra-thin shells for capsule templates

Shin-Hyun Kim,^a Jin Woong Kim,^b Jun-Cheol Cho^c and David A. Weitz^{*a}

Correction for ‘Double-emulsion drops with ultra-thin shells for capsule templates’ by Shin-Hyun Kim et al., *Lab Chip*, 2011, 11, 3162–3166.

In the section “Diameter and shell thickness of double-emulsion drops” there are errors in eqn (2) and in the sentence that begins “In the same fashion, we calculate the thickness of the middle layer of double-emulsion drops which are produced at each values of Q_1/Q_2 and plot the results in Fig. 3c”. The equation should be

$$\frac{t}{R} = 1 - \left(1 + \frac{Q_2}{Q_1} \right)^{-1/3}.$$

The sentence should read “In the same fashion, we calculate the thickness of the middle layer of double-emulsion drops which are produced at each values of Q_2/Q_1 and plot the results in Fig. 3c”.

In the caption for Fig. 3c, “Relative thickness of shell to radius of the double-emulsion drops (t/R) as a function of Q_1/Q_2 .” should read “Relative thickness of shell to radius of the double-emulsion drops (t/R) as a function of Q_2/Q_1 .” In addition, the x-axis is incorrectly labelled with “ Q_1/Q_2 ”. The x-axis should be “ Q_2/Q_1 ”. A corrected version of Fig. 3c is shown.

Fig. 3 (c) Relative thickness of shell to radius of the double-emulsion drops (t/R) as a function of Q_2/Q_1 .

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a School of Engineering and Applied Sciences and Department of Physics, Harvard University, Cambridge, Massachusetts, USA. E-mail: weitz@seas.harvard.edu; Tel: +1 617 495 3275

^b Department of Applied Chemistry, Hanyang University, Ansan, South Korea

^c Amore-Pacific Co. R&D Center, Yongin, South Korea