MedChemComm

RESEARCH ARTICLE

Cite this: Med. Chem. Commun., 2017, 8, 1659

Received 11th May 2017, Accepted 20th June 2017

DOI: 10.1039/c7md00247e

rsc.li/medchemcomm

Bitopic fluorescent antagonists of the A_{2A} adenosine receptor based on pyrazolo[4,3e][1,2,4]triazolo[1,5-c]pyrimidin-5-amine functionalized congeners†

Romain Duroux,‡^{ab} Antonella Ciancetta, D‡^a Philip Mannes,^a Jinha Yu,^a Shireesha Boyapati,^{ac} [Eliz](http://orcid.org/0000-0002-8005-4599)abeth Gizewski,^d Said Yous,^b Francisco Ciru[ela,](http://orcid.org/0000-0001-8104-1493) ^{nee} John A. Auchampach, \mathbf{D}^d Zhan-Guo Gao^a and Kenneth A. Jacobson \mathbf{D}^{*a}

A pyrazolo[4,3-e][1,2,4]triazolo[1,5-c]pyrimidin-5-amine antagonist of the A_{2A} adenosine receptor (AR) was functionalized as amine congeners, fluorescent conjugates and a sulfonate, and the $A_{24}AR$ binding modes were predicted computationally. The optimal n-butyl spacer was incorporated into the following $A_{2A}AR\text{-se-}$ lective (K_i, nM) conjugates: BODIPY630/650 derivative 11 (MRS7396, 24.6) and AlexaFluor488 derivative 12 (MRS7416, 30.3). Flow cytometry of 12 in hA_{2A}AR-expressing HEK-293 cells displayed saturable binding (low nonspecific) and inhibition by known $A_{2A}AR$ antagonists. Water-soluble sulfonate 13 was a highly potent (K_i = 6.2 nM) and selective A_{2A}AR antagonist based on binding and functional assays. Docking and molecular dynamics simulations predicted the regions of interaction of the distal portions of these chainextended ligands with the A_{2A} AR. The BODIPY630/650 fluorophore of 11 was buried in a hydrophobic interhelical (TM1/TM7) region, while AlexaFluor488 of 12 associated with the hydrophilic extracellular loops. In conclusion, we have identified novel high affinity antagonist probes for $A_{24}AR$ drug discovery and characterization. RESEARCH ARTICLE
 (a) Check for underse **Bitopic fluorescent antagonists of the A_{2A}

case is a adenosine receptor based on pyrazolo[4,3-**

case and enosine **receptor based on pyrazolo[4,3-**

case and enosine **fluores**

^a Molecular Recognition Section, Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bldg. 8A, Rm. B1A-19, Bethesda, MD 20892-0810, USA.

E-mail: kennethj@niddk.nih.gov; Fax: +301 480 8422; Tel: +301 496 9024 b Inserm, CHU Lille, UMR-S 1172 - JPArc - Centre de Recherche Jean-Pierre</sup> AUBERT Neurosciences et Cancer, Univ. Lille, F-59000 Lille, France

 ϵ Department of Pharmaceutical Chemistry, Telangana University, Nizamabad, Telangana, India 503322

 d Department of Pharmacology, Medical College of Wisconsin, 8701 Watertown Plank Road, Milwaukee, Wisconsin 53226, USA

 e Unitat de Farmacologia, Departament Patologia i Terapeutica Experimental, Facultat de Medicina, IDIBELL, Universitat de Barcelona, 08907 L'Hospitalet de Llobregat, Spain

† Electronic supplementary information (ESI) available: Chemical synthesis, characterization data, HPLC analysis, pharmacological studies, off-target screening and additional molecular modeling procedures/results. 3D coordinates of the hA_{2A}AR in complex with 11 (docking pose). 3D coordinates of the hA_{2A}AR in complex with 12 (docking pose, BM2). Video of 30 ns of MD simulations of the hA_{2A}AR in complex with 3. Video S1. Trajectory visualization (left panel) and ligand–protein interaction energy profile (right panel) of 30 ns of membrane MD simulation of the 3-hA_{2A}AR complex. Video of 30 ns of MD simulations of hA_{2A} -AR in complex with 12. Video S2. Trajectory visualization of 30 ns of membrane MD simulation of the $12-hA_{2A}AR$ complex starting from the docking binding mode denoted "BM2": side view facing TM5, TM6, and TM7 (left panel), and side view facing TM4, TM5 and TM6 (right panel). See DOI: 10.1039/ c7md00247e

‡ These authors contributed equally.

Introduction

The development of selective agonists and antagonists of the four subtypes of adenosine receptors (ARs) has been extensively explored.^{1–3} Antagonists of the G_s protein-coupled A_{24} AR are sought for as agents for treating neurodegenerative conditions such as Parkinson's disease (PD) and Alzheimer's disease (AD), and for coadministration with cancer immunotherapy.4–⁷ Caffeine, the most consumed psychostimulant in the world, acts as a nonselective AR antagonist and readily enters the brain to antagonize the $A_{2A}AR$ at doses generally consumed. Epidemiological evidence showing a lower occurrence of AD and PD with modest caffeine intake points to the possibility that caffeine consumption is neuroprotective. Indeed, A2AAR antagonists can exert a neuroprotective effect on excitotoxicity in animal models; one mechanism appears to enhance the activity of an A_1AR , which forms a heterodimer with the $A_{2A}AR$, to inhibit glutamate release.⁸ $A_{2A}R$ antagonists also control microglia-mediated neuroinflammation.⁹ Therefore, it is possible that $A_{2A}AR$ antagonists may also delay neurodegenerative disease progression. In the striatum, $A_{2A}AR$ antagonists act to boost dopaminergic signaling, and thus provide symptomatic relief to reduce the motor deficits in PD without inducing dyskinesia.

Certain selective $A_{2A}AR$ antagonists, e.g. 1–4 (Chart 1), have been shown to enter the brain at sufficient levels for imaging and efficacy in PD models.¹⁰⁻¹² A caffeine-like 1,3,7-trialkylxanthine, istradefylline 1, is approved in Japan for treating PD, and its safety and efficacy in reducing in off-time (when PD symptoms return) in levodopa-treated patients were established in a 52 week trial.¹⁰ A tricyclic pyrazolo-triazolopyrimidine derivative 4, which binds potently and selectively to the $A_{2A}AR$, was in clinical trials for PD.¹²

In the periphery, elevated adenosine levels present in the microenvironment of tumors lead to a suppressed immune response, to shift the T cell response away from an aggressive state capable of attacking tumors.⁶ Increasing evidence supports the use of AR antagonists in cancer treatment, either with selectivity for the $A_{2A}AR$ alone or with dual selectivity for the A_{2A}AR and A_{2B}AR. Cytokine production in CD8⁺ chimeric antigen receptor (CAR) T cells was increased, and both CD8⁺ and CD4⁺ CAR T cells were activated upon blocking the $A_{2A}AR$ ⁷. Thus, there is great interest in the discovery of novel A2AAR antagonists to act either centrally or peripherally, and new tools for ligand discovery are needed.

Various fluorescent probes have been developed over the past few years for characterizing ARs ,¹³⁻¹⁵ and their use in fluorescence polarization (FP), fluorescence resonance energy

transfer (FRET) and flow cytometry (FCM) has been proven to be feasible. Moreover, these ligands can provide a better understanding of receptor location, function and regulation. Most of the reports on fluorescent $A_{2A}AR$ antagonists used 5-amino-7-(3-(4-methoxy)phenylpropyl)-2-(2-furyl)pyrazolo[4,3 e]-1,2,4-triazolo[1,5-c]pyrimidine (SCH442416, 3) as the pharmacophore due to the p-methoxyphenylpropyl side chain moiety of the antagonist, which served as the site for attachment of functionalized chains through an ether linkage.

Much of the ligand development for ARs now benefits from a detailed structural knowledge of the human (h) $A_{2A}AR$ ¹⁶⁻²⁰ More than two dozen high-resolution X-ray crystallographic $A_{2A}AR$ structures with bound agonists or antagonists have been determined and used for the in silico screening of chemical libraries. $A_{2A}AR$ complexes with antagonist 4- $(2-[7-amino-2-(2-fury1)[1,2,4]triazolo[2,3-a][1,3,5]triazin-5-yl$ amino]ethyl)phenol (ZM241385, 2) show that the heterocyclic pharmacophore binds to residues in the orthosteric binding pocket that also coordinates with the adenine moiety of AR agonists.^{16,17} Although no X-ray structure of the $3-A_{2A}AR$ complex has been determined, we used the $2-A_{2A}AR$ structure as the structural template for molecular modeling. The hypothetical docking pose of 3 and its AlexaFluor488 derivative 5

Chart 1 Structures of selective A_{2A}AR antagonists and the target series: (A) widely used pharmacological probes and clinical candidates (1-4); (B) reported fluorescent probe 5 derived from 3 and functionalized amine congeners of varied chain length 6 explored in this study.

established most of the conserved interactions in the orthosteric binding site. $13,14$ The modeling also predicted stabilizing interactions of the tethered fluorophore with specific charged and H-bonding residues of the second extracellular loop (EL2).

Conjugate 5 was not optimal for fluorescent binding due to its moderate hA_{2A}AR affinity ($K_i = 111$ nM).¹³ Also, its green light emission presents difficulties in fluorescence microscopy due to cell autofluorescence. Thus, there remains a need for $A_{2A}AR$ antagonist fluorescent probes of higher affinity and compatibility with microscopy. A BODIPY650/655 conjugate containing a secondary amine in the linking chain was reported to have a K_i value of 15 nM for the $A_{2A}AR$, but its utility was not established.¹³ With this aim, we explored further the SAR of the distal region of this chemical series by varying the chain length of the spacer group and the terminal fluorophore in order to enhance the affinity, selectivity and photophysical properties.

Results and discussion

We initially prepared a series of primary amine congeners of the A2AAR antagonist 3 by extending the p-methoxyphenylpropyl chain, which is predicted to lie outside the orthosteric binding site of the pharmacophore and be able to reach accessory sites.¹³ The congener 6a ($n = 2$) was reported earlier, 13 and we hypothesized that extension of the alkyl spacer could enhance the affinity or selectivity for this receptor, perhaps by establishing H-bond or electrostatic interactions with the polar residues in the EL region of the A_{2A}AR. The synthesis of the pyrazolo^{[4},3-e][1,2,4]triazolo[1,5c]pyrimidin-5-amine derivatives is shown in Scheme 1.

The synthesis of fluorescent ligands 9–12 was accomplished in 4 steps from commercially available 3. The latter was demethylated to 7 by the action of $BBr₃$ and then treated with an excess of methyl-2-bromoacetate (12 eq.) and Cs_2CO_3 as a base to provide ester 8 in 90% yield. Other methods, such as using NaH as the base with 1 eq. of methyl 2-bromoacetate in DMF, were attempted but led to the formation of di and tri-alkylated compounds. Amide synthesis was then performed by treatment of ester 8 with a series of alkyl diamines to give the amine congeners 6a–e. Two of these amines, 6a ($n = 2$) and 6c ($n = 4$), were reacted with activated fluorophore moieties in DMF in the presence of $Et₃N$ to

Scheme 1 Synthesis of A_{2A}AR antagonist functionalized congeners of 6 and their fluorescent derivatives 9-12. (a) BBr₃, CH₂Cl₂, rt, 4 h; (b) methyl 2-bromoacetate, Cs₂CO₃, MeOH, 40 °C, overnight; (c) diaminoalkane, MeOH (9:1 v/v), rt, overnight; (d) activated fluorophore (AlexaFluor647 N-hydroxysuccinimidyl ester for 9 and 10, BODIPY 630/650 N-hydroxysuccinimidyl ester for 11, AlexaFluor488 carboxylic acid, 2,3,5,6 tetrafluorophenyl ester for 12), Et₃N, DMF, rt, overnight.

afford, after semi-preparative HPLC, compounds 9–12 in good yield (40–60%): cyanine 5 red fluorescent AlexaFluor647 $(n = 2, 9; n = 4, 10)$, red fluorescent BODIPY630/650 $(n = 4, 10)$ 11) and green fluorescent AlexaFluor488 ($n = 4$, 12). Both fluorophores are commonly incorporated into ligand tools for chemical biology.²¹

The affinity of the amine congeners and other antagonists was measured using standard radioligand binding assays for hA₁, A_{2A} and A₃ARs (Table 1).^{1,13,14,22} Membranes of human embryonic kidney (HEK)293 cells expressing the AR of interest were used in the assay. Primary amine congeners 6a–6d containing spacers of 2–5 methylenes were similar in terms of hA_{2A}AR affinity (K_i 6–9 nM), with the butylamino congener 6c displaying the highest $A_{2A}AR$ selectivity in the series. The affinity of this compound for the hA_3AR was determined to be 6.0 μ M, with only slightly higher hA₁AR affinity. Homologation to 6 methylenes in compound 6e lowered $A_{2A}AR$ affinity. Among the fluorescent conjugates of butylamino congener 6c, *i.e.* 10, 11, and 12, the highest $A_{2A}AR$ affinity and selectivity were observed with the BODIPY630/650 fluorophore 11 and AlexaFluor488 12 (Fig. 1). However, fluorescent 11 and 12 exhibited only a 4-fold weaker $A_{2A}AR$ affinity with respect to the parent amino derivative 6c. Interestingly, Reasonsh Article

and (and the Composition Simulation 2012) in Access Articles. The mediation of the state and proportion and the state of the Creative Common Simulation 2017. And the state are state as point in common th

AlexaFluor488 conjugate 12 was at least as potent in terms of A_{2A} AR affinity as its shorter homologue 5. No dependence of the affinity on the chain length was evident when comparing AlexaFluor647 conjugates 9 and 10. Affinity for the mouse (m)ARs was also measured for the selected compounds by methods described, 22 and 11 was particularly potent and selective for the mA_{2A}AR ($K_i = 2.1$ nM), in contrast to 12 ($K_i =$ 585 nM).

We performed molecular modeling analysis to identify possible binding modes of fluorescent conjugates 11 and 12 using the high resolution 2-A2AAR X-ray structure (PDB ID: 4EIY).¹⁷ We first docked reference compound 3 at the $hA_{2A}AR$ by retaining several water molecules observed in the X-ray structure as described (ESI†). In the corresponding docking pose (Fig. 2A), the pyrazolotriazolopyrimidine core of 3 docked in the orthosteric binding site similar to the triazolopyrimidine core of 2, with a π - π stacking interaction of the aromatic core with F168 (EL2), and a H-bonding network with N253 (6.55, using standard GPCR notation²³) and E169 (EL2). The methoxy-phenyl substituent of 2 pointed toward the extracellular (EC) side of the receptor and interacted with the side chain of Glu169 (EL2) through a water molecule. The binding mode of 3 described above was validated

 ${\sf Table~1}$ AR binding affinity determined for a series of pyrazolo[4,3-e][1,2,4]triazolo[1,5-c]pyrimidin-5-amine derivatives (R¹, as in Scheme 1). Human ARs, unless otherwise noted (m indicates mouse)

 a Competition radioligand binding assays were conducted with membranes prepared from HEK-293 cells expressing recombinant A_1 , A_{2A} , or A₃ARs (human) unless otherwise noted. Their incubation was performed for 1 h at 25 °C. The radioligands used were: A₁AR, [³H]8-cyclopentyl-1,3-dipropylxanthine ([³H]DPCPX, 0.5 nM) 14; A_{2A}AR, [³H]ZM241385 2 (1.0 nM) or from published data¹³ (as noted) [³H]2-[p-(2-
carboxyethyl)phenyl-ethylamino]-5'-N-ethylcarboxamidoadenosine ([³H]CGS21680, 10 nM) methyluronamide ([¹²⁵Ι]Ι-AB-MECA, 0.2 nM) 16. Nonspecific binding was determined using 10 μM 8-[4-[[[[(2-aminoethyl)amino]carbonyl]methyl]oxy]phenyl]-1,3-dipropylxanthine (XAC) 17 (A1AR and A2AAR) or 10 μM adenosine-5′-N-ethyluronamide (NECA) 18 (A3AR). HEK-293 cells expressing recombinant mA₁, mA_{2A}, or mA₃ARs were used. Values are expressed as the mean \pm SEM from 3 independent experiments. The cell lines were from the American Type Culture Collection (ATCC, Manassas, VA), and the c were from the American Type Culture Collection (ATCC, Manassas, VA), and the cDNA for the ARs was obtained from [cdna.org.](http://cdna.org) ¹ Kumar et al. and Kecskés et al.^{13,14} ^c Percent inhibition at 10 µM. ^d Fluorophore moiety, as shown in Scheme 1 (R²). ^e Using [³H]16 as the radioligand.^{*j*} 11, MRS7396; 12, MRS7416; 13, MRS7352.

Fig. 1 Antagonist radioligand ($[{}^{3}H]2$, 1.0 nM) binding inhibition curves for the hA2AAR for four antagonist fluorescent conjugates. Compound numbers: MRS7322 9, MRS7395 10, MRS7396 11 and MRS7416 12. Membranes from HEK-293 cells expressing the hA_{2A}AR were used, and their incubation was performed for 1 h at 25 °C. Results are expressed as the mean \pm SEM. The K_i values from three independent experiments are listed in Table 1.

using 30 ns molecular dynamics (MD) simulation. During the simulation (Video S1, replica analysis reported in Table S1†), the methoxy-phenyl moiety folded toward transmembrane domain 2 (TM2) and established a $\pi-\pi$ stacking interaction with Y271 (7.36), while the aromatic core maintained the H-bond network observed in the initial docking pose. Notably, the most energetically favored ligand–protein complex featured the same interaction pattern described above (Fig. S1†).

The potent fluorescent conjugate 11 was then docked in the $A_{2A}AR$ structure (Fig. 2B) with the same water molecules retained as in the $3-A_{2A}AR$ complex.¹⁷ The pyrazolotriazolopyrimidine core established the same interactions observed for 3 in the orthosteric binding site. The fluorophore linker pointed toward the EC side and folded back toward the TM bundles by directing the fluorophore group to an aromatic pocket at the interface between TM1 and TM7 (transparent surface in Fig. 2B). To explore other orientations of the linker at the receptor's EC side, we docked compound 6c, the amino precursor of 11, at the $hA_{2A}AR$. To sample all the possible H-bond acceptor/donor partners on the EC side of the receptor, we removed water molecules interacting with E169 during the docking. We then clustered the binding modes obtained according to the linker orientation. This selection resulted in two alternative binding modes (Fig. S2†) that were subsequently subjected to MD validation (30 ns of simulation run in triplicate for each binding mode, see Table S1†). In the most energetically favored docking mode (hereby referred to as "BM1", orange carbon sticks in Fig. S2,† docking score = -12.077 kcal mol⁻¹), the tail was oriented toward TM4 and TM5 with the amide moiety establishing a H-bond with the side chain of E169, while the terminal amine group H-bonded with the backbone of E169 (EL2) and the side chain of K150 (EL2). This latter H-bond was expected to be unstable in a dynamic environment due to competition in the formation of a salt-bridge between K150 and D170 (EL2). In the alternative binding mode (BM2,

green carbon sticks in Fig. S2,† docking score = −10.994 kcal mol⁻¹), the tail was oriented toward TM1 and TM2 and did not establish additional interactions. MD trajectory analysis revealed that BM1 achieved ligand–protein complexes that were more energetically favored (data not shown). Fig. S3† depicts the two $11-A_{2A}AR$ complexes with the most favorable ligand–protein interaction energy (IE, values differed by less than 2 kcal mol⁻¹ and were considered equivalent) obtained for BM1 that features different orientations of the tail. We therefore searched for hydrophobic and aromatic regions in the proximity of the terminal amine moiety to investigate the compatibility of these orientations with the fluorophore insertion. Specifically, we searched for receptor regions rich in hydrophobic and aromatic residues at 5, 13, and 14 Å from the nitrogen atom of the terminal alkylamino group of 6c (Fig. S3B†). The choice of distances reflected the analysis of the N-group distance to aromatic rings in both the docked and the energy minimized three-dimensional structures of 11 (Fig. S3A†). Of the two possible orientations of 11, only one displayed hydrophobic and aromatic residues at distances compatible with the insertion of the aromatic fluorophore. Notably, the aromatic/hydrophobic region is located at the interface between TM1 and TM7 (Fig. S3C†), thus suggesting the same orientation of the ligand as observed from the docking analysis. Nonetheless, we cannot exclude that 11 might explore different regions on the receptor's EC side. Open Access Article. Published on 22 June 2017. Downloaded on 5/17/2024 8:26:21 PM. This article is licensed under a [Creative Commons Attribution 3.0 Unported Licence.](http://creativecommons.org/licenses/by/3.0/) **[View Article Online](https://doi.org/10.1039/c7md00247e)**

The fluorescent conjugate 12 (considered as the species carrying a −2 net charge) was docked in the $A_{2A}AR$ structure by following the same procedure described for compound 11 (ESI†). The docking output suggested two equally plausible orientations of the fluorophore (Fig. 2C and D). In one docking pose (BM1, cyan carbon sticks in Fig. 2C, docking score = -11.490 kcal mol⁻¹), the fluorophore group was projected toward EL3. In the alternative binding mode (BM2, purple carbon sticks, docking score = -11.279 kcal mol⁻¹), the fluorophore group interacted with the residues in EL2 and EL3. The interaction pattern of the core was the same as those for the other members of this chemical series: H-bonds with N253 and E169 (EL2); $π$ -π stacking with H252 (6.52), F168 (EL2), and Y271 (7.36). In the MD simulation starting from BM1 (30 ns run in triplicate, see Table S1†) the fluorophore group and the linker fluctuated on the EC surface of the receptor without engaging in specific interactions except for a H-bond involving the fluorophore carboxylate moiety (data not shown). On the other hand, the simulations starting from BM2 (30 ns, replica analysis in Table S1†) achieved more energetically favored ligand–protein complexes. The simulations returned the ligand–protein complex characterized by the lowest IE value for all trajectories and converged in a unique binding mode (Fig. S4†) featuring the fluorophore group stacked between EL2 and EL3. In such a conformation, the ligand established a tight network of H-bonds and salt-bridges between charged residues in EL2 and EL3 and the polar/charged counterparts in the fluorophore moiety. In particular, during the MD simulation (visualization of run 3 trajectory selected as an example, Video

Fig. 2 Molecular modeling of antagonist binding to the hA_{2A}AR. Details of the binding site of the X-ray structure of the receptor modeled with various ligands docked: (A) known antagonist 3, with the retention of a subset of water molecules found in the high resolution A_{2A} AR structure;¹⁷ (B) the BODIPY630/650-labeled antagonist 11, showing the most energetically favorable orientation of the terminal fluorophore chain; side (C) and top (D) views of the two possible orientations of the fluorophore group of derivative 12. Residues establishing polar (dashed orange lines) and π - π interactions with the docked ligands are represented as sticks. Aromatic residues establishing hydrophobic contacts with the terminal fluorophore of compound 11 are represented as transparent surfaces with colors matching the corresponding TM domain. Non-polar hydrogen atoms are omitted.

S2†) the sulfonic groups established salt bridges with K153 (persistent) and K150 (intermittent) in EL2, one of the amine moieties interacted with the sidechain of E169 (EL2), and the carboxylate moiety of the ligand replaced E169 (EL2) in the salt bridge with H264 (EL3). Moreover, for most of the total simulation time, the pyrazolotriazolopyrimidine core maintained its interaction pattern with N253 (6.55) and F168 (EL2), while the linker between the core and the fluorophore group was anchored to TM7, EL2 and TM2 through H-bond interactions with the backbone of S67 (2.65) and the sidechains of Q157 (EL2) and Y271 (7.36), respectively (Video S2†).

Thus, both 11 and 12 are bitopic in the sense that each bridged two separate domains of the $A_{2A}AR$, *i.e.* the orthosteric binding site that is well defined in X-ray structures $16,17$ and an additional domain. The BODIPY630/650 fluorophore of 11 is predicted to be buried in a hydrophobic region, while AlexaFluor488 of 12 associates with the hydrophilic ELs. The bitopic nature of these conjugates does not necessarily imply allosteric modulation of the orthosteric action, 28 which is unexplored in this series.

In order to further explore the ligand interactions with the EC surface, we prepared an aryl sulfonate 13, which contained a terminal group capable of multiple polar interactions (Scheme S1, ESI†). This terminal phenylsulfonic acid moiety also would allow for π - π interactions with aromatic residues. A further benefit of incorporating a sulfonate group is that it carries a permanent negative charge at physiological pH and would prevent penetration into the blood brain barrier,²² a useful characteristic for a pharmacological probe for

MedChemComm Research Article

in vivo studies.²⁴ 13 was potent and selective for binding to the h and $mA_{2A}ARs$, with 671- and 426-fold selectivity compared to the hA_1AR and hA_3AR , respectively. 13 binding to the mA_1AR and mA_3AR was insignificant.

The binding of 13 was modeled using the same docking procedure and MD validation described for 6c. Fig. 3 depicts the three alternative binding modes of 13. In the most energetically favorable pose (BM1, blue carbon sticks) the sulfophenyl ring established a π - π stacking interaction with H264 (EL3) and the sulfonic group engaged in an H-bond interaction with the residue backbone. In the other two docking poses (BM2 and BM3, magenta and orange carbon sticks, respectively) the sulfophenyl ring established a π -cation interaction with K153 (EL2) and the sulfonic group interacted with the side chain of either K153 (EL2; BM2, magenta) or S156 (EL2; BM3, orange). During the MD simulation, the ligand's sulfophenyl tail fluctuated considerably (high averaged root mean square deviation values, Table S1†), thus demonstrating the instability of the interaction pattern predicted in the initial docking poses. On the other hand, the three different initial poses converged in a unique binding mode, as the ligand–protein complexes with the lowest IE value returned by each different binding mode featured the same conformation (Fig. S5†). In particular, the 7-phenylpropyl ring moved toward TM7 to establish a $\pi-\pi$ stacking interaction with Y271 (7.36). NedChemCorem

In this area of the AARs, with CP-1 and 426celab edecking comparison in points/article. In this characteristic in the article is article in the article in the article in the specific in the specific interview

Two representative derivatives were shown to be $A_{2A}AR$ antagonists in a cyclic AMP assay (Fig. S6†). The activation curve for the known agonist 15 right-shifted in a parallel manner at fixed concentrations of 11 and 13. The EC_{50} of 15 shifted from 0.89 ± 0.17 nM to 128 ± 35 nM in the presence of 11 (1000 nM) and to 10.2 ± 2.3 nM in the presence of 13 (100 nM). Off-target binding activities at 45 diverse receptors were determined by $PDSP^{25}$ for selected compounds: 6c and 13 (ESI†). The only K_i values below 10 μM were: 6c, 3.36 μM, $5HT_{2A}$ serotonin receptor; 1.92 μ M, $5HT_{2B}$ receptor. Thus, the primary amine precursor of the fluorescent conjugates 10–12 was not promiscuous in its interaction with other proteins, $i.e.$ this chemical series is not associated with pan-assay interference compounds.²⁶

Compounds 11 and 12 were tested as fluorescent tracers for flow cytometry of HEK-293 cells expressing the $hA_{2A}AR$ (Fig. 4A). 11 proved to have high nonspecific binding, with a low level of specific binding amounting to <25% of the total and therefore was not optimal for flow cytometric analysis. Thus, 11 tended to bind non-specifically to hydrophobic membranes by undetermined mechanisms. However, the more hydrophilic 12 displayed low nonspecific binding by this method, and its binding was saturable with a K_d value of 45.4 nM (Fig. 4B). For inhibition studies, the cells were coincubated for one hour prior to flow cytometry with fluorescent probe 12 (10 nM) and a test antagonist. The binding of 12 was inhibited by two known $A_{2A}AR$ antagonists, nonxanthine 3 and xanthine 17, with the expected range of potency.²⁷ The K_i values were 3.8 and 17.2 nM (Fig. 5), respectively, which corresponded closely to the reported values of 4.1 and 18 nM at $hA_{2A}AR^{1,13}$ Agonist inhibition of the competitive binding of 12 was complex (ESI†).

Conclusions

We succeeded in identifying useful $A_{2A}AR$ probes by systematically varying the chain length of amine-functionalized congeners of a potent antagonist and coupling the primary amine having an optimal length to various fluorophores. We successfully enhanced their $A_{2A}AR$ affinity compared to known fluorescent A_{2A} AR ligands.¹⁵ Conjugates 11 and 12 were potent and selective antagonist probes for the $A_{2A}AR$, but 12 was more promising for characterization of the hA2AAR in whole cells by flow cytometry. Molecular modeling suggested that the fluorophore of 11 interacted with

Fig. 3 Modeling of antagonist 13 binding to the hA_{2A}AR: side (A) and top (B) views of the three possible orientations (BM1, blue; BM2, magenta; BM3, orange) of the p-sulfophenyl tail. Residues establishing polar (dashed orange lines) and $\pi-\pi$ interactions with the docked ligands are represented as sticks. Non-polar hydrogen atoms are omitted.

Fig. 4 Saturation of fluorescent antagonists 11 (A) and 12 (B) in hA2AAR-expressing HEK-293 cells measured using flow cytometry following a 1 h incubation at 37 °C. Non-specific binding was determined with 10 μ M 3. Results are expressed as the mean \pm SEM. The K_d value calculated from Fig. 4B is listed in the text.

hydrophobic regions between TMs. However, the hydrophilic fluorophore of 12 was coordinated to the ELs, consis-

Fig. 5 Inhibition of the specific binding of fluorescent antagonist 12 (10 nM, 1 h incubation at 37 °C) in hA_{2A} AR-expressing HEK-293 cells measured using flow cytometry. Competing antagonists were: nonxanthine 3 and xanthine 17. Non-specific binding was determined with 10 μ M 3. Results are expressed as the mean \pm SEM. The K_i values are listed in the text.

tent with its lower overall hydrophobicity and more favorable whole cell binding characteristics compared to 11 under the present conditions. Evaluation of 11 and 12 by confocal microscopy studies remains to be performed. Thus, we have introduced antagonist ligands displaying high A_{2A}AR affinity and selectivity that may serve as versatile tools to better study this receptor.

Conflict of interest

The authors declare no competing interests.

Abbreviations

Acknowledgements

This research was supported in part by the Intramural Research Program of the NIH, National Institute of Diabetes and Digestive and Kidney Diseases. RD was supported by Inserm, Univ. Lille, France. We thank Dr. John Lloyd (NIDDK) for mass spectroscopy. SB was supported by UGC, India. JA acknowledges support from the National Heart, Lung, and Blood Institute (R01HL133589).

Notes and references

- 1 C. E. Müller and K. A. Jacobson, Biochim. Biophys. Acta, Biomembr., 2011, 1808, 1290–1308.
- 2 G. Yuan, N. G. Gedeon, T. C. Jankins and G. B. Jones, Expert Opin. Drug Discovery, 2015, 10, 63–80.
- 3 P. G. Baraldi, M. A. Tabrizi, S. Gessi and P. A. Borea, Chem. Rev., 2008, 108, 238–263.
- 4 C. V. Gomes, M. P. Kaster, A. R. Tomé, P. M. Agostinho and R. A. Cunha, Biochim. Biophys. Acta, Biomembr., 2011, 1808, 1380–1399.
- 5 C. Laurent, S. Burnouf, B. Ferry, V. L. Batalha, J. E. Coelho, Y. Baqi, E. Malik, E. Mariciniak, S. Parrot, A. Van der Jeugd, E. Faivre, V. Flaten, C. Ledent, R. D'Hooge, N. Sergeant, M. Hamdane, S. Humez, C. E. Müller, L. V. Lopes, L. Buée and D. Blum, Mol. Psychiatry, 2016, 21, 97–107.
- 6 S. M. Hatfield and M. Sitkovsky, Curr. Opin. Pharmacol., 2016, 29, 90–96.
- 7 P. A. Beavis, M. A. Henderson, L. Giuffrida, J. K. Mills, K. Sek, R. S. Cross, A. J. Davenport, L. B. John, S. Mardiana, C. Y. Slaney, R. W. Johnstone, J. A. Trapani, J. Stagg, S. Loi, L. Kats, D. Gyorki, M. H. Kershaw and P. K. Darcy, J. Clin. Invest., 2017, 127, 929–941.
- 8 M. T. Armentero, A. Pinna, S. Ferré, J. L. Lanciego, C. E. Müller and R. Franco, Pharmacol. Ther., 2011, 132, 280–299.
- 9 M. H. Madeira, R. Boia, A. F. Ambrósio and A. R. Santiago, Mediators of Inflammation, 2017, 2017, 4761081, DOI: 10.1155/2017/4761081.
- 10 T. Kondo and Y. Mizuno, Clin. Neuropharmacol., 2015, 38, 41–46.
- 11 A. F. Ramlackhansingh, S. K. Bose, I. Ahmed, F. E. Turkheimer, N. Pavese and D. J. Brooks, Neurology, 2011, 76, 1811–1816.
- 12 X. Zhou, S. Khanapur, A. P. Huizing, R. Zijlma, M. Schepers, R. A. J. O. Dierckx, A. van Waarde, E. F. J. de Vries and P. H. Elsinga, J. Med. Chem., 2014, 57, 9204–9210.
- 13 T. S. Kumar, S. Mishra, F. Deflorian, L. S. Yoo, K. Phan, M. Kecskés, A. Szabo, B. A. Shinkre, Z. G. Gao, W. C. Trenkle and K. A. Jacobson, Bioorg. Med. Chem. Lett., 2011, 21, 2740–2745.
- 14 M. Kecskés, T. S. Kumar, L. Yoo, Z. G. Gao and K. A. Jacobson, Biochem. Pharmacol., 2010, 80, 506–511.
- 15 F. Ciruela, V. Fernández-Dueñas and K. A. Jacobson, Neuropharmacology, 2015, 98, 58–67.
- 16 V. P. Jaakola, M. T. Griffith, M. A. Hanson, V. Cherezov, E. Y. T. Chien, J. R. Lane, A. P. IJzerman and R. C. Stevens, Science, 2008, 322, 1211.
- 17 W. Liu, E. Chun, A. A. Thompson, P. Chubukov, F. Xu, V. Katritch, G. W. Han, C. B. Roth, L. H. Heitman, A. P. IJzerman, V. Cherezov and R. C. Stevens, Science, 2012, 337, 232–236.
- 18 C. J. Langmead, S. P. Andrews, M. Congreve, J. C. Errey, E. Hurrell, F. H. Marshall, J. S. Mason, C. M. Richardson, N. Robertson, A. Zhukov and M. Weir, J. Med. Chem., 2012, 55, 1904–1909.
- 19 V. Katritch, V. P. Jaakola, J. R. Lane, J. Lin, A. P. IJzerman, M. Yeager, I. Kufareva, R. C. Stevens and R. Abagyan, J. Med. Chem., 2010, 53, 1799.
- 20 D. Rodríguez, Z. G. Gao, S. M. Moss, K. A. Jacobson and J. Carlsson, J. Chem. Inf. Model., 2015, 55, 550–563.
- 21 L. D. Lavis and R. T. Raines, ACS Chem. Biol., 2008, 3, 142–155.
- 22 S. Paoletta, D. K. Tosh, A. Finley, E. Gizewski, S. M. Moss, Z. G. Gao, J. A. Auchampach, D. Salvemini and K. A. Jacobson, J. Med. Chem., 2013, 56, 5949–5963.
- 23 J. A. Ballesteros and H. Weinstein, Methods Neurosci., 1995, 25, 366.
- 24 J. L. Carlin, D. K. Tosh, C. Xiao, R. A. Piñol, Z. Chen, D. Salvemini, O. Gavrilova, K. A. Jacobson and M. L. Reitman, J. Pharmacol. Exp. Ther., 2016, 356, 474–482.
- 25 J. Besnard, G. F. Ruda, V. Setola, K. Abecassis, R. M. Rodriguiz, X. P. Huang, S. Norval, M. F. Sassano, A. I. Shin, L. A. Webster, F. R. Simeons, L. Stojanovski, A. Prat, N. G. Seidah, D. B. Constam, G. R. Bickerton, K. D. Read, W. C. Wetsel, I. H. Gilbert, B. L. Roth and A. L. Hopkins, Nature, 2012, 492, 215–220. Open Common Common Access Articles. Published on 22 June 2017. A Licensed on 22 June 2017. Downloaded on 22 June 2017. Downloaded on 5.26:21 PM. The Licensed under Access Article is licensed under Access Article is license
	- 26 C. Aldrich, C. Bertozzi, G. I. Georg, L. Kiessling, C. Lindsley, D. Liotta, K. M. Merz Jr., A. Schepartz and S. Wang, ACS Chem. Biol., 2017, 12, 575–578.
	- 27 C. E. Müller and K. A. Jacobson, Xanthines as adenosine receptor antagonists, in Methylxanthines, Handb. Exp. Pharmacol., ed. B. B. Fredholm, Springer, 2011, vol. 200, pp. 151–199.
	- 28 G. Vauquelin, D. Hall and S. J. Charlton, Br. J. Pharmacol., 2015, 172, 2300–2315.