



Cite this: *New J. Chem.*, 2017,
41, 8673

DOI: 10.1039/c7nj90060k

rsc.li/njc

Correction: Understanding the reactivity of bis(propargyl) aromatic esters towards GAP: a theoretical exploration

Sanjeevani H. Sonawane,^a Shaibal Banerjee,^b Arun K. Sider,^a Mahadev B. Talawar^{*a}
and Md Abdul Shafeeuulla Khan^{*a}

Correction for 'Understanding the reactivity of bis(propargyl) aromatic esters towards GAP: a theoretical exploration' by Sanjeevani H. Sonawane *et al.*, *New J. Chem.*, 2017, DOI: 10.1039/c7nj01172e.

The authors would like to correct an error in the sentence beginning on Page 4, left column, line 52. This sentence should read:

In the reactant adduct **6R**, the isocyanate group of **6** and hydroxy moiety of **GAP-2** come close at distances of 2.460 Å (N···H) and 2.981 Å (C···O) with an O–H bond distance of 0.964 Å, whereas in the product **6P**, there was complete urethane linkage formation with the corresponding parameters N–H, C–O, and O···H of 1.005 Å, 1.361 Å, and 2.311 Å, respectively.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Energetic Materials Research Division, High Energy Materials Research Laboratory (Defence Research & Development Organization), Pune, 411 021, India.
E-mail: talawar.mb@hemrl.drdo.in, maskhan@hemrl.drdo.in

^b Defence Institute of Advanced Technology (Defence Research & Development Organization), Girinagar, Pune, 411 025, India

