Cite this: *Nanoscale*, 2017, **9**, 16485

Correction: Bismuth oxyiodide coupled with bismuth nanodots for enhanced photocatalytic bisphenol A degradation: synergistic effects and mechanistic insight

Shunqin Luo,^a Jinjia Xu,^b Zijing Li,^a Chen Liu,^c Jiawei Chen,^{*a} Xin Min,^d Minghao Fang^d and Zhaohui Huang^{*d}

DOI: 10.1039/c7nr90222k

rsc.li/nanoscale

Correction for 'Bismuth oxyiodide coupled with bismuth nanodots for enhanced photocatalytic bisphenol A degradation: synergistic effects and mechanistic insight' by Shunqin Luo *et al.*, *Nanoscale*, 2017, DOI: 10.1039/c7nr05320g.

The authors have noticed an error in the published article. In the sentence beginning "The κ_{ET} from pristine..." the calculated value of " $2.3 \times 10^8 \text{ s}^{-1}$ " should instead read as " $0.23 \times 10^9 \text{ s}^{-1}$ "

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aState Key Laboratory of Biogeology and Environmental Geology, School of Earth Sciences and Resources, China University of Geosciences (Beijing), 29 Xueyuan Road, Beijing 100083, China. E-mail: chenjiawei@cugb.edu.cn

^bMolecular Design & Function Group, National Institute for Materials Science (NIMS), 1-2-1 Sengen, Tsukuba, Ibaraki 305-0047, Japan

^cEnvironmental Protection Center, China Waterborne Transport Research Institute, 8 Xitucheng Road, Beijing 100088, China

^dSchool of Materials Science and Technology, China University of Geosciences (Beijing), 29 Xueyuan Road, Beijing 100083, China. E-mail: huang118@cugb.edu.cn