



Cite this: *Org. Biomol. Chem.*, 2017, **15**, 7339

Selective lysine modification of native peptides via aza-Michael addition†

Hongli Chen, * Rong Huang, Zhihong Li, Wei Zhu, Jiakang Chen, Yuexiong Zhan and Biao Jiang*

A series of vinylsulfonamides were synthesized and screened for site-selective modification of the ϵ -amino group of lysine-bearing free α -amine residues. *N*-Methyl-*N*-phenylethanesulfonamide has emerged as an applicable reagent and has been developed for efficient and highly selective modification of the lysine residue of native peptides in the presence of a free N-terminus via aza-Michael addition. We demonstrated that functional *N*-phenylvinylsulfonamide derivatives with a fluorescent moiety or drug could also be conjugated to the lysine residue of octreotide and insulin with high specificity, without modifying the N-terminus. Our method provides a promising strategy for site-selective lysine functionalization in native peptides with a free N-terminus.

Received 27th July 2017,
Accepted 21st August 2017

DOI: 10.1039/c7ob01866e

rsc.li/obc

Introduction

The modification of native peptides at a single amino acid or specific site with synthetic moieties is a significant challenge.^{1–3} Among the approximately 20 primary amino acids that compose proteins, only a subset can serve as appropriate targets for bioconjugation. Many strategies have been developed to target nucleophilic natural amino acid residues, of which cysteine and lysine residues are the most popular labeling sites. Cysteine is capable of being involved in a chemoselective reaction due to its superior nucleophilic properties.^{4–7} However, the low abundance of free cysteine residues due to its tendency to form disulfide bonds limits its application. Lysine residues are abundant in native proteins, and methods to modify primary amines can provide versatile techniques to label lysine residues.^{8,9} However, lysine conjugation methods often modify the N-terminus.^{10,11} Site-specific modification of the N-terminus is easier to achieve because the ϵ -amino group of lysine is more difficult to deprotonate due to its higher pK_a (~ 10) compared to the N-terminal α -amino group ($pK_a \sim 8$). A number of strategies for selective N-terminal modification of native peptides or proteins have been developed.^{12–18} In contrast, site-specific functionalization of the ϵ -amino group of lysine has rarely been studied.^{19,20} Typically, the most common reagents for amine modification, *N*-hydroxysuccinimides (NHS) esters, also react with serine,

tyrosine and histidine, affording heterogeneous bioconjugates.²¹ Although the stepwise addition of the substoichiometric amounts of an NHS ester can achieve a single lysine-selective modification, tediousness and low efficiency are the problems.²² Indeed, low reactivity and the generation of byproducts are the major problems encountered when attempting to selectively label an amine. The Michael-type addition is also an attractive strategy for protein conjugation. Michael acceptors, such as maleimides, vinyl sulfones and acrylamides, have been frequently utilized in the modification of proteins. Vinyl sulfonamides, with a similar structure to vinyl sulfones, are easier to prepare and have also received some attention in applications for bioconjugation.^{23–26}

Although the lysine ϵ -amino group requires a higher pH to deprotonate, it has higher intrinsic nucleophilicity. Therefore, selective lysine modification can be achieved by carefully adjusting the nucleophilicity and basicity. Herein, we report a class of vinylsulfonamides which site-specifically modify the lysine residue in native peptides rather than the N-terminus via aza-Michael addition.

Results and discussion

We first synthesized a series of vinylsulfonamides (Fig. 1, compounds 1–7). We found that activated phenylamines were the easiest to use to produce divinylsulfonamides (Fig. 1, compounds 8–10), which would be developed as efficient reagents for peptide stapling in our coming research. Then, we began to screen the abilities of different vinylsulfonamides to promote selective ϵ -amino group modification. Our initial study found that triethylenediamine (DABCO) buffer was a good medium

The institute for Advanced Immunochemical Studies, ShanghaiTech University, 99 Haike Road, Pudong, Shanghai, 201210, P.R. China.

E-mail: chenhl@shanghaitech.edu.cn, jiangbiao@shanghaitech.edu.cn

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c7ob01866e







Fig. 1 The structure of vinylsulfonamides 1–10.

for the aza-Michael addition to take place. Therefore we employed an *L*-lysine methyl ester (**11**) to react with vinylsulfonamides (**1–7**) in DABCO buffer and CH₃CN (1 : 1) at pH = 7. Based on RP-HPLC analysis, compounds **1–2** did not react with Lys-OMe (ESI, Table S1,† entries 1 and 2). Compounds **3–4** showed poor selectivity (ESI, Table S1,† entries 3 and 4). Compounds **5–6** displayed good selectivity, however the conversion was poor (<20%) (ESI, Table S1,† entries 5 and 6). *N*-Methyl-*N*-phenylethanesulfonamide (compound **7**) emerged as a potential reagent with good selectivity and moderate conversion (~30%) (ESI, Table S1,† entry 7). We further screened a series of alkaline buffers at pH 7–9 for the reaction of compound **7** and Lys-OMe (**11**) (ESI, Table S1,† entries 8–17). Unfortunately, the selectivity or conversion became bad under all of those reaction conditions.

Compound **7** was allowed to react with Lys-OMe (**11**, 3 eq.) in the presence of trimethylamine (Et₃N 6 eq.) in CH₃CN. Compound **7** was consumed, and only one high yield product (**12**, purified yield: 87%) was obtained, which was mono-labeled based on the HRESIMS of the [M + H]⁺ and [M + Na]⁺ (ion peaks at 358.1917 and 380.1615, respectively). The attached position was the ε-amino group rather than the α-amino group based on HSQC and HMBC analysis; key correlations were observed between H-8 and C-10, C-13, C-14 (Scheme 1).

Having observed the high yield and good selectivity of the *N*-methyl-*N*-phenylethanesulfonamide (compound **7**)-based conjugate, we next used this method to label peptides. Octreotide (**13**), a synthetic octapeptide with the sequence H₂N-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-ol (disulfide bridge Cys2–Cys7) that is one of the most extensively studied somatostatin analogues, was selected to evaluate the lysine site-specific modification. Octreotides have been widely investigated for conjugation with toxic drugs for targeted delivery into SSTR-selective cancer cells. The studies have revealed that a mixture of products (modification at the N-terminus, the lysine residue, or both) was produced when octreotide was allowed to react with NHS esters.^{27–30} Octreotide was treated with compound **7** (3 eq.) in the presence of Et₃N (6 eq.) in a



Scheme 1 Selective modification of the ε-amino group.

CH₃CN and H₂O (1 : 1) solution at room temperature. The reaction proceeded smoothly, and octreotide was almost completely converted to a single product (**14**) after 4 h (Fig. 2A), as





Fig. 2 Stepwise modification of the lysine residue and N-terminus of octreotide. (A) The reaction scheme. (B) RP-HPLC of the reaction. (C) The HRESIMS spectrum of product **14**. (D) The MS² spectrum of product **14**. (E) Peptide fragments of compound **14**. (F) The HRESIMS spectrum of product **15**. (G) The MS² spectrum of product **15**. (H) Peptide fragments of compound **15**.



sion of insulin was poor (<10%). The effect of pH was examined based on the DBU buffer. By screening buffers from pH 7 to pH 12, we found that the conversion of insulin was low

when the pH was below 8, and the reaction became complicated when the pH was more than 10. Moderate conversion (52%) and excellent selectivity (almost a single product **19** in 49% yield and 90% yield based on the recovered insulin) were observed in pH 9 buffer adjusted with DBU (Fig. 4A). The iso-



Scheme 2 Synthesis of functional vinylsulfonamides **20** and **21**.




Fig. 5 Selective lysine modification of octreotide with functional vinylsulfonamides **20** and **21**. (A) The reaction scheme. (B) The HRESIMS and MS² spectrum of product **30**. (C) The MS² spectrum of product **30**. (D) The HRESIMS and MS² spectrum of product **31**.




Fig. 6 Selective lysine modification of insulin with functional vinylsulfonamides **20** and **21**. (A) The reaction scheme. (B) The HRESIMS and MS² spectrum of product **32**. (C) The HRESIMS and MS² spectrum of product **33**. (D) RP-HPLC of the reaction showing that compound **21** was conjugated to insulin.



lated product **19** exhibited signals with multiple charges at 1001.6223 $[M + 6H]^{6+}$, 1201.7446 $[M + 5H]^{5+}$ and 1502.1787 $[M + 4H]^{4+}$, demonstrating that it was a mono-labeled product (Fig. 4B). After the reduction of the disulphide bonds and in-solution digestion of compound **19** by chymotrypsin, peptide fragments were obtained and analyzed by MS² (Fig. 4C). The modification site was determined as B29 (lysine) based on the fragment (GFFYTPKT+ compound **7**) with m/z 579.2698 $[M + 2H]^{2+}$, which corresponded well to the calculated M.W. of 1156.5263.

An effective bioconjugation method has to have the ability to attach functional groups of interest, such as fluorescent probes and drugs. Therefore, we further explored vinylsulfonamides as versatile building blocks for the site-specific prepa-

ration of more interesting bioconjugates. Coumarin- and combretastatin A-4 (CA4, a potent cytotoxic agent)-functionalized vinylsulfonamides (**20** and **21**) were synthesized. The synthesis of product **20** was carried out by condensation of *N*-(2-aminoethyl)benzeneamine (**22**) and coumarin-3-carboxylic acid (**23**) to form the amide (**24**), followed by treatment with 2-chloroethanesulfonyl chloride (**25**) to afford the desired product **20**. Compound **21** was prepared in a similar manner: Conjugation of CA4 derivative **28**, easily obtained from CA4 (**26**) according to the reported procedure,³⁶ with compound **22** yielded compound **29**, which was subsequently allowed to react with compound **25** to obtain the target product **21** (Scheme 2).

Based on the successful incorporation of the probes into vinylsulfonamides, their selective conjugation ability for the




Fig. 7 Modification of somatostatin **34**. (A) The reaction scheme. (B) The HRESIMS spectrum of product **35**. (C) The HRESIMS spectrum of the reduced product of compound **35**. (D) The MS² spectrum of the reduced product of compound **35**. (E) Peptide fragments. (F) The HRESIMS spectrum of product **36**. (G) The HRESIMS spectrum of the reduced product of compound **36**. (H) The MS² spectrum of the reduced product of compound **36**. (I) Peptide fragments.



peptides, octreotide and insulin was investigated. As previously described, octreotide was allowed to react with compounds **20** and **21** in the presence of Et_3N , leading to the mono-labeled derivatives **30** and **31** with high yield and successful selectivity (Fig. 5A). For compound **30**, the mass peak of m/z 709.2782 $[\text{M} + 2\text{H}]^{2+}$ and 1417.5403 $[\text{M} + \text{H}]^+$ was found in the HRESIMS spectrum, matching its calculated M.W. of 1416.5341 g mol^{-1} (Fig. 5B). After in-gel digestion by chymotrypsin, the MS^2 spectrum of compound **30** was obtained. The degradation product (m/z 1109.4239) was detected under MS^2 conditions due to the instability of the attached probe **20**. Compound **20** was degraded to a residue with m/z 308.1147 when it was directly used for the MS^2 experiment under the same conditions. The modified peptide fragment Lys-Thr-Cys-Thr-ol with m/z 527.1955 included a fragment from probe **20** (m/z 399.04 – 308.11 = 90.93) and the unmodified peptide fragment $\text{H}_2\text{N-D-Phe-Cys-Phe-D-Trp}$ with m/z 584.2334, indicating labeling at lysine (Fig. 5C). For compound **31**, the mono-lysine-modified feature was also determined by HRESIMS with signals at m/z 801.3308 $[\text{M} + 2\text{H}]^{2+}$ and 1602.6505 $[\text{M} + \text{H}]^+$ (calculated value: 1600.6441) and MS^2 with an unmodified peptide fragment $\text{H}_2\text{N-D-Phe-Cys-Phe-D-Trp}$ (m/z 584.2324) (Fig. 5D).

In the case of insulin, to conjugate with compound **20** or **21**, specific mono-modification of the B29 lysine was performed using the above optimized method with DBU buffer at $\text{pH} = 9$ (Fig. 6A). Products **32** and **33** were also confirmed by MS and MS^2 (Fig. 6B and C). Compound **21** was an excellent coupling reagent for insulin that resulted in successful conversion (>95%), yield (92%) and selectivity (almost a single product) (Fig. 6D).

Somatostatin (**34**) which has a free N-terminus and two lysine residues with the sequence H-Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Thr-Ser-Cys-OH (disulfide bridge Cys3–Cys14) was employed to investigate whether there was a preference for a single lysine residue. Somatostatin was reacted with compound **7** in DBU buffer ($\text{pH} = 9$) (Fig. 7A). The resulting HRESIMS spectrum revealed that a mono-labeled product **35** (9%) was obtained, with signals at 612.2921 $[\text{M} + 3\text{H}]^{3+}$ and 917.9351 $[\text{M} + 2\text{H}]^{2+}$ (Fig. 7B). Similarly, after the reduction of product **35** with TCEP (Fig. 3C), peptide fragments were obtained and identified by MS^2 (Fig. 7D and E). The single K9 residue of somatostatin was modified. However, the conversion was low and the reaction system became complicated with the prolonged reaction time. When somatostatin (**34**) was reacted with compound **7** (3 eq.) in the presence of Et_3N (6 eq.) in a CH_3CN and H_2O (1 : 1), a main di-labelled product (**36**) was yielded (63%) with signals at 678.3126 $[\text{M} + 3\text{H}]^{3+}$ and 1016.9655 $[\text{M} + 2\text{H}]^{2+}$ (Fig. 7F). After the reduction of product **36** with TCEP (Fig. 7G), the resulting HRESIMS spectrum showed that A1 and K9 were modified (Fig. 7H–I).

Conclusions

Selective and effective modification of a lysine residue in the presence of an N-terminus is anticipated to be difficult due to

the lower pK_a of the N-terminal position. Here, a useful and efficient method for utilizing vinylsulfonamide-based aza-Michael addition to selectively modify lysine instead of the N-terminus on peptides was developed. Furthermore, we demonstrated that functional vinylsulfonamide derivatives could also be produced *via* reaction with a fluorescent moiety or drug and achieve the selective bioconjugation. The CA4-conjugated probe showed excellent reactivity and selectivity to modify the B29 lysine of insulin. This is a promising method for native peptide modification where the N-terminus is crucial for peptide bioactivity. However, peptides that contain multiple lysine residues will result in a complicated situation. And our method is more beneficial for peptides that lack an accessible free Cys residue which has superior reactivity. Efforts to develop more useful vinylsulfonamide reagents and application of this method to modify peptides or proteins are currently underway.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by ShanghaiTech university.

Notes and references

- 1 N. Krall, F. P. da Cruz, O. Boutureira and G. J. Bernardes, *Nat. Chem.*, 2016, **8**, 103–113.
- 2 O. Koniev and A. Wagner, *Chem. Soc. Rev.*, 2015, **44**, 5495–5551.
- 3 O. Boutureira and G. J. Bernardes, *Chem. Rev.*, 2015, **115**, 2174–2195.
- 4 A. Abbas, B. Xing and T. P. Loh, *Angew. Chem., Int. Ed.*, 2014, **53**, 7491–7494.
- 5 I. M. Serafimova, M. A. Pufall, S. Krishnan, K. Duda, M. S. Cohen, R. L. Maglathlin, J. M. McFarland, R. M. Miller, M. Frodin and J. Taunton, *Nat. Chem. Biol.*, 2012, **8**, 471–476.
- 6 D. Kalia, P. V. Malekar and M. Parthasarathy, *Angew. Chem., Int. Ed.*, 2016, **55**, 1432–1435.
- 7 S. B. Gunnoo and A. Madder, *ChemBioChem*, 2016, **17**, 529–553.
- 8 H. G. Lee, G. Lautrette, B. L. Pentelute and S. L. Buchwald, *Angew. Chem., Int. Ed.*, 2017, **56**, 3177–3181.
- 9 C. L. Tung, C. T. Wong, E. Y. Fung and X. Li, *Org. Lett.*, 2016, **18**, 2600–2603.
- 10 S. T. Larda, D. Pichugin and R. S. Prosser, *Bioconjugate Chem.*, 2015, **26**, 2376–2383.
- 11 S. Diethelm, M. A. Schafroth and E. M. Carreira, *Org. Lett.*, 2014, **16**, 3908–3911.
- 12 D. Chen, M. M. Disotuar, X. Xiong, Y. Wang and D. H.-C. Chou, *Chem. Sci.*, 2017, **8**, 2717–2722.



- 13 J. I. MacDonald, H. K. Munch, T. Moore and M. B. Francis, *Nat. Chem. Biol.*, 2015, **11**, 326–331.
- 14 Z. W. Lai, A. Petrera and O. Schilling, *Curr. Opin. Chem. Biol.*, 2015, **24**, 71–79.
- 15 A. C. Obermeyer, J. B. Jarman and M. B. Francis, *J. Am. Chem. Soc.*, 2014, **136**, 9572–9579.
- 16 A. O. Chan, C. M. Ho, H. C. Chong, Y. C. Leung, J. S. Huang, M. K. Wong and C. M. Che, *J. Am. Chem. Soc.*, 2012, **134**, 2589–2598.
- 17 G. Xu, S. B. Shin and S. R. Jaffrey, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 19310–19315.
- 18 J. M. Gilmore, R. A. Scheck, A. P. Esser-Kahn, N. S. Joshi and M. B. Francis, *Angew. Chem., Int. Ed.*, 2006, **45**, 5307–5311.
- 19 M. Chilamari, L. Purushottam and V. Rai, *Chem. – Eur. J.*, 2017, **23**, 3819–3823.
- 20 L. Purushottam, S. R. Adusumalli, M. Chilamari and V. Rai, *Chem. Commun.*, 2017, **53**, 959–962.
- 21 N. Abello, H. A. Kerstjens, D. S. Postma and R. Bischoff, *J. Proteome Res.*, 2007, **6**, 4770–4776.
- 22 X. Chen, K. Muthoosamy, A. Pfisterer, B. Neumann and T. Weil, *Bioconjugate Chem.*, 2012, **23**, 500–508.
- 23 J. Dadova, M. Vrabel, M. Adamik, M. Brazdova, R. Pohl, M. Fojta and M. Hocek, *Chem. – Eur. J.*, 2015, **21**, 16091–16102.
- 24 J. L. Furman, M. Kang, S. Choi, Y. Cao, E. D. Wold, S. B. Sun, V. V. Smider, P. G. Schultz and C. H. Kim, *J. Am. Chem. Soc.*, 2014, **136**, 8411–8417.
- 25 E. H. Suh, Y. Liu, S. Connelly, J. C. Genereux, I. A. Wilson and J. W. Kelly, *J. Am. Chem. Soc.*, 2013, **135**, 17869–17880.
- 26 J. Dadova, P. Orsag, R. Pohl, M. Brazdova, M. Fojta and M. Hocek, *Angew. Chem., Int. Ed.*, 2013, **52**, 10515–10518.
- 27 H. Y. Zhang, W. Q. Xu, Y. W. Wang, E. Omari-Siaw, Y. Wang, Y. Y. Zheng, X. Cao, S. S. Tong, J. N. Yu and X. M. Xu, *Int. J. Pharm.*, 2016, **502**, 98–106.
- 28 M. Huo, Q. Zhu, Q. Wu, T. Yin, L. Wang, L. Yin and J. Zhou, *J. Pharm. Sci.*, 2015, **104**, 2018–2028.
- 29 D. H. Na, K. C. Lee and P. P. DeLuca, *Pharm. Res.*, 2005, **22**, 743–749.
- 30 T. Yin, Q. Wu, L. Wang, L. Yin, J. Zhou and M. Huo, *Mol. Pharm.*, 2015, **12**, 3020–3031.
- 31 A. N. Zaykov, J. P. Mayer and R. D. DiMarchi, *Nat. Rev. Drug Discovery*, 2016, **15**, 425–439.
- 32 D. Jacob, M. Joan Taylor, P. Tomlins and T. S. Sahota, *J. Fluoresc.*, 2016, **26**, 617–629.
- 33 D. H. Chou, M. J. Webber, B. C. Tang, A. B. Lin, L. S. Thapa, D. Deng, J. V. Truong, A. B. Cortinas, R. Langer and D. G. Anderson, *Proc. Natl. Acad. Sci. U. S. A.*, 2015, **112**, 2401–2406.
- 34 T. H. Jensen, S. Havelund, P. K. Nielsen and J. Markussen, *J. Am. Chem. Soc.*, 2005, **127**, 6158–6159.
- 35 K. Thibaudeau, R. Leger, X. Huang, M. Robitaille, O. Quraishi, C. L. Soucy, N. B. Gagnon, P. V. Wyk, V. Paradis, J. P. Castaigne and D. Bridon, *Bioconjugate Chem.*, 2005, **16**, 1000–1008.
- 36 A. Kamal, A. Mallareddy, M. Janaki Ramaiah, S. N. Pushpavalli, P. Suresh, C. Kishor, J. N. Murty, N. S. Rao, S. Ghosh, A. Addlagatta and M. Pal-Bhadra, *Eur. J. Med. Chem.*, 2012, **56**, 166–178.

