Organic & Biomolecular Chemistry

COMMUNICATION

Cite this: Org. Biomol. Chem., 2017, 15, 9071

Received 27th September 2017, Accepted 19th October 2017 DOI: 10.1039/c7ob02408h

[rsc.li/obc](www.rsc.li/obc)

The enantioselective addition of 1-fluoro-1-nitro (phenylsulfonyl)methane to isatin-derived ketimines†

M. Urban, \mathbf{D}^{a} M. Franc,^a M. Hofmanová,^a I. Císařová^b and J. Veselý \mathbf{D}^{*a}

An asymmetric organocatalytic addition of fluorinated phenylsulfonylnitromethane to isatin-derived ketimines was developed. The reaction was efficiently catalyzed by a chiral tertiary amine, cinchonine. This methodology provides a new type of optically active compound with two adjacent quaternary carbon stereocenters in good yield (up to 96%), with moderate diastereoselectivity (up to 5.7 : 1 dr) and excellent enantioselectivity (up to 98/96% ee).

Introduction

Preparation of enantiomerically pure organic compounds bearing a quaternary stereogenic center is one of the most challenging research areas in asymmetric catalysis.¹ In addition, a quaternary 3-amino-2-oxindole moiety is a ubiquitous motif in natural products and pharmaceutical agents.² Among the methods developed for the preparation of such compounds, enantioselective addition of nucleophiles to isatin-derived ketimines is straightforward and probably the most efficient approach. Easy access to N -Boc isatin imines³ together with its modulate reactivity towards nucleophiles and the ability to easily remove Boc protecting group from the adducts have made those imines as attractive substrates for a number of catalytic addition reactions.⁴ Not only metal-based complexes (Pd, Rh, Ni, $)^5$ but also small organic molecules (H-bonding donors) can effectively catalyze these processes. To date a variety of organocatalytic addition reactions have been developed, including the Strecker reaction, the Mannich reaction and its vinylogous variant, the aza-Morita–Baylis–Hillman reaction, cycloadditions and others.⁶ Because of their impact in medicinal research and chemical biology, the development

of efficient methods for the selective introduction of fluorocontaining groups into bioactive molecules is an area of high interest.7 However, the enantioselective addition reactions of fluoro-containing nucleophiles to isatin-derived ketimines have rarely been described.⁸ Besides α-fluoro-bis(phenylsulfonyl)methane (FBSM),⁹ α-fluoro-nitro(phenylsulfonyl)methane (FNSM) has also been recognized as a versatile nucleophile in various organocatalytic reactions, as demonstrated by Prakash and Olah¹⁰ and others.¹¹ In this context, we herein report the highly enantioselective addition of α-fluoronitro(phenylsulfonyl)-methane to isatin-derived ketimines under mild organocatalytic conditions affording the fluoro-containing compounds bearing adjacent quaternary stereogenic centers. COMMUNICATION

The enantioselective addition of 1-fluoro-1-nitro
 $\frac{1}{2}$
 $\frac{1}{2}$

Results and discussion

At the outset we decided to study the nucleophilic addition of α-fluoro-nitro(phenylsulfonyl)methane (FNSM, 2a) to tert-butyl (1-benzyl-2-oxoindolin-3-ylidene)carbamate (1a) in the presence of cinchona alkaloids (Fig. 1). Initially, the reaction was performed with 1a and 2a in a 1 : 1.1 ratio at room temperature in toluene (Table 1, entry 1). It provided the desired adduct 3a/ 3a′ within a few hours in high yields (varying from 93 to 98%) and with moderate diastereoselectivity (up to $3.5:1$). Unfortunately, the obtained enantiomeric purity of 3a was unsatisfactory, and further screening of the reaction conditions was required. Initially, we tested the influence of temperature on the course of the reaction (Table SI 1, ESI†). Cinchonine (CN, Fig. 1) has shown the most promising results in terms of effectiveness and enantiocontrol in the model reaction performed at a lower temperature (−50 °C, entry 5). Full conversion of isatin-derived ketimine 1a was reached within 18 hours and adduct 3a was isolated in high yield (85%), with moderate diastereoselectivity (3.5 : 1) and excellent enantioselectivity of both diastereoisomers (96/95% ee). It is noteworthy that a further decrease in the reaction temperature led to a significant drop in the yield of 3a (Table 1, entry 6).

^aDepartment of Organic Chemistry, Charles University, Hlavova 8, 12843 Prague, Czech Republic. E-mail: jxvesely@natur.cuni.cz; http://orgchem.cz/vesely/

 b Department of Inorganic Chemistry, Charles University, Hlavova 8, 12843 Prague, Czech Republic

[†]Electronic supplementary information (ESI) available: Experimental procedure and spectral data for all prepared compounds with copies of the $^{1} \mathrm{H}$ NMR, $^{13} \mathrm{C}$ NMR and 19F NMR and HPLC chromatographs. CCDC 1575684. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7ob02408h

Fig. 1 Bifunctional organocatalysts screened in the reaction of ketimine 1a with FNSM 2a.

Table 1 Optimization studies on the reaction of ketimine 1a and FNSM 2a

	N -Boc	PhO ₂ S $\ddot{}$	NO ₂	Catalyst (20 mol%)	$Boc - NH$	NO ₂ SO_2Ph
Ω Ph 1a		2a		Toluene Temperature	3a/a'	O Ph
Entry	Catalyst	Temp. $\lceil \circ \text{C} \rceil$	Time [h]	Yield ^a $\lceil \% \rceil$	dr^{b}	ee $\left\lceil \frac{6}{10} \right\rceil$
1	QN	25	18	98	1.5:1	$-3/-3$
$\overline{2}$	QD	25	$\overline{4}$	94	1.2:1	19/62
3	CN	25	4	97	2:1	77/82
4	CD	25	4	93	1.6:1	$-60/-70$
5	CN	Ω	8	95	2.8:1	85/85
6	CN	-50	18	82	3.5:1	96/96
7	CN	-78	168	68	4:1	97/96

^a Isolated yield after column chromatography. b Determined by ¹⁹F NMR of the reaction mixture. ^c Determined by HPLC analysis of the purified product for two diastereoisomers.

Next, we assessed the influence of the solvent in the reaction of 1a and 2a (Table SI 1, ESI†). As shown in Table 1, apart from toluene, the addition reaction proceeded well in several nonpolar chlorinated and etheric solvents, but with lower diastereo- and enantioselectivity and also yield (Table 2, entries 1–6). On the other hand no reaction was observed in polar protic and aprotic solvents, such as methanol and DMSO. Screening of the solvents and temperature is summarized in the ESI† (Tables SI 1) in detail.

Next, we focused on the role of the catalyst in the enantiomeric addition of 2a to isatin-derived ketimine 1a with respect to efficiency and stereoselectivity (Table 2). Besides cinchona alkaloids, screened initially, chiral thioureas such as Takemoto's bifunctional catalyst A and Soós's catalysts B, C Sharpless bases and squaramides D, E were examined in the reaction between 1a and 2b (Fig. 1). Surprisingly, none of the above-mentioned organocatalysts catalysed the model reaction

Table 2 Solvent optimization studies on the reaction of ketimine 1a with FNSM 2a

N ^{-Boc} $\ddot{}$ Ph		PhO ₂ S ₂ NO ₂	CN (20 mol)	$Boc - NH$	NO ₂ SO_2 Ph
			Toluene -50° C		റ
	1a (1.0 eq.)	2a (1.1 eq.)		3a/a'	∙Ph
Entry	Solvent	Time $[h]$	Yield ^a [%]	dr ^b	ee $\left\lceil \frac{6}{6} \right\rceil$
1	Toluene	18	85	3.5:1	96/95
2	DCM	19	57	2:1	79/86
3	CHCl ₃	19	84	3:1	88/89
4	THF	96	65	1.9:1	22/35
5	MTBE	96	75	3:1	65/63
6	EtOAc	96	18	2.8:1	61/63
7	MeOH	96			
8^d	DMSO	96			

^a Isolated yield after column chromatography. $\frac{b}{c}$ Determined by ¹⁹F NMR of the reaction mixture. ^c Determined by HPLC analysis of the purified product for two diastereoisomers. d Reaction at 25 °C.

as effectively as cinchonine (Table 3, entries 1–6, more details in the ESI†). We also screened the dependence of the reaction efficiency and selectivity on the loading of catalysts. Interestingly, no significant change in selectivity and yield was observed when 5 mol% of the catalyst was used. On the other hand a lower catalyst loading dramatically influenced the reaction time (Table 3, entry 9). A reduced yield was observed, when 2.5 mol% of the catalyst was used, nevertheless, the selectivity of the reaction was retained (Table 3, entry 10).

Once we uncovered the optimized reaction conditions for the enantioselective nucleophilic addition of FNSM to isatinderived ketimine 1a, we proceeded with the scope of the process (Scheme 1). First we studied the reactivity of FNSM towards ketimines 1a–1h having different protecting groups on

Table 3 Catalyst optimization studies on the reaction of ketimine 1a and FNSM 2a

	$N - Boc$ PhO ₂ S ₂ ÷ Ph 1a(1.0eq.)	NO ₂ 2a(1.1eq.)	CN (20 mol%) Toluene -50° C	$Boc - N$ 3a/a'	NO ₂ SO_2 Ph Ρh
Entry	Catalyst	Time [h]	Yield ^{<i>a</i>} [%]	dr^{b}	ee $\left\lceil \frac{6}{6} \right\rceil$
1	Takemoto's A	18	96	3.3:1	$-27/ -25$
2	Soós's with ON B	18	96	3.7:1	64/56
3	Soós's with CD D	20	96	5.9:1	70/62
4	(DHO)2AON	24	90	4:1	35/28
5	Rawal's with ON E	24	89	4.3:1	18/47
6	Rawal's with CN F	168	76	4.5:1	$-75/-16$
7	$CN(10 \text{ mol})$	24	80	3.5:1	96/95
8	$CN(5 \text{ mol})\%$	36	85	3.7:1	96/96
9	CN (2.5 mol%)	48	68	3.6:1	96/95
10		168			

^a Isolated yield after column chromatography. b Determined by ¹⁹F NMR of the reaction mixture. ^c Determined by HPLC analysis of the purified product for two diastereoisomers.

Scheme 1 Substrate scope of ketimines 1a–p and fluorinated sulfones 2a–3 in the organocatalytic alkylation reaction. Isolated yield after column chromatography. Determined by ^{19}F NMR of the reaction mixture. Determined by HPLC analysis of the purified product for two diastereoisomers.

the nitrogen of the 3-amino-2-oxindole moiety. The desired products 3a/3a′–3g/3g′ were prepared in good yields with high to excellent enantioselectivity (93–99% ee for the major diastereoisomer and 93–95% ee for the minor diastereoisomer). A moderate degree of diastereoselectivity $(3:1$ to $5:1)$ was obtained in the reaction of 2a with most ketimines 1a–1h. The highest diastereocontrol was observed in the reaction with Boc

protected ketamine $1d$ (dr $5:1$). On the other hand, the reaction between 2a and ketimine 1h with a tosyl protecting group proceeded with low diastereo- (dr 1 : 1.9) and enantioselectivity (70/75% ee); the corresponding adduct 3h was isolated in low yield (31%) as a mixture of diastereoisomers.

Subsequently, we focused on ketimine derivatives with different substitutions on the aromatic ring (1i-1o). Substitution at the 5-position on the aromatic ring of ketimines led to the formation of the corresponding products 3i–3l in high yield (81–95%) and excellent enantiomeric excess for both enantiomers (major diastereoisomer 96–98% ee, minor diastereoisomer 94–96% ee). On the other hand, diastereoselectivity reached moderate values (dr 3.7 : 1–5.7 : 1). A significant change in reactivity was observed, when ketimine 1o substituted at the 4-position with bromine was used. The formation of the corresponding adduct 3o was not observed even after a prolonged reaction time (96 h). This can be probably caused due to the higher steric hindrance of the imine group with the bulky bromine substituent at the 4-position, because all other adducts 3l, 3m and 3n having bromine atoms at positions 5-,6- and 7 were obtained in high yields with good levels of enantioselectivity. Replacement of the Boc group for the Cbz group on the nitrogen of the imine moiety led to a considerable drop in enantioselectivity (75/64% ee). The corresponding product 3p was obtained in 88% yield and with good diastereoselectivity (dr 5.1 : 1).

Next, we turned our attention to other types of fluorinated sulfones 2b–e. The nitro group was replaced by various functional groups such as nitrile, ester, acetyl, phenylcarbonyl groups, etc. In all cases significantly lower reactivity towards 1a under the optimized reaction conditions (QN, toluene, −50 °C) was observed. Due to this reason we approached to set up the reactions with appropriate sulfones 2b–e at 25 °C. It showed up that apart from the model reaction between 1a and 2a, reactions proceeded smoothly only with 2-fluoro-2-(phenylsulfonyl)acetonitrile (2b), affording product 3q in high yield (94%) with low diastereoselectivity (dr 2.3:1) and unsatisfactory enantioselectivity (3/32% ee). Both the subsequential turn of temperature down to −50 °C and also the change of the solvent from toluene to DCM did not lead to improved selectivity of the reaction (dr 2.9 : 1, 30/59% ee, Scheme 1).

Next, the potential versatility of a phenylsulfanyl group as a traceless activating group was verified. The standard procedure for the removal of phenylsulfonyl group with magnesium failed.¹² Nevertheless, desulfonation of enantioenriched adducts 3a was accomplished using AIBN as a radical agent in the presence of BNAH as a hydrogen transfer agent (Table 4).¹³ α-Fluoronitro alkane 4 was isolated in 96% yield as a mixture of diastereoisomers $(dr = 1:1)$ with the enantioselectivity retained (95/96% ee). Furthermore, Boc deprotection of 4a and 4b was successfully performed using TFA (Table 4).¹⁴ The corresponding amines 5a and 5b were obtained in good yields 48% and 47%, respectively, with excellent enantiomeric excess (97/96%).

To determine the absolute configuration of alkylated products 3a, we performed single-crystal X-ray diffraction analysis of 3a, obtained from the reaction of ketimine 1a with fluoro-

Table 4 Further transformation of alkylation product 3a

^{*a*} Reaction under microwave radiation. $\frac{b}{b}$ Isolated yield after column chromatography. ^c Determined by HPLC analysis of the purified product for two diastereoisomers.

Fig. 2 View of the molecular structure of 3a, the displacement ellipsoids at the 30% probability level. The dashed line indicates the intramolecular hydrogen bond N2–H2⋯O7, N2⋯O7 2.894(2) Å; angle on H2 138.7°.¹⁵

Fig. 3 Proposed transition state.

nitro(phenylsulfonyl)methane 2a (Fig. 2). As shown in Fig. 3, the C3 and C1' stereogenic centres have the (S) and (R) absolute configuration, respectively.

On the basis of the crystallographic analysis of 3a an envisaged transition-state structure was proposed. As shown in Fig. 3, cinchonine can act as a dual catalyst and activate both reaction substrates via hydrogen bonding. The tertiary amine of the chinuclidine moiety deprotonates FNSM forming the corresponding anionic nucleophile that preferentially approaches the ketimine from the Re-face due to the H-bonding of the ketimine and hydroxyl groups of cinchonine.

Conclusion

In summary, we have explored the stereoselective fluoroalkylation of isatin-derived ketimines with α-fluoronitro(phenylsulfonyl) methane. The reaction was effectively catalyzed with cinchonine furnishing the corresponding adducts in good yields, with good diastereo- (dr up to $6:1$) and high enantioselectivity (up to 98% ee). The reactivity of isatin-derived ketimines toward other fluorinated compounds was also studied. The outcome of the reported procedure was demonstrated by the preparation of rarely available β-fluoro-β-nitro amines using a desulfonylation/Boc-removal reaction sequence. Further studies and synthetic applications based on the reported methodology are currently ongoing in our laboratory.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Jan Veselý gratefully acknowledges the Czech Science Foundation (no. 16-23597S) for financial support. Michal Urban thanks the Charles University Grant Agency (grant number 392315) for financial support. We also thank Dr Hybelbauerová for NMR service and Dr Štícha for MS analysis.

Notes and references

1 For excellent reviews and perspective articles on the catalytic asymmetric construction of quaternary stereogenic centers, see: (a) Y. Liu, S.-J. Han, W.-B. Liu and B. M. Stoltz, Acc. Chem. Res., 2015, 48, 740; (b) K. W. Quasdorf and L. E. Overman, Nature, 2014, 516, 181; (c) I. Marek, Y. Minko, M. Pasco, T. Mejuch, N. Gilboa, H. Chechik and J. P. Das, J. Am. Chem. Soc., 2014, 136, 2682; (d) A. Y. Hong and B. M. Stoltz, Eur. J. Org. Chem., 2013, 2745; (e) J. P. Das and I. Marek, Chem. Commun., 2011, 47, 4593; (f) M. Shimizu, Angew. Chem., Int. Ed., 2011, 50, 5998; (g) B. Wang and Y. Q. Tu, Acc. Chem. Res., 2011, 44, 1207; (h) C. Hawner and A. Alexakis, *Chem. Commun.*, 2010, 46, 7295.

- 2 (a) T. Oost, G. Backfisch, S. Bhowmik, M. M. van Gaalen, H. Geneste, W. Hornberger, W. Lubisch, A. Netz, L. Unger and W. Wernet, Bioorg. Med. Chem. Lett., 2011, 21, 3828; (b) M. Rottmann, C. McNamara, B. S. K. Yeung, M. C. S. Lee, B. Zou, B. Russell, P. Seitz, D. M. Plouffe, N. V. Dharia, J. Tan, S. B. Cohen, K. R. Spencer, G. Gonzalez-Paez, L. Lakshiminarayana, A. Goh, R. Suwanarusk, T. Jegla, E. K. Schmitt, H.-P. Beck, R. Brun, F. Nosten, L. Renia, V. Dartois, T. H. Keller, D. A. Fidock, E. A. Winzeler and T. T. Diagana, Science, 2010, 329, 1175; (c) V. V. Vintonyak, K. Warburg, H. Kruse, S. Grimme, K. Hübel, D. Rauh and H. Waldmann, Angew. Chem. Int. Ed., 2010, 49, 5902; (d) T. Shimazaki, M. Iijima and S. Chaki, Eur. J. Pharmacol., 2006, 543, 63; (e) A. K. Ghosh, G. Schiltz, R. S. Perali, S. Leshchenko, S. Kay, D. E. Walters, Y. Koh, K. Maeda and H. Mitsuya, Bioorg. Med. Chem. Lett., 2006, 16, 1869; (f) K. Bernard, S. Bogliolo and J. Ehrenfeld, Br. J. Pharmacol., 2005, 144, 1037; (g) G. Gilles and S. L. Claudine, Stress, 2003, 6, 199; (h) M. Ochi, K. Kawasaki, H. Kataoka and Y. Uchio, Biochem. Biophys. Res. Commun., 2001, 283, 1118; (i) L. Chevolot, A. M. Chevolot, M. Gajhede, C. Larsen, U. Anthoni and C. Christophersen, J. Am. Chem. Soc., 1985, 107, 4542.
- 3 W. J. Yan, D. Wang, J. C. Feng, P. Li, D. Zhao and R. Wang, Org. Lett., 2012, 14, 2512.
- 4 For recent reviews from the area of enantioselective additions to N-carbamoylimines, see: (a) J. Veselý and R. Rios, Chem. Soc. Rev., 2014, 43, 611; (b) E. Marcantoni and M. Petrini, Adv. Synth. Catal., 2016, 358, 3657.
- 5 (a) Z. Liu, X. Feng and H. Du, Org. Lett., 2012, 14, 3154; (b) T. Rajasekaran, G. Karthik, B. Sridhar, S. K. Kumar and B. V. S. Reddy, Eur. J. Org. Chem., 2014, 2221; (c) L.-Y. Mei, X.-Y. Tang and M. Shi, Org. Biomol. Chem., 2014, 12, 1149; (d) C. S. Marques and A. J. Burke, Eur. J. Org. Chem., 2016, 806; (e) S. Nakamura, K. Hyodo, M. Nakamura, D. Nakane and H. Masuda, Chem. – Eur. J., 2013, 19, 7304; (f) T. Arai, E. Matsumura and H. Masu, Org. Lett., 2014, 16, 2768; (g) T. Arai, K. Tsuchiya and E. Matsumura, Org. Lett., 2015, 17, 2416; (h) H. Zheng, X. Liu, C. Xu, Y. Xia, L. Lin and X. Feng, Angew. Chem., Int. Ed., 2015, 54, 10958.
- 6 (a) N. Hara, S. Nakamura, M. Sano, R. Tamura, Y. Funahashi and N. Shibata, Chem. – Eur. J., 2012, 18, 9276; (b) Y.-L. Liu and J. Zhou, *Chem. Commun.*, 2013, 49, 4421; (c) D. Wang, J. Liang, J. Feng, K. Wang, Q. Sun, L. Zhao, D. Li, W. Yan and R. Wang, Adv. Synth. Catal., 2013, 355, 548; (d) F. L. Hu, Y. Wei, M. Shi, S. Pindi and G. Li, Org. Biomol. Chem., 2013, 11, 1921; (e) T.-Z. Li, X.-B. Wang, F. Sha and X.-Y. Wu, Tetrahedron, 2013, 69, 7314; (f) X.-B. Wang, T.-Z. Li, F. Sha and X.-Y. Wu, Eur. J. Org. Chem., 2014, 739; (g) J. George, B. Sridhar and B. V. S. Reddy, Org. Biomol. Chem., 2014, 12, 1595; (h) X. Zhao, T.-Z. Li, J.-Y. Qian, F. Sha and X.-Y. Wu, Org. Biomol. Chem., 2014, 12, 8072; (i) T.-Z. Li, X.-B. Wang, F. Sha and X.-Y. Wu, J. Org. Chem., 2014, 79, 4332;

 (j) Y. Guo, Y. Zhang, L. Qi, F. Tian and L. Wang, RSC Adv., 2014, 4, 27286; (k) Z. Tang, Y. Shi, H. Mao, X. Zhu, W. Li, Y. Cheng, W.-H. Zheng and Ch. Zhu, Org. Biomol. Chem., 2014, 12, 6085; (l) A. Kumar, J. Kaur, S. S. Chimni and A. K. Jassal, RSC Adv., 2014, 4, 24816; (m) Y.-H. Wang, Y. L. Liu, Z.-Y. Cao and J. Zhou, Asian J. Org. Chem., 2014, 3, 429; (n) S. Nakamura, S. Takahashi, D. Nakane and H. Masuda, Org. Lett., 2015, 17, 106; (o) A. Kumar, V. Sharma, J. Kaur, N. Kumar and S. S. Chimni, Org. Biomol. Chem., 2015, 13, 5629; (p) Y. Yoshida, M. Sako, K. Kishi, H. Sasai, S. Hatakeyama and S. Takizawa, Org. Biomol. Chem., 2015, 13, 9022; (q) S. Nakamura and S. Takahashi, Org. Lett., 2015, 17, 2590; (r) O. D. Engl, S. P. Fritz and H. Wennemers, Angew. Chem., Int. Ed., 2015, 54, 8193; (s) M. Montesinos-Magraner, C. Vila, R. Cantón, G. Blay, I. Fernández, M. C. Muñoz and J. R. Pedro, Angew. Chem., Int. Ed., 2015, 54, 6320; (t) T. Liu, W. Liu, X. Li, F. Peng and Z. Shao, J. Org. Chem., 2015, 80, 4950; (u) X. Bao, B. Wang, L. Cui, G. Zhu, Y. He, J. Ou and Y. Song, Org. Lett., 2015, 17, 5168; (v) M.-X. Zhao, L. Jing, H. Zhou and M. Shi, RSC Adv., 2015, 5, 75648; (w) K. Zhao, T. Shu, J. Jia, G. Raabe and D. Enders, Chem. – Eur. J., 2015, 21, 3933; (x) Y. Zhu, E. G. Zhang, C. Luo, X. Li and J.-P. Cheng, Tetrahedron, 2015, 71, 4090; (y) F. I. Amr, C. Vila, G. Blay, M. C. Muñoz and J. R. Pedro, Adv. Synth. Catal., 2016, 358, 1583; (z) C. Vila, F. I. Amr, G. Blay, M. C. Muñoz and J. R. Pedro, Chem. – Asian J., 2016, 11, 1532; (aa) K. Zhao, Y. Zhi, X. Li, R. Puttreddy, K. Rissanen and D. Enders, Chem. Commun., 2016, 52, 2249; (ab) B. X. Feng, J.-D. Yang, J. Li and X. Li, Tetrahedron Lett., 2016, 57, 3457; (ac) P. Cheng, W. Guo, P. Chen, Y. Liu, X. Du and C. Li, Chem. Commun., 2016, 52, 3418; (ad) M. M. Magraner, C. Vila, A. R. Patiño, G. Blay, I. Fernández, M. C. Muñoz and J. R. Pedro, ACS Catal., 2016, 6, 2689; (ae) W. Guo, Y. Liu and C. Li, Org. Lett., 2017, 19, 1044; (af) S. Hajra and B. Jana, Org. Lett., 2017, 19, 4778; (ag) J. Dai, D. Xiong, T. Yuan, J. Liu, T. Chen and Z. Shao, Angew. Chem., Int. Ed., 2017, 56, 12697; (ah) C. Cheng, X. Lu, L. Ge, J. Chen, W. Cao, X. Wu and G. Zhao, Org. Chem. Front., 2017, 4, 101. Open Access Article. Published on 19 October 2017. Downloaded on 12/20/2024 11:33:18 PM. This article is licensed under a [Creative Commons Attribution-NonCommercial 3.0 Unported Licence.](http://creativecommons.org/licenses/by-nc/3.0/) **[View Article Online](https://doi.org/10.1039/c7ob02408h)**

- 7 (a) K. Muller, C. Faeh and F. Diederich, Science, 2007, 317, 1881; (b) S. Purser, P. R. Moore, S. Swallow and V. Gouverneur, Chem. Soc. Rev., 2008, 37, 320; (c) W. K. Hagmann, J. Med. Chem., 2008, 51, 4359; (d) Fluorine in Bioorganic and Medicinal Chemistry, ed. I. Ojima, Wiley-Blackwell, Chichester, U.K., 2009; (e) D. J. O'Hagan, Fluorine Chem., 2010, 131, 1071; (f) G. Valero, X. Companyo and R. Rios, *Chem. - Eur. J.*, 2011, 17, 2018; (g) E. P. Gillis, K. J. Eastman, M. D. Hill, D. J. Donnelly and N. A. Meanwell, J. Med. Chem., 2015, 58, 8315; (h) Y. Zhou, J. Wang, Z. Gu, S. Wang, W. Zhu, J. L. Acena, V. A. Soloshonok, K. Izawa and H. Liu, Chem. Rev., 2016, 116, 422.
- 8 (a) J. Liu, L. Zhang and J. Hu, Org. Lett., 2008, 10(23), 5377; (b) J. Liu and J. Hu, Chem. - Eur. J., 2010, 16, 11443; (c) J.-S. Yua and J. Zhou, Org. Biomol. Chem., 2015, 13,

10968; (d) X. Liu, J. Zhang, L. Zhao, S. Ma, D. Yang, W. Yan and R. Wang, J. Org. Chem., 2015, 80, 12651; (e) J.-S. Yu and J. Zhou, Org. Chem. Front., 2016, 3, 298; (f) X.-L. Zeng, Z.-Y. Deng, C. Liu, G. Zhao, J.-H. Lin, X. Zheng and J.-Ch. Xiao, J. Fluorine Chem., 2017, 193, 17. For other fluoroalkylation reagents, see: Ch. Ni, M. Hu and J. Hu, Chem. Rev., 2015, 115, 765.

9 For selected studies, see: (a) T. Fukuzumi, N. Shibata, M. Sugiura, H. Yasui, S. Naka-mura and T. Toru, Angew. Chem., Int. Ed., 2006, 45, 4973; (b) S. Mizuta, N. Shibata, Y. Goto, T. Furukawa, S. Nakamura and T. Toru, *J. Am.* Chem. Soc., 2007, 129, 6394; (c) T. Furukawa, N. Shibata, S. Mizuta, S. Nakamura, T. Toru and M. Shiro, Angew. Chem., Int. Ed., 2008, 47, 8051; (d) Ch. Ni, L. Zhang and J. Hu, J. Org. Chem., 2008, 73, 5699; (e) A.-N. Alba, X. Company, A. Moyano and R. Rios, Chem. – Eur. J., 2009, 15, 7035; (f) F. Ullah, G.-L. Zhao, L. Deiana, M. Zhu, P. Dziedzic, I. Ibrahem, P. Hammar, J. Sun and A. Córdova, Chem. – Eur. J., 2009, 15, 10013; (g) X. Shen, L. Zhang, Y. Zhao, L. Zhu, G. Li and J. Hu, Angew. Chem., Int. Ed., 2011, 50, 2588; (h) W. Yang, X. Wei, Y. Pan, R. Lee, B. Zhu, H. Liu, L. Yan, K.-W. Huang, Z. Jiang and Ch.-H. Tan, Chem. – Eur. J., 2011, 17, 8066; (i) T. Furukawa, J. Kawazoe, W. Zhang, T. Nishimine, E. Tokunaga, T. Matsumoto, M. Shiro and N. Shibata, Angew. Chem., Int. Ed., 2011, 50, 9684; (j) S.-L. Zhang, H.-X. Xie, J. Zhu, H. Li, X.-S. Zhang, J. Li and W. Wang, Nat. Commun., 2011, 211(2), 1; (k) W. Huang, Ch. Ni, Y. Zhao and J. Hu, New J. Chem., 2013, 37, 1684; (l) K. Matsuzaki, T. Furukawa, E. Tokunaga, Communication Open Access Article 2017. Downloaded on 19 October 2017. Downloaded on 12/2024 12:33:19 PM. This article is licensed under a Creative Commons Access Articles. The expectation-NonCommons Access Articles. The

T. Matsumoto, M. Shiro and N. Shibata, Org. Lett., 2013, 15(13), 3282.

- 10 (a) G. K. S. Prakash, X. Zhao, S. Chacko, F. Wang, H. Vaghoo and G. A. Olah, Beilstein J. Org. Chem., 2008, 4, 17; (b) G. K. S. Prakash, F. Wang, T. Stewart, T. Mathew and G. A. Olah, Proc. Natl. Acad. Sci. U. S. A., 2009, 106, 4090.
- 11 For selected studies, see: (a) F. Ullah, G. L. Zhao, L. Deiana, M. Zhu, P. Dziedzic, I. Ibrahem, P. Hammar, J. Sun and A. Cordova, Chem. – Eur. J., 2009, 15, 10013; (b) M. Kamlar, N. Bravo, A.-N. R. Alba, S. Hybelbauerova, I. Cisarova, J. Vesely, A. Moyano and R. Rios, Eur. J. Org. Chem., 2010, 5464; (c) Y. Pan, Y. Zhao, T. Ma, Y. Yang, H. Liu, Z. Jiang and C.-H. Tan, Chem. – Eur. J., 2010, 16, 779; (d) H. W. Moon and D. Y. Kim, Bull. Korean Chem. Soc., 2012, 33(9), 2845; (e) M. Kamlar, P. Putaj and J. Veselý, Tetrahedron Lett., 2013, 54, 2097; (f) G. K. S. Prakash, L. Gurung, P. V. Jog, S. Tanaka, T. E. Thomas, N. Ganesh, R. Haiges, T. Mathew and G. A. Olah, Chem. – Eur. J., 2013, 19, 3579.
- 12 Y. Kazuta, A. Matsuda and S. Shuto, J. Org. Chem., 2002, 67, 1669.
- 13 Y. Seo, H. Kim, D. W. Chae and Y. G. Kim, Tetrahedron: Asymmetry, 2014, 25, 625.
- 14 M. Montesinos-Magraner, C. Vila, R. Cantón, G. Blay, I. Fernández, M. C. Muñoz and J. R. Pedro, Angew. Chem., Int. Ed., 2015, 54, 6320.
- 15 X-ray crystallographic data have been deposited with the Cambridge Crystallographic Data Centre (CCDC) under deposition numbers CCDC 1575684† for 3a.