



Cite this: *Org. Biomol. Chem.*, 2017, **15**, 1268

Correction: Facile functionalization at the C4 position of pyrimidine nucleosides *via* amide group activation with (benzotriazol-1-yloxy) tris(dimethylamino)phosphonium hexafluorophosphate (BOP) and biological evaluations of the products

Hari K. Akula,^{a,b} Hariprasad Kokatla,^a Graciela Andrei,^c Robert Snoeck,^c Dominique Schols,^c Jan Balzarini,^c Lijia Yang^a and Mahesh K. Lakshman*^{a,b}

Correction for 'Facile functionalization at the C4 position of pyrimidine nucleosides *via* amide group activation with (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP) and biological evaluations of the products' by Hari K. Akula, *et al.*, *Org. Biomol. Chem.*, 2017, DOI: 10.1039/c6ob02334g.

DOI: 10.1039/c7ob90013a
rsc.li/obc

The authors regret that Table 6 footnote *b* was given incorrectly. The correct footnote is given below.

^bEC₅₀ = 50% effective concentration or compound concentration required to inhibit virus replication by 50%.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aDepartment of Chemistry and Biochemistry, The City College of New York, 160 Convent Avenue, New York, New York, 10031, USA. E-mail: mlakshman@ccny.cuny.edu

^bThe Ph.D. Program in Chemistry, The Graduate Center of the City University of New York, New York, New York 10016, USA

^cLaboratory of Virology and Chemotherapy, Rega Institute for Medical Research, Herestraat 49, Postbus 1043, 3000 Leuven, Belgium

