Cite this: *Org. Biomol. Chem.*, 2017, **15**, 7474

Correction: Metal-free oxidative ring contraction of benzodiazepinones: an entry to quinoxalinones†

Hasan Mtiraoui,^{‡a} Kevin Renault,^{‡b} Morgane Sanselme,^{c,d} Moncef Msaddek,^{*a} Pierre-Yves Renard^b and Cyrille Sabot^{*b}

DOI: 10.1039/c7ob90126g

rsc.li/obc

Correction for 'Metal-free oxidative ring contraction of benzodiazepinones: an entry to quinoxalinones' by Hasan Mtiraoui, et al., *Org. Biomol. Chem.*, 2017, **15**, 3060–3068.

The authors regret that there were errors in some of the structures. The reactions to obtain compounds **2h** and **2i** were in fact carried out from 7-methoxy-benzodiazepinones **1h** and **1i**, and not from the regioisomer 8-methoxy-benzodiazepinones as reported. Consequently, the 6-methoxyquinoxalinones **2h** and **2i** were formed (and not the regioisomer 7-methoxyquinoxalinones). Subsequent compounds **5**, **6** and **8** (Fig. 2, and Scheme 4) prepared from **2i** should be modified accordingly.

The correct structures and corrected Fig. 2 and Scheme 4 are shown below.

^aUniversité Monastir, Laboratory of Heterocyclic Chemistry Natural Products and Reactivity/CHPNR, Department of Chemistry, Faculty of Science of Monastir, 5000 Monastir, Tunisia

^bNormandie Univ, CNRS, UNIROUEN, INSA Rouen, COBRA (UMR 6014), 76000 Rouen, France. E-mail: cyrille.sabot@univ-rouen.fr

^cNormandie Univ, France

^dUNIROUEN, Laboratoire SMS EA3233, 1 rue Tesniere, France, F-76821 Mont Saint Aignan, France

† Electronic supplementary information (ESI) available: ¹H and ¹³C NMR, 2D NMR data in support of the formation of 6-methoxyquinoxalinones **2h** and **2i**, photo-physical data for all new compounds, and X-ray crystallographic data (CIF files) for compound **2a**. CCDC 1519664. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7ob00205j

‡ These authors contributed equally to this work.

Fig. 2 (a) Preparation of the hydroxyl-containing quinoxalinone **5**. (b) Fluorescence emission spectrum of quinoxalinone **2i** (λ_{ex} 366 nm, λ_{em} 475 nm) in PBS 7.4 at 25 °C, before (dashed line) and after (black line) the addition of NaBH_4 .

Scheme 4 Application to the preparation of the fluorescently labelled cholesterol derivative **8**.

In addition, compounds **2h**, **2i**, **5**, **6** and **8** were named incorrectly in the Experimental section and should also be amended accordingly.

The original ESI has been updated with a corrected version showing the corrected structures and including 2D NMR data in support of the formation of 6-methoxyquinoxalinones **2h** and **2i**.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.