



Cite this: *Inorg. Chem. Front.*, 2017, **4**, 1200

# Stoichiometry of lanthanide(III) complexes with tripodal aminophosphonic ligands – a new solution to an old problem†

Rafał Janicki,<sup>\*a</sup> Joanna Gałęzowska<sup>\*b</sup> and Anna Mondry<sup>a</sup>

The  $\text{Eu}^{3+}$  and  $\text{Gd}^{3+}$  complexes with an *N*-(methylene-2-pyridine)-*N,N*-di(methylenephosphonate) ligand ( $\text{H}_4\text{NP}_2\text{py}$ ), an analogue of nitrilotris(methylphosphonic) acid ( $\text{H}_6\text{NTP}$ ), were synthesized and structurally characterized by X-ray single crystal diffraction. The determined crystal structures ( $[\text{C}(\text{NH}_2)_3][\text{Ln}(\text{NP}_2\text{py})_2] \cdot 12\text{H}_2\text{O}$ ) are the first example of a monomeric  $\text{Ln}^{3+}$  complex encapsulated by two tripodal aminophosphonic ligands. Each of the  $\text{NP}_2\text{py}$  anions coordinates to  $\text{Ln}^{3+}$  through two oxygen atoms from each monodentate phosphonic group, amine nitrogen and pyridine nitrogen atoms, filling thus 8 coordination sites of  $\text{Ln}^{3+}$ . The luminescence properties of  $[\text{C}(\text{NH}_2)_3][\text{Eu}(\text{NP}_2\text{py})_2] \cdot 12\text{H}_2\text{O}$  crystals were studied and compared with those of  $\text{Eu}-\text{NP}_2\text{py}$  complexes in solution. Speciation analysis of  $\text{Ln}-\text{NP}_2\text{py}$  complexes ( $\text{Ln} : \text{NP}_2\text{py} = 1 : 2$ ), performed by luminescence and potentiometric methods, showed that both  $[\text{Ln}(\text{NP}_2\text{py})]^-$  and  $[\text{Ln}(\text{NP}_2\text{py})_2]^{5-}$  species may exist in solution. However, the formation of the latter one occurs in alkaline solutions at pH as high as 8. By implementing the Specific Ion Interaction Theory (SIT) it was possible to calculate the thermodynamic stability constants of the  $[\text{Eu}(\text{NP}_2\text{py})]^-$  and  $[\text{Eu}(\text{NP}_2\text{py})_2]^{5-}$  complexes. The corresponding  $\log \beta_{\text{EuI}}^0$  and  $\log \beta_{\text{EuL}_2}^0$  values are  $16.3 \pm 0.11$  and  $19.5 \pm 0.15$ , respectively.

Received 10th April 2017,

Accepted 12th May 2017

DOI: 10.1039/c7qi00191f

rsc.li/frontiers-inorganic

## Introduction

The chemistry of metal-phosphonates has attracted increasing attention over the last few decades due to the variety of pro-

erties that can be introduced *via* a phosphonic moiety.<sup>1</sup> (Poly) aminopolyphosphonates are very strong complexing agents for alkaline earth, transition and rare earth metal ions.<sup>2</sup> Owing to the high basicity of the oxygen atoms,<sup>3,4</sup> phosphonates interact with cations more strongly than their carboxylic analogues and are able to create more stable complexes.<sup>5,6</sup> Phosphonate complexes with luminescent, paramagnetic lanthanide ions<sup>7,8</sup> play a significant role as promising phosphors for new technologies and potential contrast agents in magnetic resonance imaging (MRI)<sup>9</sup> as well as paramagnetic shift reagents for biologically important cations (*e.g.*  $\text{Na}^+$ ).<sup>10</sup> Owing to its good pharmacokinetics, biodistribution and bone-seeking properties the  $^{153}\text{Sm}-\text{EDTP}$  complex,<sup>1a,6e-h</sup> known as QUADRAMET®, was approved by the US FDA for use of painful skeletal metastases in humans.<sup>1d,11</sup>

In general, polyaminopolyphosphonic acids are the analogues of the nitrilotris(methylenephosphonic) acid (herein after referred to as NTP). To date the NTP ligand has found many applications<sup>12</sup> driven in many cases by coordination properties of this ligand in  $\text{Ln}^{3+}$  complexes.<sup>13</sup> Because the majority of its applications are related to the NTP properties in aqueous solution, it seems particularly important to study its properties in this medium. However, the formation of the  $\text{Ln}^{3+}$  complexes with ligands of the NTP type in solution remains still not fully understood. For example it is unclear whether bis(nitrilotris(methylenephosphonato))lanthanide complexes are

<sup>a</sup>Faculty of Chemistry, University of Wrocław, F. Joliot-Curie 14, 50-383 Wrocław, Poland. E-mail: rafal.janicki@chem.uni.wroc.pl

<sup>b</sup>Department of Inorganic Chemistry, Wrocław Medical University, Borowska 211A, 50-556 Wrocław, Poland. E-mail: joanna.galezowska@umed.wroc.pl

†Electronic supplementary information (ESI) available: Table 1S. The fitting parameters for eqn (3), determined from the spectra of the  $^5\text{D}_0 \rightarrow ^7\text{F}_{1,2,4}$  transitions of the  $[\text{Eu}(\text{NP}_2\text{py})_2]^{5-}$  complex in aqueous solution; Table 2S. Selected luminescent data of  $\text{Eu}-\text{NP}_2\text{py}$  and  $\text{Tb}-\text{NP}_2\text{py}$  systems; Table 3. The complex formation constants of  $\text{NP}_2\text{py}$  with  $\text{Sm}^{3+}$ ,  $\text{Eu}^{3+}$ ,  $\text{Gd}^{3+}$  and  $\text{Tb}^{3+}$  at 25 °C,  $I = 0.1 \text{ mol dm}^{-3}$  ( $\text{NaClO}_4$ ), in 1 : 1 and 1 : 2 molar ratios  $\text{Ln} : \text{L}$ ; Table 4S. Stability constants of  $\text{NP}_2\text{py}$  with  $\text{Eu}^{3+}$  at 25 °C in different ionic strengths (KCl); Fig. 1S. Phosphorescence spectrum of  $[\text{C}(\text{NH}_2)_3][\text{Gd}(\text{NP}_2\text{py})_2] \cdot 12\text{H}_2\text{O}$  crystals at 77 K; Fig. 2S. The plot of luminescence quantum yields of  $\text{Tb} : \text{NP}_2\text{py}$  1 : 1 and 1 : 2 systems *versus* pH; Fig. 3S. UV spectra of the  $\text{NP}_2\text{py}$  ligand and  $\text{Tb} : \text{NP}_2\text{py}$  complex at various pH; Fig. 4S. Plot of  $q_{\text{H}_2\text{O}}$  *versus* pH of solution for the  $\text{Eu}-\text{NP}_2\text{py}$  complex with  $\text{Eu} : \text{NP}_2\text{py}$  ratios of 1 : 1 and 1 : 2. The  $\Delta q_{\text{H}_2\text{O}}$  is the difference between  $q_{\text{H}_2\text{O}}$  for 1 : 1 and 1 : 2 solutions at the same pH; Fig. 5S. Plot of  $q_{\text{H}_2\text{O}}$  *versus* pH of solution for the  $\text{Tb}-\text{NP}_2\text{py}$  complex with  $\text{Tb} : \text{NP}_2\text{py}$  ratios of 1 : 1 and 1 : 2; Fig. 6S. Species distribution curves of  $\text{H}_6\text{NP}_2\text{py}$  acid (L). Charges omitted for brevity; Fig. 7S. Species distribution curves for the  $\text{Ln}^{3+}-\text{NP}_2\text{py}$  complexes as a function of pH for a 1 : 1  $\text{Ln} : \text{L}$  molar ratio (A) and 1 : 2  $\text{Ln} : \text{L}$  (B). Equimolar complexes are marked in blue, biscomplexes in red color,  $I = 0.1 \text{ M NaClO}_4$ , 25 °C. CCDC 1541894 and 1541895. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7qi00191f



formed in solution at all.<sup>14–16</sup> Moreover no crystal structures of monomeric Ln<sup>3+</sup> complexes with tripodal aminophosphonic ligands have been reported so far. The high negative charge of the NTP ligand (–6) and the presence of rather bulky phosphonic groups are probably the main factors destabilizing the [Ln(NTP)<sub>2</sub>]<sup>9–</sup> complexes in solution.

The aim of this paper is to study the chemical properties of a modified NTP ligand – *N*-(methylene-2-pyridine)-*N,N*-di(methylenephosphonic) acid (hereinafter referred to as NP<sub>2</sub>py – Scheme 1) to establish finally whether, [Ln(NP<sub>2</sub>py)<sub>2</sub>]<sup>5–</sup> complexes may be indeed formed in solution.<sup>17</sup>

We have focused our attention on the stoichiometry and stability of species in the solution of the Ln–NP<sub>2</sub>py system by implementing various physical methods such as X-ray crystallography, luminescence spectroscopy, potentiometry as well as a theoretical approach based on Specific Ion Interaction Theory (SIT).<sup>18</sup> Because coordination of aminopolyposphonates to Ln<sup>3+</sup> cations has not been completely clear, we believe that the presented data may be helpful in understanding the interaction of this important class of ligands with lanthanide ions in solution, in general.

## Results and discussion


### Crystal structures

The crystal structures of compounds of the following formula [C(NH<sub>2</sub>)<sub>3</sub>]<sub>5</sub>[Ln(NP<sub>2</sub>py)<sub>2</sub>]·12H<sub>2</sub>O (Ln = Eu or Gd) were determined. The crystals of both compounds are isostructural and belong to the *P*1 space group. The asymmetric unit comprises one [Ln(NP<sub>2</sub>py)<sub>2</sub>]<sup>5–</sup> complex anion, five guanidinium cations out of which one is disordered and twelve water molecules out of which two are disordered. The monomeric [Ln(NP<sub>2</sub>py)<sub>2</sub>]<sup>5–</sup> complex anions are completely deprotonated and their negative charge is compensated by [C(NH<sub>2</sub>)<sub>3</sub>]<sup>+</sup> cations.


Two anionic [NP<sub>2</sub>py]<sup>4–</sup> ligands are bound to the Ln<sup>3+</sup> cation. Each of them is linked with the Ln<sup>3+</sup> by two oxygen atoms from both monodentate phosphonic groups, one tertiary nitrogen atom and one pyridine nitrogen atom, filling thus four coordination sites of the Ln<sup>3+</sup> cation. The molecular structure of the [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5–</sup> complex is presented in Fig. 1. The selected Ln–L bond lengths are presented in Table 1.

As is seen the Eu–L and Gd–L bond lengths are similar within the error of bond length estimation. The Ln–L bond lengths increase in the order Ln–O < Ln–N<sub>(py)</sub> < Ln–N<sub>(am)</sub> which reflects the sequence of ligand donor properties.<sup>19</sup>

The coordination polyhedron of the Ln<sup>3+</sup> cation may be described as a distorted prism (Fig. 2). The Ln<sup>3+</sup> cation lies in




**Scheme 1** Chemical structure of *N*-(methylene-2-pyridine)-*N,N*-di(methylenephosphonic acid) – H<sub>4</sub>NP<sub>2</sub>py.



**Fig. 1** Molecular structure of the [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5–</sup> complex; the H atoms are omitted for clarity.

**Table 1** Selected Ln–L bond lengths in Å

Bond lengths	Eu	Gd
Ln–O21	2.315(2)	2.308(5)
Ln–O13	2.337(2)	2.328(5)
Ln–O15	2.319(2)	2.316(5)
Ln–O24	2.353(2)	2.347(5)
<b>Ln–O(aver.)</b>	<b>2.331(18)</b>	<b>2.325(17)</b>
Ln–N12(am)	2.770(3)	2.764(7)
Ln–N21(am)	2.751(3)	2.744(7)
<b>Ln–N(am aver.)</b>	<b>2.761(13)</b>	<b>2.754(14)</b>
Ln–N11(py)	2.596(3)	2.592(7)
Ln–N22(py)	2.657(3)	2.634(6)
<b>Ln–N(py aver.)</b>	<b>2.626(43)</b>	<b>2.613(29)</b>



**Fig. 2** The coordination polyhedra of Eu<sup>3+</sup> in the [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5–</sup> complex anion.

the general position, but the approximate symmetry of the complex anion is *C*<sub>i</sub>, as shown in Fig. 2.

The [Ln(NP<sub>2</sub>py)<sub>2</sub>]<sup>5–</sup> complexes are held together by a network of hydrogen bonds and electrostatic interactions. Owing to the high negative charge of the anionic [Ln(NP<sub>2</sub>py)<sub>2</sub>]<sup>5–</sup> complex the distance between the neighbouring anions is large and the anionic complexes are well separated from each other; the shortest Eu⋯Eu' and Gd⋯Gd' distances are 10.47(7) Å and 10.37(7), respectively.



### Luminescence spectra of the $[\text{C}(\text{NH}_2)_3]_5[\text{Eu}(\text{NP}_2\text{py})_2]\cdot 12\text{H}_2\text{O}$ crystal

Out of  $\text{Ln}^{3+}$  the  $\text{Eu}^{3+}$  f-f transitions are characterized by a simple and informative spectral pattern that is why we have chosen this cation for detailed spectroscopic studies.

The luminescence spectra of the  $[\text{C}(\text{NH}_2)_3]_5[\text{Eu}(\text{NP}_2\text{py})_2]\cdot 12\text{H}_2\text{O}$  crystal were observed when the  $\text{Eu}^{3+}$  was excited into f-f transitions as well as into ligand states. The luminescence excitation spectrum of the crystals under study is presented in Fig. 3.

Narrow peaks attributed to the f-f transitions of the  $\text{Eu}^{3+}$  and broad band of the ligand ( ${}^1\pi \rightarrow {}^1\pi^*$ ,  ${}^1n \rightarrow {}^1\pi^*$ ) are observed in the spectrum. The presence of the latter band indicates that the energy transfer from the ligand to the  $\text{Eu}^{3+}$  states occurs. The luminescence spectrum of the  $[\text{C}(\text{NH}_2)_3]_5[\text{Gd}(\text{NP}_2\text{py})_2]\cdot 12\text{H}_2\text{O}$  complex indicates that the position of the triplet state of the ligand is centered at  $\sim 437$  nm, Fig. 1S.†<sup>20</sup> The luminescence spectrum of the  $[\text{C}(\text{NH}_2)_3]_5[\text{Eu}(\text{NP}_2\text{py})_2]\cdot 12\text{H}_2\text{O}$  crystal excited into the ligand band is shown in Fig. 4. The emission intensities of the  ${}^5\text{D}_0 \rightarrow {}^7\text{F}_j$  transitions in relation to the  ${}^5\text{D}_0 \rightarrow {}^7\text{F}_1$  one are also included in this figure.

Five bands attributed to the electronic transitions from the excited  ${}^5\text{D}_0$  to the lower lying  ${}^7\text{F}_{0,1,2,3,4}$  states are observed in the spectrum. All these transitions are induced electric dipole (ED) transitions, except for the  ${}^5\text{D}_0 \rightarrow {}^7\text{F}_1$ , which is of magnetic dipole (MD) character. Thus the  ${}^5\text{D}_0 \rightarrow {}^7\text{F}_1$  transition may be used as a reference since its intensity does not depend on the chemical surrounding of  $\text{Eu}^{3+}$ .<sup>21</sup> The  ${}^5\text{D}_0 \rightarrow {}^7\text{F}_0$  and  ${}^5\text{D}_0 \rightarrow {}^7\text{F}_3$  transitions are forbidden due to the selection rules. For this reason, the intensities of these transitions are usually very low as compared with those of  ${}^5\text{D}_0 \rightarrow {}^7\text{F}_{1,2,4}$  transitions. Since the ground  ${}^7\text{F}_0$  and the emitting  ${}^5\text{D}_0$  states are nondegenerate, the  ${}^5\text{D}_0 \leftrightarrow {}^7\text{F}_0$  transitions consist of a single unsplit line for the given  $\text{Eu}^{3+}$  surrounding. Therefore there is a one to one correspondence between the number of lines in the spectrum of this transition and the number of chemically distinct environ-




Fig. 3 Luminescence excitation spectrum ( $\lambda_{\text{em}} = 592.35$  nm) of the  $[\text{C}(\text{NH}_2)_3]_5[\text{Eu}(\text{NP}_2\text{py})_2]\cdot 12\text{H}_2\text{O}$  crystals.




Fig. 4 Luminescence spectrum of the crystal under study ( $\lambda_{\text{ex}} = 276$  nm).

ments of the  $\text{Eu}^{3+}$  ion. The energy levels of the  ${}^7\text{F}_{1,2,3,4}$  multiplets and their crystal field (e) as well as vibronic (v) components found in the RT spectra are presented in Table 2.

Because  ${}^5\text{D}_0 \rightarrow {}^7\text{F}_{0,2,3,4}$  transitions are strictly forbidden in centrosymmetric systems, the presence of these transitions in the luminescence spectrum indicates that the  $C_1$  symmetry of the  $[\text{Eu}(\text{NP}_2\text{py})_2]^{5-}$  complex in the crystal is approximate. The determined luminescence lifetime of the  ${}^5\text{D}_0$  state at RT is 1840  $\mu\text{s}$ .

The next step of our studies was the analysis of the spectroscopic properties of the  $\text{Eu}-\text{NP}_2\text{py}$  complex in aqueous solution.

### Luminescence spectra of the $\text{Eu}-\text{NP}_2\text{py}$ and $\text{Tb}-\text{NP}_2\text{py}$ systems in solution

#### $\text{Ln}-\text{NP}_2\text{py}$ complexes with the metal to ligand ratio 1 : 1.

Taking into account  $\text{Eu}-\text{L}$  bond lengths it may be reasonably assumed that the encapsulation of the  $\text{Ln}^{3+}$  cation by the  $\text{NP}_2\text{py}$  ligand is achieved by the coordination of phosphonic and pyridine groups. The binding of the N(imino) atom to the

Table 2 The energy of the crystal field and vibronic lines of the selected multiplets for the  $4f^6$  configuration at RT.  $\Delta_{\text{CFS}}$  is the energy difference between the highest and lowest CF components of the  ${}^{2S+1}L_J$  multiplet

${}^{2S+1}L_J$	$\text{E}/\text{cm}^{-1}$		${}^{2S+1}L_J$	$\text{E}/\text{cm}^{-1}$	
${}^7\text{F}_1$	180	e	${}^7\text{F}_3$	1800	e
	452	e		1880	e
	512	e		1979	v
	$\Delta_{\text{CFS}}$	332		2131	e
				$\Delta_{\text{CFS}}$	331
${}^7\text{F}_2$	889	e	${}^7\text{F}_4$	2577	e
	919	e		2717	v
	967	v		2901	e
	1050	v		2935	e
	1145	v		3007	v
	1259	v		3064	e
	1287	v		3144	e
	1440	e		3256	e
1585	v	3269	e		
$\Delta_{\text{CFS}}$	550		$\Delta_{\text{CFS}}$	692	




$\text{Ln}^{3+}$  is forced by the former ones. In this context, it seems to be important to indicate how the coordination of phosphonate and pyridine groups depends on the pH of a solution. In order to find the binding mode of the  $\text{NP}_2\text{py}$  ligand and the stoichiometry of the formed  $\text{Ln-NP}_2\text{py}$  species, the luminescence spectra of  $\text{Eu-NP}_2\text{py}$  and  $\text{Tb-NP}_2\text{py}$  complexes in aqueous solution were recorded. The selected luminescence spectra of the  $\text{Eu-NP}_2\text{py}$  complex at different pH are presented in Fig. 5.


At pH 2.26 the intensity of the hypersensitive  $^5\text{D}_0 \rightarrow ^7\text{F}_2$  transition (1.08) is similar to that of the magnetic dipole  $^5\text{D}_0 \rightarrow ^7\text{F}_1$  transition and higher than the relative intensity for the aqua ion (0.68). The increase of pH brings about an increase of the relative intensity of the  $^5\text{D}_0 \rightarrow ^7\text{F}_2$  transition from 1.08 (pH = 2.26) to 2.74 (pH = 10.5). It means that the  $\text{NP}_2\text{py}$  ligand starts to coordinate at pH  $\sim$ 2.26. Under these conditions competing reactions undergo between the protonation and coordination of phosphonate groups to the  $\text{Eu}^{3+}$ .

The protonation of the  $\text{N}_{\text{py}}$  atom at low pH prevents the  $\text{Eu-N}_{\text{py}}$  interaction and consequently may hinder the L-Eu energy transfer. Therefore the question arises whether the efficiency of an antenna effect as a function of pH may serve as a coordination probe for establishing the binding of pyridine to the  $\text{Eu}^{3+}$ . In general the  $\text{Tb}^{3+}$  compounds are more convenient to study the energy transfer process in comparison with those of  $\text{Eu}^{3+}$  for which the ligand band may be imposed on the L-M charge transfer spectra of  $\text{Eu}^{3+}$ .<sup>22</sup> Accordingly, to find the pH range in which the coordination of the pyridyl group to the  $\text{Ln}^{3+}$  occurs, the luminescence spectra of the  $\text{Tb-NP}_2\text{py}$  complex at various pH with an excitation wavelength of 266 nm were recorded (Fig. 6). The oscillator strengths of the  $^1\pi \rightarrow ^1\pi^*$ ,  $^1n \rightarrow ^1\pi^*$  transition calculated from absorption spectra versus pH are also presented in the insert of this figure.

As can be seen the largest increase of the luminescence intensity of all f-f transitions is observed between pH 3 and 5.5. Simultaneously the oscillator strength of the ligand band monotonically decreases. These results indicate that the changes of the efficiency of the antenna effect may be used to study the coordination modes of chromophore groups to the  $\text{Tb}^{3+}$ . The determined values of luminescence quantum yields



**Fig. 5** Luminescence spectra of the  $\text{Eu-NP}_2\text{py}$  complex ( $\lambda_{\text{ex}} = 393$  nm,  $\text{Eu} : \text{NP}_2\text{py} = 1 : 1$ ) in aqueous solution at different pH.




**Fig. 6** Luminescence spectra of the  $\text{Tb-NP}_2\text{py}$  1:1 complex at different pH. The plot of the oscillator strengths of the  $^1\pi \rightarrow ^1\pi^*$ ,  $^1n \rightarrow ^1\pi^*$  transition versus pH is shown in the insert.

$\Phi_{\text{Tb}^{3+}}$  of the  $\text{Tb} : \text{NP}_2\text{py}$  1:1 and 1:2 systems (Fig. 2S<sup>†</sup>) may confirm this conclusion.

It is also worth considering the changes of the energy of the ligand centered transitions, caused by coordination of the  $\text{NP}_2\text{py}$  ligand to the  $\text{Tb}^{3+}$  (Fig. 3S<sup>†</sup>). Because the  $^1\pi \rightarrow ^1\pi^*$ ,  $^1n \rightarrow ^1\pi^*$  transitions are particularly sensitive to the interaction of the N pyridine atom with the metal cation, a remarkable bathochromic shift of the ligand band ( $\sim 550$   $\text{cm}^{-1}$ ) is observed in the spectra of the  $\text{Tb-NP}_2\text{py}$  complex at pH above 6 as compared with the spectra of the free ligand (see Fig. 3S). The observed changes of the f-f and ligand bands above pH  $\sim$  3 support the conclusion that the coordination of the pyridyl group to the  $\text{Tb}^{3+}$  occurs at pH between 3.0 and 5.5. The coordination of the  $\text{NP}_2\text{py}$  ligand is completed above pH = 6.

**Ln-NP<sub>2</sub>py complexes with the metal to ligand ratio 1 : 2.** The luminescence spectra of the  $\text{Eu-NP}_2\text{py}$  system with the  $\text{Eu} : \text{L} = 1 : 2$  in solutions at different pH are presented in Fig. 7.



**Fig. 7** Luminescence spectra of the crystal and solutions of  $\text{Eu}^{3+} : \text{NP}_2\text{py} = 1 : 2$  and 1:1. The bands attributed to the  $[\text{Eu}(\text{NP}_2\text{py})]^-$  species are marked with asterisks.




As can be seen in Fig. 7 the spectral pattern of f-f bands strongly depends on pH. A qualitative comparison between the spectra of the crystal and solutions of the Eu:NP<sub>2</sub>py = 1 : 2 complex demonstrates that at pH = 10.5 [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5-</sup> is a predominant species. As the pH decreases, the CF components of individual multiplets of the [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5-</sup> complex vanish and below pH 8 they completely disappear. It may suggest that the formation of the [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5-</sup> complex starts at pH ~ 8. The presence of additional bands (marked with an asterisk in the spectra in Fig. 7) may indicate that the other species, different to that found in the crystal, also exist in solution.

Owing to the simplicity of the <sup>5</sup>D<sub>0</sub> → <sup>7</sup>F<sub>0</sub> spectrum it was used to monitor the changes in the nearest coordination sphere of the Eu<sup>3+</sup> cation. The luminescence spectra of the <sup>5</sup>D<sub>0</sub> → <sup>7</sup>F<sub>0</sub> transition of solutions and the crystal are presented in Fig. 8.

Two well separated <sup>5</sup>D<sub>0</sub> → <sup>7</sup>F<sub>0</sub> peaks in the spectrum of the solution of the Eu:NP<sub>2</sub>py = 1 : 2 system at pH 10.5 are observed. The comparison of the energy of these peaks with those in the crystal and the solution of Eu : NP<sub>2</sub>py = 1 : 1 at pH = 10.5 enabled us to attribute the higher energy peak at 17330 cm<sup>-1</sup> to the [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5-</sup> species, while the lower energy peak centered at 17286 cm<sup>-1</sup> to the [Eu(NP<sub>2</sub>py)]<sup>-</sup> species. Both <sup>5</sup>D<sub>0</sub> → <sup>7</sup>F<sub>0</sub> peaks are separated by 56 cm<sup>-1</sup>. This difference is probably caused by the change of the number of donor atoms and/or number of water molecules coordinated to the Eu<sup>3+</sup> (see section “Number of coordinated H<sub>2</sub>O molecules to the Eu<sup>3+</sup> cation” in the ESI†).

**Quantitative analysis of the EuL + L ⇌ EuL<sub>2</sub> reaction.** In the next step we focused our attention on a quantitative description of the reaction (1) expressed by the equilibrium constant *K*<sub>12</sub>



$$K_{12} = \frac{[\text{Eu}(\text{NP}_2\text{py})_2^{5-}]}{[\text{Eu}(\text{NP}_2\text{py})^-] \cdot [\text{NP}_2\text{py}^{4-}]} \quad (2)$$




Fig. 8 Luminescence spectra of the <sup>5</sup>D<sub>0</sub> → <sup>7</sup>F<sub>0</sub> transition of the Eu-NP<sub>2</sub>py complexes in the crystal and in solutions at pH = 10.5.

To determine the *K*<sub>12</sub> constant we used the method based on the fitting of the spectra of the crystals to those of solutions.<sup>23</sup> The spectrum of the crystal under study served us as a model of 1 : 2 species while the spectrum of Eu-NP<sub>2</sub>py 1 : 1 at pH = 10.5 was used as the spectrum of a pure 1 : 1 complex. However owing to the dynamic effects,<sup>23,24</sup> the spectra of f-f transitions in the solution are broadened in comparison to the spectra of the crystal. For this reason the simulated spectra of the [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5-</sup> species derived from the spectra of the crystal were artificially broadened. The equation that was used to calculate the simulated spectrum of the [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5-</sup> species in solution is as follows:

$$J_{\text{calc}} = \left( \sum_{i=1}^z A_{12} J_{\text{max}} \frac{\exp \left[ -0.3 \cdot \frac{\lambda - \lambda_{\text{max}}}{B_{12} \cdot \Delta \lambda} \right]}{\left[ 1 + \left( \frac{\lambda - \lambda_{\text{max}}}{B_{12} \cdot \Delta \lambda} \right)^2 \right]} \right) \quad (3)$$

where:

*J*<sub>calc</sub> – intensity at a given value of λ;

*z* – the number of peaks for the given <sup>5</sup>D<sub>0</sub> → <sup>7</sup>F<sub>*j*</sub> transition;

*J*<sub>max</sub> – the maximum intensity of the peak;

λ<sub>max</sub> – the wavelength of the peak maximum;

Δλ – the half-width of a peak;

*A*<sub>12</sub> and *B*<sub>12</sub> are adjustable parameters connected with artificial broadening of bands.

The final set of parameters used to calculate the simulated spectrum of the [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5-</sup> species is presented in Table 1S (ESI†). The simulated and experimental spectra of compounds under study are shown in Fig. 9.

Next, the simulated spectra of the <sup>5</sup>D<sub>0</sub> → <sup>7</sup>F<sub>1,2,4</sub> transitions were used to calculate the molar fractions χ of [Eu(NP<sub>2</sub>py)]<sup>-</sup> and [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5-</sup> species (Table 3).

The average molar fractions of [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5-</sup> and [Eu(NP<sub>2</sub>py)]<sup>-</sup> species are ~81% and ~19%, respectively. These results are consistent, within the error of experimental limits (±10%), regardless of the transition taken for the calculation.




Fig. 9 Simulated spectra of the [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5-</sup> (red) and [Eu(NP<sub>2</sub>py)]<sup>-</sup> (blue) species. The fitted and experimental spectra of the Eu : NP<sub>2</sub>py 1 : 2 system at pH 10.5.



**Table 3** Molar fractions  $\chi$  of  $[\text{Eu}(\text{NP}_2\text{py})]^-$  and  $[\text{Eu}(\text{NP}_2\text{py})_2]^{5-}$  species in aqueous solution at pH = 10.5 determined from various  $^5\text{D}_0 \rightarrow ^7\text{F}_{1,2,4}$  transitions

Transition	$\chi$ $[\text{Eu}(\text{NP}_2\text{py})]^-$	$\chi$ $[\text{Eu}(\text{NP}_2\text{py})_2]^{5-}$	$\log K_{12}$
$^5\text{D}_0 \rightarrow ^7\text{F}_1$	25% $\pm$ 10%	75% $\pm$ 10%	4.2 $\pm$ 0.5
$^5\text{D}_0 \rightarrow ^7\text{F}_2$	18% $\pm$ 10%	82% $\pm$ 10%	4.5 $\pm$ 0.5
$^5\text{D}_0 \rightarrow ^7\text{F}_4$	15% $\pm$ 10%	85% $\pm$ 10%	4.6 $\pm$ 0.5
Average	19(5)	81(5)	4.4 $\pm$ 0.5

The estimated averaged value of  $\log K_{1-2}$  for the reaction (1) is about 4.4  $\pm$  0.5.

It should be noted that the presented speciation analysis for the case of f-elements is not always feasible, because the complexes existing in equilibrium are usually characterized by similar spectral patterns as it was shown previously by Horrocks *et al.* for the case of lanthanide complexes with NTP.<sup>15</sup>

To support the presented results the analysis was extended by potentiometric measurements.

### Potentiometric study

**Protonation of the ligand.** We performed potentiometric titrations of a free ligand in order to determine its acid-base properties. The fully protonated form of the  $\text{H}_4\text{NP}_2\text{py}$  possesses six dissociable protons  $[\text{H}_6\text{L}]^{2+}$  (see Fig. 6S†) two at each phosphonic group, one at the pyridine moiety and one on the central tertiary N amine atom. However, it was possible to determine only four protonation constants ( $K_1$ – $K_4$ ) because the  $K_5$  and  $K_6$  values corresponding to the dissociation of one proton from each of the  $\text{PO}_3\text{H}_2$  groups can be determined only for solutions at pH below 2. This is beyond the available range for standard potentiometric titrations, and the  $\log K_5$  and  $\log K_6$  values were determined previously by the NMR method.<sup>17</sup> Measurements in different ionic strengths (KCl): 0.075 M, 0.1 M, 0.2 M, 0.5 M, 1 M and 2 M were also performed. The calculated  $\log K_n$  values are listed in Table 4.

The most basic constant  $\log K_1 = 10.63(2)$  (Table 4) corresponds to the protonation of the nitrogen of the tertiary amino group and is two logarithmic units higher than that of  $\text{H}_6\text{NTP}$  acid.<sup>14,25–27</sup> The values of this constant for NTP, determined by potentiometric methods in 0.1 M ionic strengths in different electrolytes at 25 °C were found to be: 12.7(1),<sup>14</sup> 12.8(2),<sup>25</sup> 12.20(6),<sup>26</sup> 12.5.<sup>27</sup> Such a decrease of the highest protonation

**Table 4** The protonation constants of  $\text{H}_4\text{NP}_2\text{py}$  in 25 °C in different ionic strengths and background electrolytes

$I$ [M]		$\log K_1$ (HL)	$\log K_2$ (H <sub>2</sub> L)	$\log K_3$ (H <sub>3</sub> L)	$\log K_4$ (H <sub>4</sub> L)
0.1	NaClO <sub>4</sub>	10.63(2)	6.74(6)	5.61(8)	2.23(9)
0.075	KCl	10.87(1)	6.55(2)	5.47(2)	2.17(2)
0.1	KCl	10.83(1)	6.61(2)	5.29(2)	1.60(6)
0.2	KCl	10.78(1)	6.41(2)	5.37(2)	2.00(3)
0.5	KCl	10.59(1)	6.27(2)	5.25(2)	1.60(6)
1	KCl	10.50(1)	6.17(2)	5.26(2)	2.01(4)
2	KCl	10.51(1)	6.19(2)	5.27(2)	1.60(8)

constant of  $\text{H}_4\text{NP}_2\text{py}$  is clearly caused by substitution of one of the phosphonic groups by the pyridine moiety. The same phenomenon has been noticed by Kurzak *et al.*<sup>28</sup> in *N*-3-picolyliminodi(methylenephosphonic) acid ( $\text{NP}_3\text{py}$ ); the  $\log K$  value of the tertiary amine was found to be 10.40(1) in 0.2 M KCl. The following deprotonation steps of  $\text{H}_4\text{NP}_2\text{py}$  release the protons from both phosphonic groups;  $\log K_2 = 6.74(4)$ ,  $\log K_4 = 2.23(5)$  and the pyridine moiety;  $\log K_3 = 5.61(4)$  (Table 4). Such deprotonation of  $\text{H}_4\text{NP}_2\text{py}$  is consistent with  $^{31}\text{P}$  NMR results<sup>17</sup> and based on those data it was concluded that the third and fourth dissociation steps occur at pH between 2 and 3 ( $\log K_4$ ) and 4–7 ( $\log K_3$ ,  $\log K_2$ ) and deprotonation of phosphonic groups is completed at pH above 8.2. The observed changes for the chemical shifts of pyridyl protons suggested that deprotonation of pyridine nitrogen occurs between pH 5 and 6, thus the  $\text{p}K_3$  value was assigned to pyridine nitrogen. This conclusion is confirmed by the observed changes in the UV-vis spectra of the  $\text{NP}_2\text{py}$  ligand. The spectrophotometric titration curve is shown in Fig. 10.

As was mentioned above, the  $^1\pi \rightarrow ^1\pi^*$  and  $^1n \rightarrow ^1\pi^*$  band observed in the UV spectrum of the  $\text{NP}_2\text{py}$  ligand is particularly sensitive to the protonation of pyridine *N* (Fig. 3S†). A remarkable decrease of the  $\epsilon$  value is observed between pH 5 and 7. This unequivocally confirms the attribution of  $\log K_3$  to the protonation of the pyridine moiety.

The minimalization of the sum equal to  $\sum (\epsilon_{\text{exp}} - \epsilon_{\text{calc}})^2$  by using eqn (4) enabled us to determine the  $\beta_n$  values which are converted into  $K_n$  values (Table 5).


$$\epsilon_{\text{calc}} = \frac{\epsilon_{\text{L}} + [\text{H}]\beta_1\epsilon_{\text{HL}} + [\text{H}]^2\beta_2\epsilon_{\text{H}_2\text{L}} + [\text{H}]^3\beta_3\epsilon_{\text{H}_3\text{L}} + [\text{H}]^4\beta_4\epsilon_{\text{H}_4\text{L}}}{1 + [\text{H}]\beta_1 + [\text{H}]^2\beta_2 + [\text{H}]^3\beta_3 + [\text{H}]^4\beta_4} \quad (4)$$

where:

$\epsilon_{\text{exp}}$ ,  $\epsilon_{\text{calc}}$  – experimental and calculated absorption molar coefficients ( $\text{M}^{-1} \text{cm}^{-1}$ ) at  $\lambda = 259.6 \text{ nm}$ ;

$\epsilon_{\text{L}} - \epsilon_{\text{H}_n\text{L}}$  – absorption molar coefficients of individual species;

$\beta_n$  – cumulative protonation constant.



**Fig. 10** Spectrophotometric titration curve of the  $\text{NP}_2\text{py}$  ligand.



**Table 5** The comparison of the protonation constants of the NP<sub>2</sub>py ligand using different methods

	<sup>31</sup> P NMR <sup>17</sup>	UV-vis	Potentiometry
log <i>K</i> (HL)	10.9(9)	11.0(9)	10.63(2)
log <i>K</i> <sub>2</sub> (H <sub>2</sub> L)	6.7(9)	6.6(9)	6.74(6)
log <i>K</i> <sub>3</sub> (H <sub>3</sub> L)	5.5(9)	5.6(9)	5.61(8)
log <i>K</i> <sub>4</sub> (H <sub>4</sub> L)	2.3(9)	2.7(9)	2.23(9)
log <i>K</i> <sub>5</sub> (H <sub>5</sub> L)	0.98(5)	—	—
log <i>K</i> <sub>6</sub> (H <sub>6</sub> L)	0.70(5)	—	—

As is seen from Table 5 the values of the respective log *K<sub>n</sub>* determined from different methods are the same, within the experimental error. Finally the calculated values of log *K<sub>n</sub>* were used to determine the speciation distribution curve, which is presented in Fig. 6S.†

**Formation constants.** Potentiometric titrations of the Ln–NP<sub>2</sub>py system (where Ln = Sm<sup>3+</sup>, Eu<sup>3+</sup>, Gd<sup>3+</sup> and Tb<sup>3+</sup>) were performed in order to determine the stoichiometry and stability constants of the formed complexes. The potentiometric measurements were performed for two Ln : NP<sub>2</sub>py molar ratios 1 : 1 and 1 : 2. The calculated stability constants of LnL = [Ln(NP<sub>2</sub>py)]<sup>−</sup> and LnL<sub>2</sub> = [Ln(NP<sub>2</sub>py)<sub>2</sub>]<sup>5−</sup> complexes (at 25 °C, *I*<sub>NaClO<sub>4</sub></sub> = 0.1 M) are collected in Table 3S† and selected values are shown in Table 6.

The species distribution curves are depicted in Fig. 7S.† The potentiometric measurements of Eu : NP<sub>2</sub>py = 1 : 1 and 1 : 2 systems at different ionic strengths (KCl) were also performed and the determined stability constants are given in Table 4S.†


Based on potentiometric data it was possible to calculate log *K*<sub>12</sub> for reaction 2, which is equal to 5.27 and similar within the limit of experimental error to that estimated from the spectroscopic data (4.4 ± 0.5).

The determined stability constants for Eu–NP<sub>2</sub>py systems were used to calculate the average ligand number ( $\bar{n}$ ) as a function of pH (Fig. 11). The data for the Eu–NTA (1 : 2) complex (where NTA is nitrilotriacetic acid) are also included in this figure for comparison purposes.<sup>29a</sup>

As is seen in Fig. 11 the coordination of the NP<sub>2</sub>py ligand to the Eu<sup>3+</sup> starts at about pH 3 for 1 : 1 and 1 : 2 complexes. In the case of the Eu : NP<sub>2</sub>py 1 : 2 system the formation of the [Ln(NP<sub>2</sub>py)<sub>2</sub>]<sup>5−</sup> complex starts at about pH 8 and it is consistent with the spectroscopic data described above. The difference between Eu–NTA and Eu–NP<sub>2</sub>py complexes with the molar

**Table 6** The stability constants of Ln<sup>3+</sup> complexes with the NP<sub>2</sub>py ligand for Ln : NP<sub>2</sub>py molar ratios of 1 : 1 and 1 : 2, at 25 °C, *I* = 0.1 mol dm<sup>−1</sup> (NaClO<sub>4</sub>)



Ln : NP <sub>2</sub> py	1 : 1 log β <sub>LnL</sub>	1 : 2 log β <sub>LnL<sub>2</sub></sub>	log β <sub>LnL<sub>2</sub></sub>
Sm <sup>3+</sup>	11.87(6)	11.83(5)	16.99(8)
Eu <sup>3+</sup>	12.85(8)	12.85(6)	18.44(9)
Gd <sup>3+</sup>	12.32(10)	12.45(6)	17.88(7)
Tb <sup>3+</sup>	13.36(7)	13.47(10)	17.86(9)

**Fig. 11** The average ligand number  $\bar{n}$  as a function of pH for Eu–NTA (1 : 2 ratio) as well as for Eu–NP<sub>2</sub>py complexes (1 : 2 and 1 : 1 ratios).

ratio 1 : 2 is clearly seen. The coordination of two NTA ligands occurs at low pH ~ 2<sup>29</sup> and is completed at pH ~ 6. As far as Eu–NP<sub>2</sub>py of the 1 : 2 system is concerned, the [Eu(NP<sub>2</sub>py)]<sup>−</sup> complex is a predominating species within the pH range 3–9 and the formation of the [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5−</sup> complex takes place at pH above 8. This may be explained qualitatively by considering the formal charge of reacting species. In the case of the Ln–NTA system the neutral LnNTA complex reacts with the H<sub>n</sub>NTA ligand to form finally [Ln(NTA)<sub>2</sub>]<sup>3−</sup> species with the formal charge −3. The formal charge of the Ln(NP<sub>2</sub>py)<sub>2</sub> complex is −5 and the formed complex is not as stable as [Ln(NTA)<sub>2</sub>]<sup>3−</sup>. Out of the considered complexes, the [Ln(NTP)<sub>2</sub>]<sup>9−</sup> one is the most negatively charged, therefore this may suggest that the stability of the complexes should change in the following order: Ln(NTA)<sub>2</sub> > Ln(NP<sub>2</sub>py)<sub>2</sub> > Ln(NTP)<sub>2</sub>. Such an assumption is consistent with potentiometric results derived from Sawada *et al.*<sup>14</sup> These authors have shown that only the [Ln(NTP)]<sup>3−</sup> complex is formed in solution, on the other hand Kozlovski *et al.*<sup>16</sup> assumed the existence of both [Ln(NTP)]<sup>3−</sup> and [Ln(NTP)<sub>2</sub>]<sup>9−</sup> species.

#### Extrapolation of [Eu(NP<sub>2</sub>py)]<sup>−</sup> and [Eu(NP<sub>2</sub>py)<sub>2</sub>]<sup>5−</sup> stability constants to zero ionic strength

The SIT approach (Specific Ion Interaction Theory)<sup>18</sup> was used for the extrapolation and correction of equilibrium data to the infinite dilution standard state. This simple method provides good estimations of activity coefficients and is recommended for use by the OECD Nuclear Energy Agency.<sup>30</sup> The extrapolation of the formation constants to ionic strength equal to zero requires the estimation of activity coefficients of all species that participated in chemical reactions. This may be performed by the SIT model. For the reactions:



According to the SIT procedure, the logarithms of stability constants at a fixed ionic strength  $\log \beta_n(I)$  are related to the corresponding logarithms of thermodynamic constants  $\beta_n^0$  by the following equations:

$$\log \beta_{\text{H}_n\text{L}}^0 = \log \beta_{\text{H}_n\text{L}}(I_m) + \log \gamma_{\text{H}_n\text{L}} - \log \gamma_{\text{L}} - n \log \gamma_{\text{H}} \quad (7)$$

$$\log \beta_{\text{EuL}_m}^0 = \log \beta_{\text{EuL}_m}(I_m) + \log \gamma_{\text{EuL}_m} - \log \gamma_{\text{Eu}} - m \log \gamma_{\text{L}} \quad (8)$$

where  $\gamma$  are activity coefficients of individual species.

The  $\log \gamma_j$  of the  $j$ -species may be calculated by using the following formula:

$$\log \gamma_j = -z_j^2 \frac{0.509\sqrt{I_m}}{1 + 1.5\sqrt{I_m}} + \varepsilon_j I_m \quad (9)$$

where:

$z$  - charge of the  $j$ -species;

$I_m$  - ionic strength of solution expressed in mol kg<sup>-1</sup>;

$\varepsilon_j$  - the interaction coefficient of ion pairs.

Combining eqn (8) or (9) with (10), the following equations are derived:

$$\log \beta_{\text{H}_n\text{L}}(I_m) + \Delta z^2 \frac{0.509\sqrt{I_m}}{1 + 1.5\sqrt{I_m}} = \log \beta_{\text{H}_n\text{L}}^0 - \Delta \varepsilon_{\text{H}_n\text{L}} I_m \quad (10)$$

$$\log \beta_{\text{EuL}_m}(I_m) + \Delta z^2 \frac{0.509\sqrt{I_m}}{1 + 1.5\sqrt{I_m}} = \log_{10} \beta_{\text{EuL}_m}^0 - \Delta \varepsilon_{\text{EuL}_m} I_m \quad (11)$$

where:

$\Delta z^2 = \sum z_{\text{products}}^2 - \sum z_{\text{substrates}}^2$ ;

$\Delta \varepsilon^2 = \sum \varepsilon_{\text{products}}^2 - \sum \varepsilon_{\text{substrates}}^2$ .

Linear regressions of eqn (10) and (11), using the experimental values of  $\log \beta(I_m)$  and  $I_m$  give the thermodynamic values of  $\log \beta_{\text{H}_n\text{L}}^0$  and  $\log \beta_{\text{EuL}_m}^0$  (Table 7).

In general, these data may be useful in finding a reliable, thermodynamic model of the complexation reaction of aminophosphonic ligands with lanthanides particularly that, to the best of our knowledge, there are no stability constants of lanthanide/actinide phosphonates extrapolated to the zero ionic strength. This is the reason for which we are unable to refer our data to other ones.

**Table 7** Formation constants  $\log \beta_{\text{AB}}^0$  of compounds under study at the zero ionic strength at 25 °C

AB	$\log \beta_{\text{AB}}^0$	$\Delta \varepsilon$	$R^2$
HL	11.8 ± 0.05	-0.31 ± 0.05	0.92
H <sub>2</sub> L	19.0 ± 0.05	-0.48 ± 0.04	0.98
H <sub>3</sub> L	24.6 ± 0.11	-0.88 ± 0.10	0.96
H <sub>4</sub> L	26.9 ± 0.26	-0.76 ± 0.25	0.76
EuL	16.3 ± 0.11	-0.82 ± 0.11	0.95
EuL <sub>2</sub>	19.5 ± 0.15	-0.46 ± 0.15	0.77

## Summary and conclusions

The compounds of the formula  $[\text{C}(\text{NH}_2)_3]_5[\text{Ln}(\text{NP}_2\text{py})_2] \cdot 12\text{H}_2\text{O}$  (where Ln = Eu, Gd) with new tripodal NP<sub>2</sub>py ligands were synthesized and their crystal structure was determined. The crystal comprises monomeric  $[\text{Eu}(\text{NP}_2\text{py})_2]^{5-}$  anions. Both ligands are bonded with the Eu<sup>3+</sup> cation by two oxygen atoms from monodentate phosphonic groups, one tertiary nitrogen atom and one pyridine nitrogen atom, filling thus the eight coordination sites of the Eu<sup>3+</sup> cation. The approximate C<sub>i</sub> symmetry of the  $[\text{Eu}(\text{NP}_2\text{py})_2]^{5-}$  complex is reflected in the luminescence spectra, in which the intensity of the magnetic dipole <sup>5</sup>D<sub>0</sub> → <sup>7</sup>F<sub>1</sub> transition dominates over the intensities of the remaining <sup>5</sup>D<sub>0</sub> → <sup>7</sup>F<sub>0,2,3,4</sub> transitions.

The luminescence spectra of Eu-NP<sub>2</sub>py and Tb-NP<sub>2</sub>py complexes in aqueous solutions at different pH with various Ln : L ratios were studied in order to find the coordination mode of the NP<sub>2</sub>py ligand and the stoichiometry of complexes. The NP<sub>2</sub>py ligand starts to coordinate through phosphonic oxygen atoms with Ln<sup>3+</sup> at pH ~ 2. The coordination of the pyridyl group to the Tb<sup>3+</sup> occurs at pH between 3.0 and 5.5. The binding of the NP<sub>2</sub>py ligand is completed above pH = 6.

This study has demonstrated that both  $[\text{Ln}(\text{NP}_2\text{py})]^-$  and  $[\text{Ln}(\text{NP}_2\text{py})_2]^{5-}$  complexes exist in solution, however, the  $[\text{Ln}(\text{NP}_2\text{py})_2]^{5-}$  starts to form at pH as high as 8. The comparison of the luminescence spectra of the  $[\text{C}(\text{NH}_2)_3]_5[\text{Eu}(\text{NP}_2\text{py})_2] \cdot 12\text{H}_2\text{O}$  crystals with those of Eu : NP<sub>2</sub>py (1 : 2) solutions enabled us to estimate the molar fraction of  $[\text{Eu}(\text{NP}_2\text{py})]^-$  (~19%) and  $[\text{Eu}(\text{NP}_2\text{py})_2]^{5-}$  (~81%) complexes at pH = 10.5, and finally, to calculate the conditional formation constant of reaction  $[\text{Eu}(\text{NP}_2\text{py})]^- + \text{NP}_2\text{py}^{4-} \rightleftharpoons [\text{Eu}(\text{NP}_2\text{py})_2]^{5-}$  ( $\log K_{12} = 4.4 \pm 0.5$ ).

The potentiometric measurements were applied to determine the stability constants of complexes under study. The stability constants are between 11.83(5) for  $[\text{Sm}(\text{NP}_2\text{py})]^-$  and 13.47(7) for  $[\text{Tb}(\text{NP}_2\text{py})]^-$  and are slightly lower than those obtained for Ln-NTP complexes.<sup>16</sup> Thus, the substitution of one phosphonic group by a pyridyl moiety slightly affects the stability of the  $[\text{Ln}(\text{NP}_2\text{py})]^-$  complex in comparison with the  $[\text{Ln}(\text{NTP})]^{3-}$  one.<sup>16</sup> On the other hand the  $\log \beta_{\text{LnL}_2}^0$  of the  $[\text{Ln}(\text{NP}_2\text{py})_2]^{5-}$  complexes are about 5–6 orders smaller in comparison with those reported for  $[\text{Ln}(\text{NTP})_2]^{9-}$ .<sup>16</sup> It should be noted, however, that the existence of  $[\text{Ln}(\text{NTP})_2]$  complexes in solution has not been proved unequivocally.<sup>14–16</sup>

Potentiometric titrations at various ionic strengths were also performed in order to find the thermodynamic stability constants of  $[\text{Eu}(\text{NP}_2\text{py})]^-$  and  $[\text{Eu}(\text{NP}_2\text{py})_2]^{5-}$  complexes. Such data are, in principle, preferable as a basis for discussion on the factors governing the stabilities of the complexes. The extrapolation of the formation constants to zero ionic strength under standard conditions was performed by the SIT procedure. The estimated  $\log \beta_{\text{EuL}}^0$  and  $\log \beta_{\text{EuL}_2}^0$  values are 16.3 ± 0.11 and 19.5 ± 0.15, respectively. For lanthanide aminophosphonate complexes there are no thermodynamic quantities in the literature therefore we refer the obtained data to the well-known  $[\text{Eu}(\text{NTA})]$  and  $[\text{Eu}(\text{NTA})_2]^{3-}$  complexes, only.<sup>29</sup> The



latter are characterized by the thermodynamic stability constants:  $\log \beta_{\text{EuNTA}}^0 = 13.34$  and  $\log \beta_{\text{Eu(NTA)}_2}^0 = 22.66$ .

Taking into account all these data one may conclude that the high negative charge of a ligand may stabilize the LnL complex, but simultaneously destabilizes the LnL<sub>2</sub> one.

## Experimental

### Materials

All chemicals used were of analytical grade. The stock solutions of anhydrous Ln<sup>3+</sup> chlorides (Aldrich) were standardized against EDTA using xylenol orange as an indicator. *N*-(methylene-2-pyridine)-*N,N*-di(methylenephosphonate) as a diammonium dihydrogen compound was synthesized as described previously.<sup>17</sup>

### Crystal preparation

The crystals of the formula [C(NH<sub>2</sub>)<sub>3</sub>]<sub>5</sub>[Ln(NP<sub>2</sub>py)<sub>2</sub>]·12H<sub>2</sub>O (Ln = Eu or Gd) were obtained in the following way:

The suspension of 0.5 mmol Ln<sub>2</sub>O<sub>3</sub> (Eu<sub>2</sub>O<sub>3</sub> Stanford Materials 99.995%, Gd<sub>2</sub>O<sub>3</sub> Stanford Materials 99.99%) and of 2.2 mmol (NH<sub>4</sub>)<sub>2</sub>H<sub>2</sub>NP<sub>2</sub>py salt in H<sub>2</sub>O were heated at 90 °C ± 5 °C. After dissolution of the reagents, the solution was alkalinized by using [C(NH<sub>2</sub>)<sub>3</sub>]<sub>2</sub>CO<sub>3</sub> to the final pH ~ 10.5 and left for crystallization. The colourless crystals were formed after three months.

The luminescence quantum yields for solutions of the Tb<sup>3+</sup>-NP<sub>2</sub>py complexes were determined in relation to fluorescein in 0.1 M NaOH ( $\Phi = 93\%$ ), by using eqn (13):

$$Q_X = Q_S \cdot \frac{n_X^2 \int_{\bar{\nu}_1}^{\bar{\nu}_2} I_X(\bar{\nu}) d\bar{\nu}}{n_S^2 \int_{\bar{\nu}_1}^{\bar{\nu}_2} A_S(\bar{\nu}) d\bar{\nu}} \cdot \frac{1 - 10^{-\epsilon_S \cdot d \cdot c_S}}{1 - 10^{-\epsilon_X \cdot d \cdot c_X}} \quad (13)$$

where  $Q$  stands for the quantum yield,  $n$  is the refraction index ( $n_X = 1.38$ ,  $n_S = 1.33$ ), the integrals were taken for the emission spectra corrected for the instrument response,  $\epsilon$  is the molar absorption coefficient,  $c$  – concentration, and  $d$  – thickness of the cuvette (1 cm). The subscripts X and S stand for the sample and the standard, respectively. The pairs of the sample and the standard spectra used in the above equation were taken at the same instrument setting (slit widths, photomultiplier voltage *etc.*). The slits were 1 nm for the excitation and 1 nm for the emission recording.

**Elemental analysis.** C<sub>21</sub>H<sub>74</sub>EuN<sub>19</sub>O<sub>24</sub>P<sub>4</sub>: N%<sub>calc</sub> – 21.24%, N%<sub>exp</sub> – 20.85%, C%<sub>calc</sub> – 20.13%, C%<sub>exp</sub> – 20.56%, H%<sub>calc</sub> – 5.95%, H%<sub>exp</sub> – 5.28%; C<sub>21</sub>H<sub>74</sub>GdN<sub>19</sub>O<sub>24</sub>P<sub>4</sub>: N%<sub>calc</sub> – 21.15%, N%<sub>exp</sub> – 22.48%, C%<sub>calc</sub> – 20.05%, C%<sub>exp</sub> – 21.56%, H%<sub>calc</sub> – 5.93%, H%<sub>exp</sub> – 5.30%.

### X-Ray crystal analysis

The crystallographic measurements were performed on an Xcalibur, Sapphire2 four-circle diffractometer with graphite-monochromatized MoK $\alpha$  radiation. The data for the crystal were collected at 100(2) K using the Oxford Cryosystems cooler and were analytically corrected for absorption with the use of the CrysAlis RED program of the Xcalibur software. The structures were solved routinely by using Patterson synthesis. The C- and N-bonded hydrogen atoms were placed in positions calculated from the geometry. The final refinement was anisotropic for all non-H atoms. The computations were performed with the SHELXS97<sup>31</sup> and SHELXL97 programs,<sup>32</sup> and the molecular graphics was prepared with XP-Interactive Molecular Graphics.<sup>33</sup> High residual peaks suggest that the C50 guanidine cation is partially distorted, however it was not possible to resolve this distortion. A summary of the conditions for the data collection and the structure refinement parameters is given in Table 8.

**Table 8** Crystal data

CCDC no.	CCDC 1541895	CCDC 1541894
Chemical formula	C <sub>21</sub> H <sub>74</sub> EuN <sub>19</sub> O <sub>24</sub> P <sub>4</sub>	C <sub>21</sub> H <sub>74</sub> GdN <sub>19</sub> O <sub>24</sub> P <sub>4</sub>
$M_r$	1252.83	1258.12
Crystal system, space group	Triclinic, $P\bar{1}$	Triclinic, $P\bar{1}$
Temperature (K)	100(2)	100(2)
$a, b, c$ (Å)	11.9255 (11), 14.1373 (12), 16.1417 (15)	11.9496 (6), 14.1528 (7), 16.1651 (9)
$\alpha, \beta, \gamma$ (°)	105.889 (8), 94.515 (7), 90.629 (7)	105.710 (5), 94.360 (4), 90.901 (4)
$V$ (Å <sup>3</sup> )	2607.8 (4)	2622.2 (2)
$Z$	2	2
$\mu$ (mm <sup>-1</sup> )	1.39	1.48
Crystal size (mm)	0.45 × 0.07 × 0.04	0.27 × 0.05 × 0.02
$T_{\text{min}}, T_{\text{max}}$	0.709, 0.958	0.816, 0.979
No. of measured, independent and observed [ $I > 2\sigma(I)$ ] reflections	51 538, 25 885, 19 822	23 976, 15 463, 8529
$R_{\text{int}}$	0.053	0.093
$(\sin \theta/\lambda)_{\text{max}}$ (Å <sup>-1</sup> )	0.851	0.737
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.061, 0.147, 1.15	0.101, 0.184, 1.05
No. of reflections	25 885	15 463
No. of parameters	601	601
No. of restraints	6	6
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å <sup>-3</sup> )	4.71, -2.10	1.26, -1.25



## Spectroscopic measurements

Luminescence spectra and luminescence lifetimes were recorded on an Edinburgh Instruments FLS 920 spectrometer at RT. UV-vis absorption spectra were recorded on a Cary 500 UV/Vis/NIR spectrophotometer.

## Potentiometric titrations

The solution studies were carried out in double-distilled water. The titrations were performed using a MOLSPIN pH-meter system equipped with a Mettler Toledo InLab® Micro electrode filled with 3 M KCl and the automatic burette. The electrode was calibrated daily against hydrogen concentrations using HClO<sub>4</sub> (Merck) or HCl (POCh) according to the procedure of Irving *et al.*<sup>34</sup> The purity and exact concentration of the ligand were determined by the method of Gran.<sup>35</sup> Alkali, carbonate-free, was standardized by titration with potassium hydrogen phthalate (Merck).

Measurements in 0.1 M ionic strength:

The ionic strength was fixed at  $I = 0.1$  M with NaClO<sub>4</sub> (VWR). The ionic product of water under the used conditions was  $10^{-13.77}$  mol<sup>2</sup> dm<sup>-6</sup>. Alkali: 0.1 M sodium hydroxide solution (Aldrich).

Measurements in various ionic strengths:

The ionic strength was fixed at  $I = 0.075$  M, 0.1 M, 0.2 M, 0.5 M, 1 M and 2 M with KCl (POCh). The ionic product of water under the used conditions was  $10^{-13.78}$ ,  $10^{-13.77}$ ,  $10^{-13.72}$ , and  $10^{-13.75}$  mol<sup>2</sup> dm<sup>-6</sup>, respectively.<sup>36</sup> Alkali: 0.1 M potassium hydroxide solution (Aldrich).

All the titrations were carried out on 2.0–3.0 ml samples at  $25 \pm 0.2$  °C under an atmosphere of Ar. SUPERQUAD<sup>37</sup> and HYPERQUAD2008<sup>38</sup> computer programs that use non-linear least-squares methods<sup>39</sup> were applied to calculate the stability constants. The results were obtained in the form of concentration overall stability constants  $\beta_{pqr} = [M_p H_q L_r] / [M]^p [H]^q [L]^r$ , where M stands for metal, H is proton and L the deprotonated form of the ligand. They are, however, a good indication of the importance of a particular species in the equilibrium. Triplicate titrations of the free ligand (number of recorded points: 150–180) and the complexes were carried out at metal to ligand ratios 1:1 and 1:2 (number of recorded points: 180–210). Each time about 120–150 and 150–190 titration points have been used for the calculations, respectively. The ligand concentration was  $0.7\text{--}1 \times 10^{-3}$  mol dm<sup>-3</sup> in all titrations. No precipitation was noticed in any titration. The distribution curve of the protonated species of L as a function of pH was calculated using the HySS2009 program.<sup>40</sup>

The ligand used for potentiometric measurements was in the form of a diammonium salt ((NH<sub>4</sub>)<sub>2</sub>H<sub>2</sub>NP<sub>2</sub>py); NP<sub>2</sub>py = [H<sub>6</sub>L]<sup>2+</sup> in order to improve its solubility in water. Therefore two ammonium cations were also considered in the overall acid–base equilibrium calculations of the compound. The determined values of the protonation constants which belong to the ammonia log  $K_{\text{NH}_4}$  vary within the range 8.98(1)–9.73(1) for different studied ionic strengths and are very close to those

previously reported in the literature: log  $K = 9.38(1)$ ,<sup>41</sup> log  $K = 9.50$ ,<sup>42</sup> log  $K = 9.10(3)$ ,<sup>43</sup> ( $I = 0.1$  M).

**CAUTION!** Although no problems were encountered in this work, transition metal perchlorate complexes are potentially explosive and should be handled with appropriate precautions.

## Acknowledgements

This work was supported by MNiSW grant no 2273/M/WCH/12.

## References

- (a) W. A. Volkert and T. J. Hoffmann, *Chem. Rev.*, 1999, **99**, 2269–2292; (b) E. Gumienna-Kontecka and J. Gałęzowska, *Coord. Chem. Rev.*, 2012, **256**, 105–124; (c) J. Goura and V. Chandrasekhar, *Chem. Rev.*, 2015, **115**, 6854–6965; (d) J. A. Amoroso, I. A. Fallis and S. J. A. Pope, *Coord. Chem. Rev.*, 2017, **340**, 198–219; (e) A. D. G. Firmino, R. F. Mendes, M. M. Antunes, P. C. Barbosa, S. M. F. Vilela, A. A. Valente, F. M. L. Figueiredo, J. P. C. Tome and F. A. A. Paz, *Inorg. Chem.*, 2017, **56**, 1193–1208.
- L. Lukeš, J. Kotek, P. Vojtisek and P. Hermann, *Coord. Chem. Rev.*, 2001, **216–217**, 287.
- (a) R. Janicki and P. Starynowicz, *Acta Crystallogr., Sect. B: Struct. Sci.*, 2010, **66**, 559–567; (b) R. Janicki, A. Kędziorki and A. Mondry, *Phys. Chem. Chem. Phys.*, 2016, **18**, 27808–27817.
- A. Mermer and P. Starynowicz, *Acta Crystallogr., Sect. B: Struct. Sci.*, 2011, **67**, 399–408.
- (a) T. Kiss, J. Balla, G. Nagy, H. Kozłowski and T. Jankowska, *Inorg. Chim. Acta*, 1987, **138**, 25–30; (b) T. Kiss and I. Lazar, in *Aminophosphonic and Aminophosphinic Acids: Chemistry and Biological Activity*, ed. P. V. Kukhar and H. R. Hudson, Wiley, NY, 2000, pp. 285–325.
- (a) G. Ch. de Witt, P. M. May, J. Webb and G. Hefter, *BioMetals*, 1996, **9**, 351–361; (b) A. D. Sherry, J. Ren, J. Huskens, E. Brücher, E. Tóth, C. F. C. G. Geraldes, M. M. C. A. Castro and W. P. Cacheris, *Inorg. Chem.*, 1996, **35**, 4604–4612; (c) J. E. Bollinger and D. M. Roundhill, *Inorg. Chem.*, 1993, **32**, 2821–2826; (d) J. Gałęzowska, R. Janicki, A. Mondry, R. Burgada, T. Bailly, M. Lecouvey and H. Kozłowski, *Dalton Trans.*, 2006, 4384–4394; (e) A. Mondry and R. Janicki, *Dalton Trans.*, 2006, 4702–4710; (f) R. Janicki and A. Mondry, *Polyhedron*, 2008, **27**, 1942–1946; (g) R. Janicki and A. Mondry, *Eur. J. Inorg. Chem.*, 2013, 3429–3438; (h) R. Janicki, M. Monteil, M. Lecouvey and A. Mondry, *Opt. Mater.*, 2013, **36**, 259–264; (i) J. Gałęzowska, R. Janicki, H. Kozłowski, A. Mondry, P. Młynarz and Ł. Szyrwił, *Eur. J. Inorg. Chem.*, 2010, 1696–1702.



- 7 (a) J. C. G. Bünzli and C. Piguet, *Chem. Soc. Rev.*, 2005, **4**, 1048–1077; (b) J. C. G. Bünzli and S. V. Eliseeva, *Chem. Sci.*, 2013, **4**, 1939–1949.
- 8 J.-G. Mao, *Coord. Chem. Rev.*, 2007, **251**, 1493–1520.
- 9 (a) A. L. Tiwari, H. Ojha, A. Kaul, A. Dutta, P. Srivastava, G. Shukla, R. Srivastava and A. K. Mishra, *Chem. Biol. Drug Des.*, 2009, **7**, 87–91; (b) É. Tóth, L. Helm and A. E. Merbach, in *The Chemistry of Contrast Agents in Medical Magnetic Resonance Imaging*, ed. A. E. Merbach, L. Helm and É. Tóth, Wiley, Chichester, UK, 2nd edn, 2013, pp. 25–76; (c) S. Marinim, Y. Huang, D. Coman and F. Hyder, *J. Biol. Inorg. Chem.*, 2014, **19**, 1385–1398; (d) M. Elhabiri, S. Abada, M. Sy, A. Nonat, P. Choquet, D. Esteban-Gómez, C. Cassino, C. Platas-Iglesias, M. Botta and L. J. Charbonnière, *Chem. – Eur. J.*, 2015, **21**, 6535–6546.
- 10 A. D. Sherry and C. F. G. C. Geraldes, in *Lanthanide Probes in Life, Chemical and Earth Sciences, Theory and Practice*, ed. J. C. Bünzli and G. Choppin, Elsevier, Amsterdam, 1989, p. 93.
- 11 US FDA Approval for Quadramet, 28/09/97 (NDA) 020570.
- 12 (a) H. Studnik, S. Liebsch, G. Forlani, D. Wiczorek, P. Kafarski and J. Lipok, *New Biotechnol.*, 2015, **32**(1), 1–6; (b) B. Nowack, *Water Res.*, 2002, **36**, 4636–4642.
- 13 (a) P. Silva, F. Vieira, A. C. Gomes, D. Ananias, J. A. Fernandes, S. M. Bruno, R. Soares, A. A. Valente, J. Rocha and F. A. Almeida Paz, *J. Am. Chem. Soc.*, 2011, **133**, 15120–15138; (b) R. F. Mendes, P. Silva, M. M. Antunes, A. A. Valente and F. A. Almeida Paz, *Chem. Commun.*, 2015, **51**, 10807–10810.
- 14 K. Sawada, M. Kuribayashi, T. Suzuki and H. Miyamoto, *J. Solution Chem.*, 1991, **20**, 829–839.
- 15 R. C. Holz, G. E. Meister and W. Dew. Horrocks Jr., *Inorg. Chem.*, 1990, **29**, 5183–5189.
- 16 E. Kozlovski, S. Aleksandrov and L. Chesnokova, *Zh. Neorg. Khim.*, 2002, **47**, 1566–1568.
- 17 R. Janicki, *J. Mol. Struct.*, 2013, **1036**, 35–41.
- 18 (a) J. N. Brønsted, *J. Am. Chem. Soc.*, 1922, **44**, 877–898; (b) J. N. Brønsted, *J. Am. Chem. Soc.*, 1922, **44**, 938–948; (c) E. A. Guggenheim, *Philos. Mag.*, 1935, **19**, 588–643; (d) G. Scatchard, *Chem. Rev.*, 1936, **19**, 309–327.
- 19 R. S. Dickins, D. Parker, J. I. Bruce and D. J. Tozer, *Dalton Trans.*, 2003, 1264–1271.
- 20 (a) M. Pan, X. L. Zheng, Y. Liu, W. S. Liu and C. Y. Su, *Dalton Trans.*, 2009, 2157–2169; (b) Q. Y. Yang, K. Wu, J. J. Jiang, C. W. Hsu, M. Pan, J. M. Lehn and C. Y. Su, *Chem. Commun.*, 2014, **50**, 7702–7704; (c) B. B. Du, a Y. X. Zhu, M. Pan, M. Q. Yue, Y. J. Hou, K. Wu, L. Yin Zhang, L. Chen, S. Y. Yin, Y. N. Fana and C. Y. Su, *Chem. Commun.*, 2015, **51**, 12533–12536; (d) E. Kasprzycka, V. A. Trush, V. M. Amirkhanov, L. Jerzykiewicz, O. Malta, J. Legendziewicz and P. Gawryszewska, *Chem. – Eur. J.*, 2017, **23**, 1318–1330.
- 21 (a) C. Görller-Walrand, C. L. Fluyt, A. Ceulemans and W. T. Carnall, *J. Chem. Phys.*, 1991, **95**, 3099; (b) C. Görller-Walrand and K. Binnemans, in *Handbook on the Physics and Chemistry of Rare Earths Vol. 25*, ed. K. A. Gschneidner and L. Eyring, Elsevier, 1998, vol. 167, pp. 121–125.
- 22 (a) M. Latva, H. Takalo, V. M. Mikkala, C. Matachescu, J. C. Rodriguez-Ubis and J. Kankare, *J. Lumin.*, 1995, **75**, 149–169; (b) L. Smętek and A. Kędziorowski, *J. Lumin.*, 2010, **130**, 1154–1159.
- 23 (a) R. Janicki and A. Mondry, *Phys. Chem. Chem. Phys.*, 2015, **16**, 26823–26831; (b) R. Janicki and A. Mondry, *Phys. Chem. Chem. Phys.*, 2015, **17**, 29558–29565.
- 24 K. Bukietyńska, A. Mondry, P. N. Thuy and P. Starynowicz, *J. Alloys Compd.*, 1995, **225**, 52–54.
- 25 C. K. Sawada, T. Araki and T. Suzuki, *Inorg. Chem.*, 1987, **26**, 1199.
- 26 G. Grossmann, K. A. Burkov, G. Hägele, L. A. Myund, S. Hermens, C. Verwey and S. M. Arat-ool, *Inorg. Chim. Acta*, 2004, **357**, 797.
- 27 V. Deluchat, J.-C. Bolliner, B. Shepaud and C. Caullet, *Talanta*, 1997, **44**, 897.
- 28 B. Kurzak, A. Kamecka, K. Kurzak, J. Jezierska and P. Kafarski, *Polyhedron*, 2000, **19**, 2083.
- 29 (a) G. Anderegg, *Pure Appl. Chem.*, 1982, **54**, 2693–2758; (b) K. Bukietyńska and A. Mondry, *Inorg. Chim. Acta*, 1987, **130**, 271–276.
- 30 I. Grenthe, F. Mompean, K. Spahiu and H. Wanner, *Guidelines for the extrapolation to zero ionic strength*, OECD Nuclear Energy Agency, Data Bank, 2013.
- 31 G. M. Sheldrick, *SHELXS-97, program for structure solution*, University of Göttingen, 1997.
- 32 G. M. Sheldrick, *SHELXL-97, program for structure refinement*, University of Göttingen, 1997.
- 33 40 XP-Interactive Molecular Graphics, v. 5.1–Bruker Analytical X-ray Systems 1998.
- 34 H. M. Irving, M. G. Miles and L. D. Pettit, *Anal. Chim. Acta*, 1967, **38**, 475–488.
- 35 G. Gran, *Acta Chem. Scand.*, 1950, **4**, 559–577; G. Gran, *Analyst*, 1952, **77**, 661–671.
- 36 F. H. Sweeton, R. E. Mesmer and C. F. Baes Jr., *J. Solution Chem.*, 1974, **3**(3), 191–214.
- 37 P. Gans, A. Sabatini and A. Vacca, *J. Chem. Soc., Dalton Trans.*, 1985, 1195–1200.
- 38 P. Gans, A. Sabatini and A. Vacca, *Talanta*, 1996, **43**, 1739–1753.
- 39 P. Gans, *Data Fitting in the Chemical Sciences*, John Wiley & Sons, Chichester, 1992.
- 40 L. Alderighi, P. Gans, A. Ienco, D. Peters, A. Sabatini and A. Vacca, *Coord. Chem. Rev.*, 1999, **184**, 311–318.
- 41 N. Saha and H. Sigel, *J. Am. Chem. Soc.*, 1982, **104**, 4100.
- 42 S. Kim and R. Martin, *J. Am. Chem. Soc.*, 1984, **106**, 1707.
- 43 W. N. Perera and G. Senanayake, *Inorg. Chem.*, 2004, **43**, 3048.

