



Cite this: *RSC Adv.*, 2017, 7, 30376

# Task-specific acidic ionic liquid-catalyzed efficient synthesis of $\beta$ -enamino lactones from alkynoates and $\beta$ -amino alcohols†

Lu Chen, \* Bin Chen, Fuying Zhao, Yibiao Li, Bin Li\* and Min Zhang

By employing task-specific acidic ionic liquid as an efficient catalyst, a new method for the straightforward synthesis of  $\beta$ -enamino lactones has been demonstrated. A series of alkynoates in combination with various  $\beta$ -amino alcohols was efficiently converted into the desired products in good to excellent yields upon isolation. The skeleton of the seven-membered ring is generated *via* tandem intermolecular hydroamination and intramolecular transesterification processes. The developed synthetic protocol furnishes the desired products in a step- and atom-economic fashion with the advantages of high yields, broad substrate scope, good functional tolerance, and operational simplicity, offering an important basis for the construction of  $\beta$ -enamino lactones.

Received 9th April 2017  
 Accepted 23rd May 2017

DOI: 10.1039/c7ra04028h

[rsc.li/rsc-advances](http://rsc.li/rsc-advances)


$\beta$ -Amino acids constitute a highly significant branch in organic chemicals, which have been found to possess diverse bioactivities and are employed as useful building blockings for the preparation of  $\beta$ -lactam antibiotics and heterocycles. In addition,  $\beta$ -amino acid derivatives frequently occur in numerous biologically active natural products.<sup>1</sup> Among various compounds investigated to date,  $\beta$ -enamino lactone derivatives, the precursors of enantiopure  $\beta$ -amino acids, are the main structural components of many bioactive natural products that have attracted significant attention.<sup>2</sup>

Owing to the abovementioned interesting functions of  $\beta$ -enamino lactone, several synthetic protocols have been nicely demonstrated during the past few decades to access these compounds.<sup>3–8</sup> For instance, Abarbri and coworkers<sup>3</sup> first reported the reaction of ethyl perfluorobut-2-ynoate with amino alcohols to generate 5-(perfluoroalkyl)-3,4-dihydro-2H-[1,4]oxazepin-7-ones *via* the intermolecular Michael addition and lactone formation. In 2005, Dechoux's group reported an efficient method for the synthesis of  $\beta$ -enamino lactones<sup>4</sup> *via* the condensation of acetonedicarboxylate with  $\beta$ -amino alcohols followed by an intramolecular cyclization step. However, during this synthesis, the reaction intermediates needed to be isolated and it required excess NaH to achieve the cyclization; moreover, an additional neutralization manipulation by adding  $\text{NH}_4\text{Cl}/\text{H}_2\text{O}$  was essential to obtain the desired products after the completion of the reaction. From the viewpoint of green

chemistry, the development of efficient shortcuts for accessing  $\beta$ -enamino lactones from easily available feedstocks would be of high significance.

As a part of our program aimed at developing new synthetic methodologies for the construction of heterocycles,<sup>17</sup> we initially had the idea to develop a ruthenium-catalyzed synthesis of ester-substituted pyrrole **3a'** from alkynoate **1a** and  $\beta$ -amino alcohol **2a** *via* dehydrogenative cyclization (Scheme 1, eqn (1)). However, we failed to obtain even traces of the anticipated product, and a small portion of  $\beta$ -enamino lactone **3a** was obtained. Further investigations showed that the ruthenium catalyst was not essential for the product formation (**3a**), whereas the presence of 10 mol% of  $\text{AlCl}_3$  was able to improve the yield of **3a** to 18% (eqn (2)). Upon a thorough investigation of this new observation, a straightforward method for the efficient synthesis of  $\beta$ -enamino lactones from alkynoates and  $\beta$ -amino alcohols using a task-specific sulfonic ionic liquid as the catalyst was realized and has been reported herein.

Our initial investigation was to develop a more efficient reaction system by choosing the synthesis of  $\beta$ -enamino lactone **3a** from diethyl but-2-ynedioate **1a** and 2-amino-2-phenylethanol **2a** as a model reaction. First, the reaction in



Scheme 1 The new observation leading to  $\beta$ -enamino lactone.

School of Chemical & Environmental Engineering, Wuyi University, Jiangmen, Guangdong Province 529090, China. E-mail: [wuyuchemcl@126.com](mailto:wuyuchemcl@126.com); [andonlee@163.com](mailto:andonlee@163.com)

† Electronic supplementary information (ESI) available: General experimental information, experimental procedure for product synthesis, full characterization data,  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of all products. See DOI: 10.1039/c7ra04028h




*t*-amyl alcohol was performed at 120 °C for 12 h, and several conventional Lewis acid catalysts were tested (Table 1, entries 1–7); it was found that Cu(OTf)<sub>2</sub> or Zn(OTf)<sub>2</sub> shows good performance in affording the desired product **3a**. However, in the absence of an acid catalyst, the desired product was not obtained (Table 1, entry 8), indicating that the acidic catalyst played a crucial role in the reaction. Moreover, attracted by the significant advantages, such as the designability, easy recovery, and reusability,<sup>9–15</sup> of task-specific acidic ionic liquids, we evaluated the utility of a sulfonic-functionalized ionic liquid<sup>16</sup> (TSFIL-1: [TMBSA]HSO<sub>4</sub>) as a catalyst. Gratifyingly, this catalyst exhibited an excellent activity in the production of product **3a** (entries 9–11), and 10 mol% catalyst loading was essential to afford a satisfactory yield (entry 9). Further, changes in the reaction temperatures led to diminished product yields (Table 1, entries 12 and 13), implying that a lower temperature significantly decreased the reaction rate, whereas increasing this parameter would lower the product stability. However, use of toluene, DMF, DMAc, DMSO, and 1,4-dioxane as the solvents proved to be totally ineffective or less effective for the formation

of products when compared with *t*-amyl alcohol (Table 1, entries 14–18). Finally, [TMBSA]HSO<sub>4</sub> was used as both the solvent and the catalyst and it only gave a 77% yield, which could be rationalized since a viscous ionic liquid was not beneficial for the interaction of substrates. Based on these results, the optimal reaction conditions were obtained and are indicated in entry 9 of Table 1.

With the availability of the optimized reaction conditions, we subsequently examined the generality of the synthetic protocol. A variety of alkynoates (**1**) in combination with β-amino alcohols (**2**) were tested. As shown in Table 2, all the reactions smoothly proceeded and furnished the desired products in good to excellent isolated yields. Diethyl but-2-ynedioate **1a** reacted with 2-amino-2-phenylethanol **2a** and 2-amino-3-phenylpropan-1-ol **2c** to provide the products **3a** and **3c** in 86% and 94% yields, respectively (Table 2, entries 1 and 3). Moreover, even a steric-hindered substrate, such as 2-amino-1,2-diphenylethanol **2b**, also underwent smooth transformation with 2-amino-2-phenylethanol **2a**, affording the desired product in 78% yield (Table 2, entry 2). Similarly, the reactions of **1a** with amino alcohols **2d** and **2e** produced the corresponding products **3d** and **3e** in excellent yields (Table 2, entries 4 and 5), respectively. Interestingly, the less reactive ethyl 3-phenylpropionate **1c** could also generate the desired coupling products in moderate to good yields, demonstrating that the developed chemistry was applicable for a broad substrate scope (Table 2, entries 6–10). Note that amino alcohols (**2a** and **2c**) with a phenyl group or a benzyl substituent could afford higher yields (Table 2, entries 6 and 9) than those with an alkyl group (Table 2, entries 1 and 3–5). Moreover, amino alcohols with an isopropyl group gave a relatively lower yield (Table 2, entry 8), presumably because of the influence of its strong electron-donating effect, thus deactivating the ester group. On the other hand, the secondary alcohols such as 1-aminopropan-2-ol **2f** reacted with alkynoate **1c** to give the corresponding product **3j** in 74% yield (Table 2, entry 10). Note that various functional groups such as 4-Cl, 4-Br, 4-F, and 4-CH<sub>3</sub>CO on the phenyl ring of alkynoates (**1d–1g**) were well tolerated, affording the corresponding products in good to excellent yields (Table 2, entries 12–14, 19 and 20). Similarly, electron-donating groups on the aryl ring of the substrates **2** were also compatible with the transformation (Table 2, entries 15–17). The retention of these functional groups would offer the potential for further molecular complexity *via* chemical transformation.

To gain insight into the reaction information, we performed the control experiments. It was found that hydroamination between alkynoate **1a** and amino alcohol **2a** completed in 5 minutes without any catalyst or additive. Then, the resulting enamine intermediate (**3a-I**) under standard conditions furnished the cyclization product **3a** in an almost quantitative yield (Scheme 2). This result, in combination with the fact that in the absence of a catalyst, product **3a** cannot be formed (Table 1, entry 8), indicates that the acidic catalyst plays a crucial role in the activation of the ester group. These findings suggest that the product formation is initiated by fast intermolecular hydroamination followed by [TMBSA]HSO<sub>4</sub>-catalyzed intramolecular transesterification.


Table 1 Screening of the reaction conditions<sup>a</sup>



Entry	Catalyst (mol%)	Temp. (°C)	Solvent (2.0 mL)	Yield <sup>b</sup> (%)
1 <sup>c</sup>	AlCl <sub>3</sub> (10)	120	<i>t</i> -Amyl alcohol	18
2	ZnCl <sub>2</sub> (10)	120	<i>t</i> -Amyl alcohol	35
3	FeCl <sub>3</sub> (10)	120	<i>t</i> -Amyl alcohol	40
4	Yb(OTf) <sub>3</sub> (10)	120	<i>t</i> -Amyl alcohol	40
5	Cu(OTf) <sub>2</sub> (10)	120	<i>t</i> -Amyl alcohol	78
6	Al(OTf) <sub>3</sub> (10)	120	<i>t</i> -Amyl alcohol	80
7	Zn(OTf) <sub>2</sub> (10)	120	<i>t</i> -Amyl alcohol	84
8	None	120	<i>t</i> -Amyl alcohol	— <sup>d</sup>
9	TSILs (10)	120	<i>t</i> -Amyl alcohol	94
10	TSILs (5)	120	<i>t</i> -Amyl alcohol	92
11	TSILs (15)	120	<i>t</i> -Amyl alcohol	90
12 <sup>c</sup>	TSILs (10)	100	<i>t</i> -Amyl alcohol	36
13	TSILs (10)	140	<i>t</i> -Amyl alcohol	87
14	TSILs (10)	120	Toluene	— <sup>d</sup>
15	TSILs (10)	120	DMF	— <sup>d</sup>
16	TSILs (10)	120	DMAc	22
17	TSILs (10)	120	DMSO	56
18	TSILs (10)	120	1,4-Dioxane	14
19	TSILs (1 mL)	120	None	77


<sup>a</sup> Reaction conditions: unless otherwise stated, the reaction mixture of dimethyl acetylenedicarboxylate (**1**) (0.5 mmol), 2-amino-2-phenylethanol (**2**) (0.6 mmol), catalyst loaded in different solvents (2.0 mL) was stirred at 120 °C for 12 h under atmospheric condition. <sup>b</sup> GC yield using *n*-hexadecane as an internal standard. <sup>c</sup> Under a N<sub>2</sub> atmosphere. <sup>d</sup> No product detected.



Table 2 Efficient synthesis of 3,4-dihydro-1,4-oxazepin-7(2H)-one derivatives<sup>a</sup>


Entry	1	2	3, yield (%)
1	1a: R <sup>1</sup> = Et, R <sup>2</sup> = CO <sub>2</sub> Et	2a: R <sup>3</sup> = Ph, R <sup>4</sup> = H	3a, 86
2	1a: R <sup>1</sup> = Et, R <sup>2</sup> = CO <sub>2</sub> Et	2b: R <sup>3</sup> = Ph, R <sup>4</sup> = Ph	3b, 78
3	1a: R <sup>1</sup> = Et, R <sup>2</sup> = CO <sub>2</sub> Et	2c: R <sup>3</sup> = PhCH <sub>2</sub> , R <sup>4</sup> = H	3c, 94
4	1b: R <sup>1</sup> = CH <sub>3</sub> , R <sup>2</sup> = CO <sub>2</sub> CH <sub>3</sub>	2a: R <sup>3</sup> = Ph, R <sup>4</sup> = H	3d, 80
5	1b: R <sup>1</sup> = CH <sub>3</sub> , R <sup>2</sup> = CO <sub>2</sub> CH <sub>3</sub>	2c: R <sup>3</sup> = PhCH <sub>2</sub> , R <sup>4</sup> = H	3e, 90
6	1c: R <sup>1</sup> = Et, R <sup>2</sup> = Ph	2a: R <sup>3</sup> = Ph, R <sup>4</sup> = H	3f, 68
7	1c: R <sup>1</sup> = Et, R <sup>2</sup> = Ph	2d: R <sup>3</sup> = CH <sub>3</sub> , R <sup>4</sup> = H	3g, 78
8	1c: R <sup>1</sup> = Et, R <sub>2</sub> = Ph	2e: R <sup>3</sup> = (CH <sub>3</sub> ) <sub>2</sub> CH, R <sup>4</sup> = H	3h, 58
9	1c: R <sup>1</sup> = Et, R <sup>2</sup> = Ph	2c: R <sup>3</sup> = PhCH <sub>2</sub> , R <sup>4</sup> = H	3i, 68
10	1c: R <sup>1</sup> = Et, R <sup>2</sup> = Ph	2f: R <sup>3</sup> = H, R <sup>4</sup> = CH <sub>3</sub>	3j, 74
11	1d: R <sup>1</sup> = Et, R <sup>2</sup> = 4-ClPh	2d: R <sup>3</sup> = CH <sub>3</sub> , R <sup>4</sup> = H	3k, 84
12	1e: R <sup>1</sup> = Et, R <sup>2</sup> = 4-BrPh	2d: R <sup>3</sup> = CH <sub>3</sub> , R <sup>4</sup> = H	3l, 89
13	1d: R <sup>1</sup> = Et, R <sup>2</sup> = 4-ClPh	2e: R <sup>3</sup> = (CH <sub>3</sub> ) <sub>2</sub> CH, R <sup>4</sup> = H	3m, 60
14	1e: R <sup>1</sup> = Et, R <sup>2</sup> = 4-BrPh	2e: R <sup>3</sup> = (CH <sub>3</sub> ) <sub>2</sub> CH, R <sup>4</sup> = H	3n, 62
15	1h: R <sup>1</sup> = Et, R <sup>2</sup> = 4-CH <sub>3</sub> Ph	2d: R <sup>3</sup> = CH <sub>3</sub> , R <sup>4</sup> = H	3o, 72
16	1i: R <sup>1</sup> = Et, R <sup>2</sup> = 4-CH <sub>3</sub> OPh	2d: R <sup>3</sup> = CH <sub>3</sub> , R <sup>4</sup> = H	3p, 70
17	1h: R <sup>1</sup> = Et, R <sup>2</sup> = 4-CH <sub>3</sub> Ph	2e: R <sup>3</sup> = (CH <sub>3</sub> ) <sub>2</sub> CH, R <sup>4</sup> = H	3q, 56
18	1i: R <sup>1</sup> = Et, R <sup>2</sup> = 4-CH <sub>3</sub> OPh	2e: R <sup>3</sup> = (CH <sub>3</sub> ) <sub>2</sub> CH, R <sup>4</sup> = H	3r, 50
19	1f: R <sup>1</sup> = Et, R <sup>2</sup> = 4-FPh	2d: R <sup>3</sup> = CH <sub>3</sub> , R <sup>4</sup> = H	3s, 70
20	1g: R <sup>1</sup> = Et, R <sup>2</sup> = 4-CH <sub>3</sub> COPh	2d: R <sup>3</sup> = CH <sub>3</sub> , R <sup>4</sup> = H	3t, 75

<sup>a</sup> Reaction conditions: unless otherwise stated, the reaction mixture of dimethyl acetylenedicarboxylate (**1**) (0.5 mmol), 2-amino-2-phenylethanol (**2**) (0.6 mmol), and [TMBSA]HSO<sub>4</sub> (10 mol%) in *t*-amyl alcohol (2.0 mL) was stirred at 120 °C for 12 h under atmospheric condition.



Scheme 2 Control experiments.

Finally, we demonstrated the utility of the developed new method. The reaction of the diethyl but-2-ynedioate **1a** with enantiopure amino alcohol (*S*)-(+)-**2a'** produced the enantiopure enamino lactone (*S*)-(+)-**3a'** in an excellent yield with retention of the chiral configuration (Scheme 3). This result shows that the task-specific acidic ionic liquid-catalyzed protocol is also

Scheme 3 Synthesis of chiral  $\beta$ -enaminolactone.

applicable for the preparation of chiral  $\beta$ -enaminolactones from chiral amino alcohols.

In summary, by employing a task-specific sulfonic ionic liquid as the catalyst, we demonstrated an environmentally friendly and straightforward approach for the versatile synthesis of  $\beta$ -enaminolactones from readily available amino alcohols and alkynoates for the first time. The synthetic protocol proceeds *via* tandem intermolecular hydroamination and intramolecular transesterification processes; moreover, it furnished the desired products in a step- and atom-economic fashion with the advantages of high isolated yields, broad substrate scope, good functional tolerance, and operational simplicity, which offers an important basis for the construction of  $\beta$ -enaminolactones.

## Acknowledgements

The authors are grateful to the funds of the Science Foundation for Young Teachers of Wuyi University (No: 2016zk03, 2014zk04 and No: 2015td01), the Scientific Research Foundation for the Returned Overseas as Chinese Scholar, State Education Ministry ([2015]1098), the Natural Science Foundation of Guangdong Province (No: 2014A030310211), and the Foundation of the Department of Education of Guangdong Province (No: 2016KTSCX144).



## Notes and references

- (a) O. W. Griffith, *Annu. Rev. Biochem.*, 1986, **55**, 855–878; (b) E. Juaristi, D. Quintania and J. Escalante, *Aldrichimica Acta*, 1994, **27**, 3–11; (c) J. S. Carey, D. Laffan, C. Thomson and M. T. Williams, *Org. Biomol. Chem.*, 2006, **4**, 2337–2347.
- (a) D. Seebach and J. L. Matthews, *Chem. Commun.*, 1997, 2015–2022; (b) R. P. Cheng, S. H. Gellman and W. F. DeGrado, *Chem. Rev.*, 2001, **101**, 3219–3232.
- S. Richard, G. Prie, A. Guignard, J. Thibonnet, J. L. Parrain, A. Duchene and M. Abarbri, *Helv. Chim. Acta*, 2003, **86**, 726–732.
- J. Alladoum and L. Dechoux, *Tetrahedron Lett.*, 2005, **46**, 8203–8205.
- J. Alladoum, V. Toum, S. Hebbe, C. Kadouri-Puchot and L. Dechoux, *Tetrahedron Lett.*, 2009, **50**, 617–619.
- Y. Ishida, N. Iwahashi, N. Nishizono and K. Saigo, *Tetrahedron Lett.*, 2009, **50**, 1889–1892.
- A. K. Singh, R. Chawla and L. D. S. Yadav, *Synthesis*, 2012, **44**, 2353–2358.
- Y. Tian, X. Wang, Q. Xiao, C. Sun and D. Yin, *J. Org. Chem.*, 2014, **79**, 9678–9685.
- M. Petkovic, K. R. Seddon, L. P. N. Rebelo and C. S. Pereira, *Chem. Soc. Rev.*, 2011, **40**, 1383–1403.
- N. Isambert, M. M. S. Duque, J.-C. Plaquevent, Y. Génisson, J. Rodriguez and T. Constantieux, *Chem. Soc. Rev.*, 2011, **40**, 1347–1357.
- R. Sheldon, *Chem. Commun.*, 2001, **23**, 2399–2407.
- T. Welton, *Chem. Rev.*, 1999, **99**, 2071–2083.
- J. P. Hallett and T. Welton, *Chem. Rev.*, 2011, **111**, 3508–3576.
- T. Welton, *Coord. Chem. Rev.*, 2004, **248**, 2459–2477.
- A. S. Amarasekara, *Chem. Rev.*, 2016, **116**, 6133–6183.
- (a) S. G. Lee, *Chem. Commun.*, 2006, 1049–1063; (b) G. Y. Zhao, T. Jiang, H. X. Gao, B. X. Han, J. Huang and D. H. Sun, *Green Chem.*, 2004, **6**, 75–77; (c) D. Q. Xu, J. Wu, S. P. Luo, J. Zhang, J. Y. Wu, X. H. Du and Z. Y. Xu, *Green Chem.*, 2009, **11**, 1246.
- (a) C. B. Bheeter, L. Chen, J.-F. Soulé and H. Doucet, *Catal. Sci. Technol.*, 2016, **6**, 2005–2049; (b) L. Chen, X. Luo and Y. Li, *Monatsh. Chem.*, 2017, **148**, 957–961; (c) J. Cai, S. Huang, R. He, L. Chen, D. Chen, S. Jiang, B. Li and Y. Li, *Org. Biomol. Chem.*, 2017, **15**, 333–337; (d) Y. Li, L. Cheng, B. Li, S. Jiang, L. Chen and Y. Shao, *ChemistrySelect*, 2016, **5**, 1092–1095.

