# **RSC Advances**



View Article Online

View Journal | View Issue

### PAPER

Check for updates

Cite this: RSC Adv., 2017, 7, 41955

# Chemoselective acylation of 2-amino-8-quinolinol in the generation of C2-amides or C8-esters<sup>†</sup>

Yongseok Park,‡<sup>a</sup> Xiang Fei,‡<sup>a</sup> Yue Yuan,<sup>a</sup> Sanha Lee,<sup>a</sup> Joonseong Hur,<sup>b</sup> Sung Jean Park,<sup>a</sup> Jae-Kyung Jung<sup>b</sup><sup>c</sup> and Seung-Yong Seo<sup>b</sup>\*<sup>a</sup>

Two different ways to carry out the chemoselective acylation of 2-amino-8-quinolinol with unique features to generate C2-amides or C8-esters were developed. The coupling reaction with a variety of carboxylic acids using EDCI and DMAP provided C8-ester derivatives, whereas *N*-heteroaromatic acids were not introduced on the C8-hydroxy group, but rather on the C2-amino group under the same conditions. To obtain C2-amides selectively, the anionic nucleophile from 2-amino-8-quinolinol was treated with less reactive acyl imidazolides or esters.

Received 10th May 2017 Accepted 23rd August 2017

DOI: 10.1039/c7ra05287a

rsc.li/rsc-advances

#### Introduction

Ester and amide functionalities are important structural components that serve as key motifs in a variety of bioactive natural products and pharmaceuticals. Amide and ester formation are fundamental reactions in chemical synthesis.1 Typically, carbodiimides such as dicyclohexylcarbodiimide (DCC) and 1-ethyl-3-(3'-dimethylaminopropyl)-carbodiimide (EDCI) are applied as coupling reagents for esterification and amidation reactions. To increase the overall rate of the coupling reaction, DMAP (4-dimethylaminopyridine), 1-hydroxy-1H-benzotriazole (HOBt) and 1-hydroxy-7-azabenzotriazole (HOAt) are commonly used for carbodiimide-mediated reactions in either stoichiometric or catalytic fashions. Meanwhile, 1,1'-carbonyl diimidazole (CDI) is used because of the enhanced stability relative to that of acid halides and viability in large-scale syntheses.<sup>2</sup> In addition, numerous other methods have been developed for ester and amide bond formation.<sup>1c</sup>

Quinolines are valuable synthons and precursors in synthetic organic chemistry, as they comprise important heterocyclic pharmacophores with a broad spectrum of biological and me-dicinal activities. In particular, 8-hydroxyquinoline (8-HQ) is a well-known metal ion chelator and represents an excellent scaffold with a wide spectrum of pharmacological applica-tions.<sup>3</sup> Similarly, 2-aminoquinoline (2-AQ) exhibits various biological activities such as anthelmintic, antiprotozoal, anti-depressant, some effectiveness in the treatment of Alzheimer's disease.<sup>4</sup> Toward that end, we paid attention to unique 2-amino-8-quinolinol (1), which contains important structural components of 8-HQ and 2-AQ.<sup>5</sup> The synthesis of its derivatives may prove to be useful, owing to the utility of the parent compounds; however, the previous works for *O*- and *N*-acylation of **1** have hardly been reported to date.<sup>6</sup> We sought the synthesis of its derivatives, especially the esterification on C8 and the amidation on C2 (Fig. 1).

Recently, we reported an acyl derivative of 2-amino-8quinolinol (1) called **SG-HQ2**. This compound is an antiallergic inflammatory agent.<sup>7</sup> Unfortunately, NMR spectroscopic studies cast doubt on the originally proposed structure, which was regarded as the amide form. To prove that the ester form was obtained as the major product of the EDCImediated coupling reaction of 2-amino-8-quinolinol (1) and 3,4,5-tris(benzyloxy)benzoic acid, we reexamined the acylation and analyzed the structure of **SG-HQ1** *via* NMR spectroscopy and X-ray crystallography.<sup>8</sup> **SG-HQ2** from the catalytic hydrogenation of SG-HG1 was determined to be the C8-ester instead of the amide as reported previously (Scheme 1).

#### Results and discussion

Encouraged by the chemoselective *O*-acylation of 2-amino-8quinolinol, our synthetic investigations regarding C2-amide derivatives began with initial studies of the 2 step synthesis



Fig. 1 2-Amino-8-quinolinol (1) and its acyl derivatives.

<sup>&</sup>lt;sup>a</sup>College of Pharmacy and Gachon Institute of Pharmaceutical Sciences, Gachon University, Incheon 21936, South Korea

<sup>&</sup>lt;sup>b</sup>College of Pharmacy, Seoul National University, Seoul 08826, South Korea

<sup>&</sup>lt;sup>c</sup>College of Pharmacy, Chungbuk National University, Cheongju, Chungbuk 28644, South Korea

<sup>†</sup> Electronic supplementary information (ESI) available: Experimental procedures, spectral and analytical data (PDF) and X-ray crystallographic analyses of SG-HQ1 (3f), 3b, and 4b (CIF). CCDC 1548392–1548394. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7ra05287a

<sup>‡</sup> Both authors contributed equally to this manuscript.



 $\mbox{Scheme 1}$  Synthesis of  $\mbox{SG-HQ1}$  and  $\mbox{SG-HQ2}$  and their revised structures.  $^{8}$ 

comprised of the diacylation with excess of aromatic acid (up to 5 equiv.) and the hydrolysis of the ester moiety (Scheme 2). C2amides of various aromatic acids were prepared, and the structures of **3b** and **4b** were confirmed by X-ray crystallography.<sup>8</sup> Interestingly, the acylation with heteroaromatic acids such as 2-picolinic acid and pyrazinecarboxylic acid afforded only the C2-amides (**4h** and **4i**), excluding C8-ester and diacyl compounds, albeit in low yield. Thus, the selective and efficient introduction of various carboxylic acids on 2-amino-8quinolinol remains a challenge,<sup>9</sup> although some C2 and C8-acyl derivatives were prepared successfully. Inspired by the unique characteristics of 2-amino-8-quinolinol, herein we present the optimization of the chemoselective *O*- or *N*-acylation.

To investigate possible synthetic routes towards C2-amide and C8-ester derivatives of 2-amino-8-quinolinol, we first examined various acylation conditions with 4-methoxybenzoic acid (Table 1). THF was selected as the most favorable solvent in terms of solubility and purification. The use of carbodiimides provided the C8-ester **3a** preferentially as the major product.



Scheme 2 Initial synthesis of C8-ester and C2-amide derivatives of 2amino-8-quinolinol. Aromatic acids including 4-methoxybenzoic (a), 4-chlorobenzoic (b), 4-(trifluoromethyl)benzoic (c), 4-*tert*-butylbenzoic (d), 3-methoxybenzoic (e), 3,4,5-tris(benzyloxy)benzoic (f), 2furoic (g), picolinic (h), and pyrazinecarboxylic (i) acids.

DMAP as an additive increased the chemical yield and ester selectivity (entry 1 and 7). However, when HOBt was employed, the formation of diacyl derivative 5a increased with that of C8-ester (entry 3-5). Though the use of HOAt provided the C2ester selectively, the isolated yield of 3a was not satisfactory (entry 6). Uranium (HATU) and phosphonium (PyBop) salts did not exhibit significant effects (entry 8 and 9). The use of 4-methoxybenzoyl chloride with iPr2NEt gave the C8-ester as the major product (entry 10). The yield decreased slightly when pyridine was used as a solvent (entry 11). Finally, we chose CDI for the amidation through the acyl imidazolide, leading to the formation of C2-amide 4a although the C8-ester formed as the major product. Higher temperature increased the formation of C2-amide slightly (entry 12–14). C2-amide 4a could be obtained in 75% yield with the ratio of 5:88:7, when the anionic nucleophile by the treatment of 2-amino-8-quinolinol with NaH reacted with the acyl imidazolide derived from 4-methoxybenzoic acid and CDI (entry 15).

The optimized acylation conditions were further investigated using a cheap and readily available ester instead of acyl imidazolide **6** with other strong bases (Table 2).<sup>10</sup> Methyl 4methoxybenzoate 7 was used as an acyl donor. Even though the use of the methyl ester provided a lower yield than acyl imidazolide **6**, the acylation resulted in the selective formation of C2-amide. When NaH and *n*-BuLi were used to generate the anionic intermediate of 2-amino-8-quinolinol, the excellent selectivity was observed (entries 2 and 3). The use of iPrMgCl as a base resulted in a significant loss of reactivity and selectivity (entry 4). The formation of C2-amide **4a** was improved as the amount of *t*-BuOK was increased (entry 5–8). The anionic nucleophile of 2-amino-8-quinolinol could mostly be converted into C2-amide rather than C8-ester.

Additive effect on reactivity and selectivity was evaluated in order to reveal the role of acyl imidazolide in C8-acylation. Some amines (such as pyridine, DMAP, and HOBt) and their HCl salts were treated in the reaction of **1** with the isolated acyl imidazolide **6** (Table 3). Without additives, C2-amide (**4a**) was obtained even though the reaction yield and ratio was slightly different with CDI-mediated acylation. However, the use of pyridine and DMAP at reflux provided C8-ester (**3a**) reversely. The use of their HCl salt (Pyr·HCl and DMAP·HCl) and HOBt provided selectively the C8-ester in good yield even at room temperature.

A variety of carboxylic acids were introduced on 2-amino-8quinolinol under the optimized acylation conditions. Either condition A (EDCI, iPr<sub>2</sub>NEt and catalytic DMAP) or condition B (PyBop and iPr<sub>2</sub>NEt) was used for the synthesis of the C8-esters (Table 4). Various substituted-benzoic acids and furoic acid (*i.e.* **3a–3f**, **3j** and **3k**) were introduced on the C8 position in good yields, regardless of electron-withdrawing and donating groups. However, *N*-heteroaromatic acids were not incorporated on the phenol group in the C8 position; C2-amides (**4h–4i**, **4l– 4m**) were isolated even in low yields instead. Interestingly, Bocleucinyl C8-ester (**3n**) seems to be unstable during flash column chromatography like heteroaromatic acid, however Boc-valinyl C8-ester (**3o**) was obtained in moderate yield.

#### Table 1 Optimization of acylation conditions<sup>4</sup>



Ratio <sup>c</sup> ( <b>3a : 4a : 5a</b> )
0:<1
0:<1
: 32
: 22
: 20
:7
:1
: 17
: 41
: 13
:19
2:8
9:24
3:33
:7

<sup>*a*</sup> The reactions were carried out three times. <sup>*b*</sup> Isolated yield of **3a** and **4a** after column chromatography. <sup>*c*</sup> Ratio of **3a**, **4a** and **5a** determined by HPLC analysis. <sup>*d*</sup> At reflux. <sup>*e*</sup> 4-Methoxybenzoyl chloride was used. <sup>*f*</sup> Reaction conditions: a THF solution of CDI and 4-methoxybenzoic acid was added to 2-amino-8-quinolinol and NaH (2 equiv.) in THF.

 Table 2
 The amidation of 2-amino-8-quinolinol with acyl imidazole and ester under anionic conditions



Entry	Acyl dollol	Dase (equiv.)	rinc (ii)	70 yielu	Ratio
1	6	NaH (2.0)	4	75	5:88:7
2	7	NaH (2.0)	24	25	2:98:0
3	7	<i>n</i> -BuLi (2.0)	4	56	0:99:1
4	7	iPrMgCl (2.0)	24	8	0:86:14
5	7	<i>t</i> -BuOK (1.0)	24	11	4:93:3
6	7	<i>t</i> -BuOK (2.0)	2	63	3:96:0
7	7	t-BuOK (2.5)	1	72	1:93:7
8	7	<i>t</i> -BuOK (3.0)	0.5	85	1:96:3
a <b>-</b> 1			(		

<sup>*a*</sup> Isolated yield of amide **3a**. <sup>*b*</sup> Ratio (**3a** : **4a** : **5a**) determined by HPLC analysis.

Next, we turned our attention to the amidation of 2-amino-8quinolinol on the C2 position. The CDI-mediated amidation of some aromatic acids with electron-withdrawing substituents provided C2-amides as the major product in contrast to carbodiimide-mediated acylation (condition C, Table 5). Nevertheless, this general amidation was accompanied by the formation of the C8-ester species, which was difficult to  
 Table 3
 The effect on additives in the reaction of 2-amino-8-quinolinol with acyl imidazole



<sup>a</sup> Reaction conditions: 2-amino-8-quinolinol (1 equiv.), acyl imidazolide (1.2 equiv.), an additive (1.5 equiv.), THF (4 mL), at room temperature.
 <sup>b</sup> Isolated yield. <sup>c</sup> Ratio determined by HPLC analysis. <sup>d</sup> At reflux.
 <sup>e</sup> Added iPr<sub>2</sub>NEt (3 equiv.).

separate. With the optimized reaction conditions in hand, the substrate scope was examined (condition D, Table 5). 4-Methoxybenzoyl derivatives (2a) was obtained in 75% yield, whereas the isolated yields of other substituted benzoic acids and heteroaromatic acids were improved up to 99% (4a vs. 4b-4m). Interestingly amides with *N*-protected amino acids (4n and

#### Table 4 Scope of C8-ester formation<sup>a,b</sup>



<sup>*a*</sup> Reaction conditions: 1 (0.2 mmol), carboxylic acid (0.4 mmol), EDCI (2.5 equiv.), DMAP (10 mol%), iPr<sub>2</sub>NEt (3 equiv.), THF (1 mL). <sup>*b*</sup> Isolated yield by flash column chromatography. <sup>*c*</sup> Not obtained C8-ester; isolated yield of C2-amide in parenthesis.

**4o**) were easily prepared from the corresponding acyl imidazolides.

By using EDCI-mediated acylation, the bulky aliphatic acyl moieties such as piperidinylcarboxy, isovaleroyl, pivaloyl and 1adamantylcarbonyl were introduced on the C8-hydroxy group successfully (3p-3s) as shown in Scheme 3. Similarly, sulfonyl chloride, benzyl chloroformate and Boc<sub>2</sub>O were used to prepare the sulfonate and carbonate derivatives (3t-3v) on the C8position in good yield, respectively. Meanwhile, the C2-amides with bulky aliphatic amides bearing piperidinylcarboxy, isovaleroyl, pivaloyl and 1-adamantylcarbonyl (4p-4s), were easily prepared from the corresponding acyl imidazolides. Finally, a similar method was used to prepare two carbamates with Cbz and Boc on C2 position, in spite of the unsatisfactory result obtained with *p*-toluenesulfonamide (4t *vs.* 4u and 4v).

Additive effects besides imidazole are shown in Table 3 to shed light on the mechanism of the chemoselective synthesis of C8-ester and C2-amide derivatives. Thus, representative amines (such as pyridine, DMAP, and HOBt) and their HCl salts were treated with the imidazolide **6** to afford the C8-ester (Scheme 4). With reactive acyl donors, C8-ester was regarded as the kinetic product. On the contrary, two C8-esters (**3a** and **3k**) were refluxed with imidazole to afford the corresponding amides (**4a** 



<sup>*a*</sup> Reaction conditions: a mixture of the carboxylic acid (1.2 equiv.) and CDI (1.3 equiv.) was added to a THF (4 mL) solution of 1 (0.20 mmol) and NaH (2 equiv.) at room temperature. <sup>*b*</sup> Isolated yield from flash column chromatography. <sup>*c*</sup> Isolated yield based on recovered starting material in parenthesis.

and 4k) in 40% and 82% yields, respectively. The observed conversion from the C8-ester to the C2-amide revealed that the C2-amide was the thermodynamically stable product. The conversion of 3k to 4k was superior to that of 3a, because of the electronic effect on substituents of benzene ring. This result is relevant to the lack of formation of C8-esters with N-heteroaromatic carbonyls (3h-3n) under most of conditions, as shown in Scheme 3. In order to determine the origin of the unique reactivity of 2-amino-8-quinolinol, 8-hydroxyquinoline (8) and 2-aminoquinoline (9) reacted with 4-methoxybenzoic acid under various coupling conditions. The yields with EDCI, PyBop, HBTU, and CDI were not satisfactory, but the acylation was successful with using the reactive acid chloride in good yields (up to 92%). As the weak reactivity of 8 and 9, it is considered to have a beneficial effect between the 2-amino and 8-hydroxy moieties for the introduction of an acyl group to 1.

To investigate the acylation pattern of **1**, we monitored the progress of C2-amide and C8-ester formation *via* CDI-mediated acylation by HPLC. Ester **3a** formed faster than amide **4a** (0–6 h). After 24 h, **4a** was the major component, while starting material **1**, **3a**, and **5a** were formed in similar amounts (almost 20%, Table 6).<sup>8</sup>



Scheme 3 The introduction of aliphaticacyl, sulfonyl and alkoxycarbonyl groups. <sup>a</sup>Reaction conditions: 1 (0.2 mmol), carboxylic acid (0.4 mmol), EDCI (2.5 equiv.), DMAP (10 mol%), iPr<sub>2</sub>NEt (3 equiv.), THF (1 mL). <sup>b</sup>Reaction conditions: a mixture of the carboxylic acid (1.2 equiv.) and CDI (1.3 equiv.) was added to a THF (4 mL) solution of 1 (0.20 mmol) and NaH (2 equiv.). <sup>c</sup>Isolated yield by flash column chromatography. <sup>d</sup>*p*-toluenesulfonyl chloride, benzyl chloroformate and Boc<sub>2</sub>O were used for 3t, 3u, and 3v respectively. <sup>e</sup>Isolated yield based on recovered starting material in parenthesis. <sup>f</sup>*p*-Toluene-sulfonyl chloride, benzyl chloroformate and Boc<sub>2</sub>O were used with imidazole for 4t, 4u, and 4v respectively, but 4t was not isolated.



Scheme 4 Mechanistic investigation.

Based on the above mechanistic investigation, a proposed reaction pathway is outlined in Scheme 5. Notably, Joussef *et al.* reported the chemoselectivity *O*-sulfonylation at C8.<sup>11</sup> For the C8-ester formation, the nitrogen nucleophile at N1 induced from the lone pair of the amino at C2 attacks acyl donors such as acid chlorides preferentially; this step is followed by an intraor intermolecular acyl transfer to the neighboring hydroxy at C8.<sup>12</sup> Meanwhile, the formation of C2-amides involves the anions of **1**, generated by strong bases such as NaH. We speculated that a sodium ion which interacts with the anion at C8 might coordinate to the nitrogen atom at N1, whereas the  
 Table 6
 The representative time-course data (HPLC) for the formation of C8-ester and C2-amide under CDI-mediated condition<sup>a</sup>



Time (h)	<b>1</b> (mmol)	3a (mmol)	<b>4a</b> (mmol)	5 <b>a</b> (mmol)
0	0.3000	0	0	0
1	0.2780	0.0201	0.0014	0.0005
3	0.2096	0.0593	0.0252	0.0060
6	0.1666	0.0697	0.0512	0.0125
12	0.0753	0.0628	0.1201	0.0418
24	0.0681	0.0656	0.1087	0.0576

 $^a$  Reaction conditions: a THF (4 mL) solution of 2-amino-8-quinolinol (1 equiv.), *p*-methoxybenzoic acid (1 equiv.), and CDI (1.3 equiv.) was stirred at reflux.



Scheme 5 Proposed reaction pathways.

second anion at C2 might attack acyl donors such as acyl imidazolides and esters. As with the unstable C8-ester (*i.e.* 3k) could be converted into the C2-amide by imidazole, the reactivity of the amino at C2 with acyl imidazolides under neutral conditions remains to be elucidated.

#### Conclusions

In conclusion, we developed chemoselective acylation of 2amino-8-quinolinol on amino group at C2 and hydroxy group at C8. The coupling reaction with a variety of carboxylic acids using EDCI and DMAP provided C8-ester derivatives in excellent yield. When the dianion of 2-amino-8-quinolinol reacts with less reactive acylimidazoles and esters, the corresponding C2amides and carbamate derivatives could be prepared as the major product. Biological evaluations of the C2-amide and C8ester derivatives of 2-amino-8-quinolinol are underway in our laboratory and will be reported in due course.

#### Experimental

All the starting materials, solvents and reagents were obtained from commercial suppliers and were used without further purification. Flash column chromatography was performed using silica gel 60 (230-400 mesh, Merck) with the indicated solvents. Thin-layer chromatography was performed using 0.25 mm silica gel plates (Merck). <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a Bruker 600 MHz spectrometer as solutions in deuterated CDCl<sub>3</sub>, methanol-d<sub>4</sub> or DMSO-d<sub>6</sub>. <sup>1</sup>H NMR data were reported in the order of chemical shift, multiplicity (s, singlet; d, doublet; t, triplet; m, multiplet and/or multiple resonances), number of protons, and coupling constant (1) in hertz (Hz). High-resolution mass spectra (HRMS) were recorded on a JEOL JMS-700 (FAB and EI) and an Agilent 6530 Q-TOF LC/MS/MS system (ESI). For conversion measurements, the samples prepared for the HPLC were analyzed at 256 nm using an Agilent 1260 HPLC system equipped with a 6 mm imes 50 mm Sunfire 5µ C18 column, in which the mobile phase was a gradient from water to acetonitrile for 30 min.

### General procedure of condition A (Table 4) for *O*-acylation at C8

To a solution of 2-amino-8-quinolinol (30 mg, 0.19 mmol) and a carboxylic acid (1.2 equiv.) in anhydrous tetrahydrofuran (4 mL) were added EDCI (1.3 equiv.), DMAP (0.5 equiv.) and *N*,*N*diisopropylethylamine (3.0 equiv.) under N<sub>2</sub> atmosphere with an ice bath. After stirring for 3 h at ambient temperature, the mixture was diluted with dichloromethane, washed by sat. NH<sub>4</sub>Cl, sat. NaHCO<sub>3</sub>, and brine, dried by MgSO<sub>4</sub>, filtered, and concentrated. The residue was purified by silica gel column chromatography to afford the desired product.

## General procedure of condition B (Table 4) for O-acylation at C8

To a solution of 2-amino-8-quinolinol (30 mg, 0.19 mmol) and a carboxylic acid (1.2 equiv.) in anhydrous tetrahydrofuran were added PyBop (1.3 equiv.) and N,N-diisopropylethylamine (3.0 equiv.) under N<sub>2</sub> atmosphere with an ice bath and stirring for 4 h at ambient temperature. The reaction mixture was diluted with dichloromethane, washed by NH<sub>4</sub>Cl, saturated NaHCO<sub>3</sub>, and brine, dried by MgSO<sub>4</sub>, filtered, and concentrated. The residue was purified by flash column chromatography to the desired product.

# General procedure of condition C (Table 5) for *N*-acylation at C2

To a solution of a carboxylic acid (1.2 equiv.) in anhydrous tetrahydrofuran was added 1,1'-carbodiimidazole (1.3 equiv.).

After stirring for 1 h under  $N_2$  atmosphere, 2-amino-8quinolinol (30 mg, 0.19 mmol) was added to the reaction mixture. After stirring for 24 h at reflux, the mixture was diluted with tetrahydrofuran, washed by NH<sub>4</sub>Cl, saturated NaHCO<sub>3</sub>, and brine, dried by MgSO<sub>4</sub>, filtered, and concentrated. The residue was purified by flash column chromatography to the desired product.

## General procedure of condition D (Table 5) for *N*-acylation at C2

To a solution of carboxylic acid (1.2 equiv.) in anhydrous tetrahydrofuran (4 mL), 1,1'-carbonyldiimidazole (1.3 equiv.) was added under N<sub>2</sub> atmosphere at ambient temperature. After stirred for 1 h. The mixture was diluted with dichloromethane, extracted by brine and dichloromethane, dried by MgSO<sub>4</sub>, filtered, and concentrated to obtain the acyl imidazolide intermediate. To a solution of 2-amino-8-quinolinol (30 mg, 0.19 mmol) and NaH (2.0 equiv.) in anhydrous tetrahydrofuran (4 mL) was added the acyl imidazolide intermediate under N<sub>2</sub> atmosphere with an ice bath. After stirring for 2 h at ambient temperature, the reaction was quenched with sat. NH<sub>4</sub>Cl, washed by dichloromethane and dried by MgSO<sub>4</sub>, filtered, and concentrated. The residue was purified by silica gel column chromatography to afford the desired product.

**2-Aminoquinolin-8-yl 4-methoxybenzoate** (3a). General procedure A, white solid, yield: 92%, and purification by column chromatography (ethyl acetate/*n*-hexane = 1 : 3,  $R_f = 0.1$ ). <sup>1</sup>H-NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.28 (dd, 2H, J = 9.0 Hz), 7.86 (d, 1H, J = 9.0 Hz), 7.53 (dd, 1H, J = 8.4 and 1.2 Hz), 7.39 (dd, 1H, J = 7.8 and 1.2 Hz), 7.26 (t, 1H, J = 7.8 Hz), 6.99 (dd, 2H, J = 9.0 and 1.8 Hz), 6.66 (d, 1H, J = 9.0 Hz), 4.85 (bs, 2H), 3.89 (s, 3H); <sup>13</sup>C-NMR (150 MHz, CDCl<sub>3</sub>)  $\delta$  165.3, 163.7, 157.0, 145.3, 140.7, 137.9, 132.5, 125.4, 125.0, 122.2, 122.1, 121.9, 113.7, 112.3, 55.5; HRMS (EI): m/z calcd for  $C_{17}H_{14}N_2O_3$ : 294.1004; found: 294.1006.

*N*-(8-Hydroxyquinolin-2-yl)-4-methoxybenzamide (4a). General procedure D, white solid, yield: 75%, and purification by column chromatography (dichloromethane/methanol =  $20 : 1, R_f = 0.25$ ). <sup>1</sup>H-NMR (600 MHz, DMSO-d<sub>6</sub>)  $\delta$  10.68 (s, 1H), 9.29 (bs, 1H), 8.34 (d, 1H, J = 9.0 Hz), 8.28 (d, 1H, J = 9.0 Hz), 8.10 (d, 2H, J = 8.4 Hz), 7.38 (d, 1H, J = 7.8 Hz), 7.35 (t, 1H, J = 7.8 Hz), 7.10 (t 3H, J = 7.5 Hz), 3.86 (s, 3H); <sup>13</sup>C-NMR (150 MHz, DMSO-d<sub>6</sub>)  $\delta$  165.9, 162.8, 152.4, 150.7, 138.4, 137.2, 130.5, 127.0, 126.3, 126.2, 118.2, 116.4, 114.1, 111.9, 55.9; HRMS (EI): m/z calcd for C<sub>17</sub>H<sub>14</sub>N<sub>2</sub>O<sub>3</sub>: 294.1004; found: 294.1003.

### Conflicts of interest

There are no conflicts to declare.

### Acknowledgements

This work was supported by grants from the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2016R1D1A1A09917300), the Pioneer Research Center Program through the NRF funded by the Ministry of Science, ICT & Future Planning (2014M3C1A3001556), and the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare (Grant number HI14C1135).

#### Notes and references

- For selected reviews of esterification and amidation, see: (a)
   F. Albericio, *Curr. Opin. Chem. Biol.*, 2004, **8**, 211–221; (b)
   S.-Y. Han and Y.-A. Kim, *Tetrahedron*, 2004, **60**, 2447–2467;
   (c) C. A. G. N. Montalbetti and V. Falque, *Tetrahedron*, 2005, **61**, 10827–10852; (d) R. M. de Figueiredo, J. S. Suppo and J. M. Campagne, *Chem. Rev.*, 2016, **116**, 12029–12122;
   (e) J. R. Dunetz, J. Magano and G. A. Weisenburger, *Org. Process Res. Dev.*, 2016, **20**, 140–177.
- 2 (a) R. Bright, D. J. Dale, P. J. Dunn, F. Hussain, Y. Kang, C. Mason, J. C. Mitchell and M. Snowden, Org. Process Res. Dev., 2004, 8, 1054–1058; (b) R. Vaidyanathan, V. G. Kalthod, D. P. Ngo, J. M. Manley and S. P. Lapekas, J. Org. Chem., 2004, 69, 2565–2568; (c) C. Larrivée-Aboussafy, B. P. Jones, K. E. Price, M. A. Hardink, R. W. McLaughlin, B. M. Lillie, J. M. Hawkins and R. Vaidyanathan, Org. Lett., 2010, 12, 324–327; (d) S. T. Heller, T. Fu and R. Sarpong, Org. Lett., 2012, 14, 1970–1973.
- 3 (a) J. P. Phillips, Chem. Rev., 1956, 56, 271–297; (b)
  M. Albrecht, M. Fiege and O. Osetska, Coord. Chem. Rev., 2008, 252, 812–824; (c) Y. N. Song, H. Xu, W. Chen, P. Zhan and X. Liu, Med. Chem. Commun., 2015, 6, 61–74.
- 4 (a) Y. Cheng, T. C. Judd, M. D. Bartberger, J. Brown, K. Chen, R. T. Fremeau, D. Hickman, S. A. Hitchcock, B. Jordan, V. Li, P. Lopez, S. W. Louie, Y. Luo, K. Michelsen, T. Nixey, T. S. Powers, C. Rattan, E. A. Sickmier, D. J. St Jean, R. C. Wahl, P. H. Wen and S. Wood, J. Med. Chem., 2011, 54, 5836–5857; (b) T. Tomioka, Y. Takahashi and T. Maejima, Org. Biomol. Chem., 2012, 10, 5113–5118; (c) M. A. Cinelli, H. Li, G. Chreifi, P. Martasek, L. J. Roman, T. L. Poulos and R. B. Silverman, J. Med. Chem., 2014, 57, 1513–1530; (d) L. Wang, J. Ferguson and F. Zeng, Org. Biomol. Chem., 2015, 13, 11486–11491.
- 5 (*a*) T. Storz, R. Marti, R. Meier, P. Nury, M. Roeder and K. Zhang, *Org. Process Res. Dev.*, 2004, **8**, 663–665; (*b*) P. K. Nayak, N. Agarwal, F. Ali, M. P. Patankar,

K. L. Narasimhan and N. Periasamy, *J. Chem. Sci.*, 2010, **122**, 847–855; (*c*) S. M. Shiriskar, N. Agarwal, R. R. Pissurlenkar and B. Ahmad, *Eur. Biophys. J.*, 2015, **44**, 193–205.

- 6 89% yield in *N*-acetylation. For details, see: (a) M. Albrecht,
  K. Witt, R. Frohlich and O. Kataeva, *Tetrahedron*, 2002, 58, 561–567. 32% yield in 8-hydroxyquinoline-2-carbonyl amide. For details, see (b) C. Deraeve, C. Boldron,
  A. Maraval, H. Mazarguil, H. Gornitzka, L. Vendier, M. Pitie and B. Meunier, *Chem.-Eur. J.*, 2008, 14, 682–696. 36% yield in ethyl malonyl amide. For details, see (c) K. Serafin,
  P. Mazur, A. Bak, E. Laine, L. Tchertanov, J.-F. Mouscadet and J. Polanski, *Bioorg. Med. Chem.*, 2011, 19, 5000–5005.
- 7 (a) I.-G. Je, H.-H. Kim, P.-H. Park, T. K. Kwon, S.-Y. Seo, T.-Y. Shin and S.-H. Kim, *Exp. Biol. Med.*, 2015, **240**, 631– 638; (b) S.-Y. Seo and W. Kang, *J. Pharm. Biomed. Anal.*, 2016, **131**, 103–106.
- 8 See ESI†, for details.
- 9 For chemoselective acylation, see: (a) M. Movassaghi and M. A. Schmidt, Org. Lett., 2005, 7, 2453–2456; (b) J. W. Bode, R. M. Fox and K. D. Baucom, Angew. Chem., Int. Ed., 2006, 45, 1248–1252; (c) S. D. Sarkar, S. Grimme and A. Studer, J. Am. Chem. Soc., 2010, 132, 1190–1191; (d) S. M. Mali, R. D. Bhaisare and H. N. Gopi, J. Org. Chem., 2013, 78, 5550–5555.
- 10 (a) Y.-J. Yoon, J. Park, B. Kim, H.-G. Lee, S.-B. Kang, G. Sung,
  J.-J. Kim and S.-G. Lee, *Synthesis*, 2012, 44, 42–50; (b)
  M. Choi, S.-W. Won, H. Jo, M. Viji, S.-Y. Seo, Y.-J. Lee,
  H.-S. Lee, H. Lee, J. T. Hong, Y.-S. Kwak and J.-K. Jung, *Tetrahedron Lett.*, 2014, 55, 6582–6584.
- 11 (a) L. Everson Da Silva, P. Teixieira De Sousa and A. C. Joussef, Synth. Commun., 2009, 39, 1378–1388; (b) M. M. Bader, T. Hamada and A. Kakuta, J. Am. Chem. Soc., 1992, 114, 6475–6479.
- 12 The  $pK_a$  values of 2-amino-8-quinolinol (1) and its conjugate acid are 10.5 and 6.7, respectively, calculated using the ACD/ Percepta 14 (ACD/Labs, Toronto, Canada). The chemical shift perturbation of the ring proton of N1 positon of 1 depending on pHs was monitored by NMR. The result also showed three plateaus exist around the calculated  $pK_a$ values, which supported that the  $pK_a$  values should be correct. For details, see ESI.<sup>†</sup>