# RSC Advances



# PAPER

Cite this: RSC Adv., 2017, 7, 51928

## Ag(I)/persulfate-catalyzed decarboxylative coupling of a-oxocarboxylates with organotrifluoroborates in water under room temperature†

Sheng Chang[,](http://orcid.org/0000-0001-5044-3334)  $\mathbb{D}^{*ab}$  Jian Feng Wang,<sup>c</sup> Lin Lin Dong,<sup>a</sup> Dan Wang,<sup>a</sup> Bo Feng<sup>\*a</sup> and Yuan Tai Shi<sup>a</sup>

The decarboxylative coupling reaction of  $\alpha$ -oxocarboxylates and organotrifluoroborates was carried out smoothly in the presence of catalytic AgNO<sub>3</sub> using  $K_2S_2O_8$  as oxidant to generate diarylketone products in high yields. The method is novel, simple, safe and efficient. Both aryl substituted potassium  $\alpha$ oxocarboxylates and organotrifluoroborates proceeded smoothly in water under room temperature. The utilization of  $\alpha$ -oxocarboxylates as acylating agent presents some elements of interest.

Received 3rd October 2017 Accepted 26th October 2017

DOI: 10.1039/c7ra10924e

rsc.li/rsc-advances

### Introduction

Arylketone is a common functionality in many types of organic compounds, for example, pharmaceuticals and natural products.<sup>1</sup> For organic chemists, it is also an extremely useful functional group for further manipulations. Consequently, it continues to inspire the development of methods for the construction of this structural element. Classical routes to synthesize aryl ketones mainly rely on the Friedel-Crafts acylation of aromatic compounds<sup>2</sup> and the oxidation of the corresponding secondary alcohols.<sup>3</sup> Transition-metal-catalyzed reactions provide many opportunities for the synthesis of aryl ketones. The coupling reaction of acylating agents with aryl halides (or aryl metallic compounds and their equivalents) has been developed under the catalysis of transition metals.<sup>4</sup>

Transition metal catalyzed decarboxylative coupling reactions have received much attention for their applications in the construction of C–C and C-hetero bonds in recent years.<sup>5</sup> Among various types of carboxylic acids (carboxylate), a-oxocarboxylates have emerged as a novel class of acyl surrogates after releasing molecular  $CO<sub>2</sub>$ .<sup>6</sup> Theoretically, there are three types of acyl reagents from the decarboxylation of a-oxocarboxylates: acyl anions, acyl cations and acyl radicals.<sup>7</sup> The direct decarboxylative acylation of potassium a-oxocarboxylates with aryl bromides/chlorides have been demonstrated by

Gooßen's group, with the  $Pd(n)/Cu(n)$  catalytic system being particularly efficient with the assistance of P-based and N-based ligands.<sup>8a-8c</sup> Then they demonstrated a one-pot threecomponent decarboxylative coupling with a-oxocarboxylates, amines and aryl bromides for the synthesis of azomethines.<sup>8d</sup> In 2014, Ji et al. reported a palladium/copper-catalyzed decarboxylative coupling of aryl iodides with a-oxocarboxylates without the need of ligands.<sup>8e</sup>  $\alpha$ -Oxocarboxylates providing acyl anions with Pd/Cu catalysis by this decarboxylation type, to afford diaryl ketones under harsh conditions (or derivatives) (Scheme 1, I). Recently, You and co-workers reported a transition-metal-free decarboxylation of a-oxocarboxylates with  $\alpha$ -bromo-acetophenone.<sup>9a</sup> In 2017, the room-temperature coupling/decarboxylation reaction between in situ generated a-oxocarboxylates with a-bromoketones was described by their group. The transformation shows excellent regioselectivity for 1,2- and 1,3-diketones by solvent-controlled.<sup>9b</sup> Acyl cations were generated in this decarboxylation manner, with the coupling of  $\alpha$ -oxocarboxylates with sp<sup>3</sup> carbon center (Scheme 1, II). Acyl radicals were formed via Ag(I)/persulfate-catalysis of potassium a-oxocarboxylates by the third decarboxylation type. Lei's group described the first example for realizing isocyanide insertion by using an acyl radical via the oxidative radical decarboxylation, to construct 6-acyl phenanthridine structure (Scheme 1, III).<sup>10a</sup> Synthesis of oxazoles by Ag(I)/persulfate-catalyzed oxidative decarboxylation–cyclization of a-oxocarboxylates and isocyanides was reported by them for the next year.<sup>10b</sup> Therefore, it is still necessary to develop more Ag(I)/persulfate-catalyzed oxidative decarboxylation of potassium a-oxocarboxylates. PAPER<br> **(A)** Check for goddes **Ag(I)/persulfate-catalyzed decarboxylative**<br>
Counter the section, 2017, 51926<br>
Coupling of  $\alpha$ -**oxocarboxylates with**<br> **or the section 2017**<br>
Coupling of  $\alpha$ -**oxocarboxylates with**<br> **or de** 

> Interestingly, there is a novel recognition that  $\alpha$ -oxocarboxylic acids can serve as aryl radical precursors via oxidative silverpromoted carbon–carbon bond cleavage in recent years.<sup>11</sup> Ag( $I$ )/ persulfate catalytic systems have been shown to be effective for

a College of Pharmacy, Jilin Medical University, Jilin, Jilin, 132013, China. E-mail: jmu\_changsheng@126.com; furfolgbec84@163.com; Tel: +86 432 6456 0532

b State Key Laboratory of Medicinal Chemical Biology, NanKai University, Tianjin, 300071, China

c Department of Radiotherapy, China-Japan Union Hospital of Jilin University, Changcun, Jilin, 130033, China

<sup>†</sup> Electronic supplementary information (ESI) available. See DOI: 10.1039/c7ra10924e



Scheme 1 Decarboxylation coupling of  $\alpha$ -oxocarboxylates.

this transformation, including direct C-H acylation of arenes<sup>12</sup> and oxidative radical decarboxylation–cyclization.<sup>13</sup> In the presence of silver $(i)$  salts, a persulfate anion disproportionates into a sulfate dianion and a sulfate radical anion, which could react with a-oxocarboxylates through an radical process, providing an acyl radical.

As continued effort on coupling of organoboronic reagents and carbonylative coupling,<sup>14</sup> we report herein a novel method for the silver-catalyzed, decarboxylative acylation of aryltri fluoroborates to form the corresponding diarylketones. It has been successfully developed using a-oxocarboxylates as acylating reagents, as well as a catalytic amount of a silver salt and a cheap inorganic oxidant. The reaction demonstrates a broad substrate scope and excellent functional-group tolerance. This catalytic method should be valuable in the synthesis of arylcarbonyl motifs.

### Results and discussion

Initially, we optimized the reaction conditions using potassium oxophenylacetate and phenyltrifluoroborate as model substrate in acetone with 5 mol% AgNO<sub>3</sub> and 1.5 equiv.  $K_2S_2O_8$  at room temperature for 1 h under air. Only trace amount of crosscoupling biphenyl ketone products was observed (Table 1, entry 1). Then we explored the silver-catalyzed reactions with a variety of reaction medium (Table 1), and 22% of yield were obtained when  $CH<sub>3</sub>CN$  was used as a solvent (Table 1, entry 2). No better results were obtained with polar aprotic solvents such as DMF, DMSO, or NMP(N-methyl-2-pyrrolidone) (Table 1, entries 3–5). The solvents (DCE, DCM, CHCl<sub>3</sub> and CCl<sub>4</sub>) were tested, and DCE gave the best result (Table 1, entries 6–9). Three additional ethereal solvents [1,4-dioxane, DME and THF] were surveyed without better results (Table 1, entries 10–12). However, the reaction did not proceed in toluene, EA or t-BuOH (Table 1, entries 13–15). Considered that the infusibility of both potassium  $\alpha$ -oxocarboxylates and aryltrifluoroborate in organic

solvent, water could be much better medium candidate for this transformation. Water is an easily available, nontoxic, nonin flammable, and renewable solvent. The unusual properties of water can lead to unusual reactivity that is not seen in traditional organic solvents. Consequently, the use of water as a solvent in synthetic organic chemistry and materials science has spread throughout the chemical community. The combination of water and metal catalysis has led in recent years to the development of a huge number of new and greener synthetic methodologies. It also provides opportunities for alternative reactivity and simplified product isolation compared to organic solvents. To our delight, sharply increase was observed with the application of water (Table 1, entry 16).

We began to screen many oxidants for the assistance of the regeneration of Ag(I). Many oxidants have been tested and showed different influence on the reaction (Table 2, entries 1-10). The cross-coupling could not proceed with  $Cu(OAc)<sub>2</sub>$ , oxygen and DDQ (Table 2, entries 2-4).  $PhI(OAc)_2$  and BPO (dibenzoyl peroxide) can enhance the efficiency to about 26–34% (Table 2, entries 5–6). In addition, inorganic peroxide are efficient to the cross-coupling. Oxone and  $(NH_4)_2S_2O_8$  can afford diphenylketone with the yield of 70% and 74% (Table 2, entries 7–8), then  $Na<sub>2</sub>S<sub>2</sub>O<sub>8</sub>$  performed better and  $K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>$  performed best (86%) (Table 2, entries 9–10). Then the effect of silver salts was investigated, and many silver( $I$ ) compounds such as AgOAc, Ag<sub>2</sub>CO<sub>3</sub>, and AgOTf were also applicable for this reaction, albeit in the slightly decreased yield (Table 2, entries 11-13). AgBF<sub>4</sub>, AgOTf and Ag2O were inferior to AgOTf (Table 2, entries 14–16). Meanwhile, the use of  $Ag_2SO_4$ ,  $AgSbF_6$  and AgF as catalysts gave no desired products (Table 2, entries 17–19). And 86% of yield were obtained when  $AgNO<sub>3</sub>$  was used as a catalyst (Table 2, entry 20). Therefore,  $AgNO<sub>3</sub>/K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>$  in water at room temperature under air for 1 hour was chosen as the optimal conditions for the synthesis of diarylketone from  $\alpha$ -oxocarboxylates and aryltrifluoroborate.

With optimal conditions in hand, the scope of this decarboxylative coupling reaction method was evaluated next (Tables

#### Table  $1$  Solvent effect of cross-coupling reaction<sup>a</sup>





<sup>a</sup> Reaction conditions: potassium oxophenylacetate (1.0 mmol), phenyltrifluoroborate (1.05 mmol), AgNO<sub>3</sub> (0.05 mmol), K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> (1.5 mmol), solvent  $(2 \text{ mL})$ ,  $25^{\circ}$ C, 1 h.  $^{b}$  Isolated yields.

#### Table 2 The screening of various oxidants and silver salts<sup>a</sup>





 $a$  Reaction conditions: potassium oxophenylacetate (1.0 mmol), phenyltrifluoroborate (1.05 mmol), silver salts (0.05 mmol), oxidant (1.5 mmol), H<sub>2</sub>O (2 mL), 25 °C, 1 h. <sup>b</sup> Isolated yields using AgNO<sub>3</sub> as the silver salt. <sup>c</sup> Isolated yields using K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> as the oxidant.

3 and 4). As shown, the reaction exhibits a broad substrate scope, with respect to both of the  $\alpha$ -oxocarboxylates and the organotrifluoroborates coupling partners. Importantly, the reaction is compatible with electron-neutral (3a), electron-rich (3b), and electron-deficient organotrifluoroborate precursors (3c–f), furnishing the products in good yields. The mild reaction conditions are compatible with a range of sensitive functional groups, such as nitro-(3c), trifluoromethyl-(3d), cyan-(3e), and even halides (fluoro-, bromo-, iodo-) (3f-h), affording diaryl ketone products with a bouquet of functional handles poised for further functionalization. Although, the organotrifluoroborate component bearing functional groups at different positions, such as methyl and chloro at 3- or 4-position of the phenyl ring were obtained in good yields (3i–l). Moreover, meta, para-di-methyl phenyltrifluoroborate and meta, para-dichloro phenyltrifluoroborate (3m and 3n) underwent the desired cross-coupling in high yields. Substrates having ortho methyl (30) or *ortho-chloro* (3p) substituents at phenyl moiety gave slightly lower reactivity, presumably due to steric reason. Finally, we were pleased to find that the reaction conditions could be readily extended to the arylation of a range of poly fluorinated substrates (3q-t). Importantly, polyfluorinated biaryl building blocks are of great interest in materials chemistry, because of favorable physicochemical properties.





<sup>a</sup> Reaction conditions: potassium oxophenylacetate (1.0 mmol), aryltrifluoroborate (1.05 mmol), AgNO<sub>3</sub> (0.05 mmol), K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> (1.5 mmol), H<sub>2</sub>O (2 mL), 25 °C, 1 h.  $<sup>b</sup>$  Isolated yields.</sup>

The scope, with respect to the  $\alpha$ -oxocarboxylates component, is also very broad (Table 4), including electron-rich (4a), and electron-deficient (4b–d). The reaction was shown to tolerate a wide variety of functionality on the aryl ring including methoxy-, cyan-, fluoro- and trifluoromethyl- (4a–d). Yields were reduced when using highly electron deficient arenes with substituents such as two  $CF_3$  (4e). The reaction could accommodate substrates containing a chloro, bromo or iodo substituent, which can later be applied to various cross-coupling reactions with metal catalysts (4f–h). Good yield was achieved using para-methyl and para-nitro substituents, to afford corresponding diarylmethanone (4i–j). Meta-substituted examples such as 3-methyl and 3-nitro (4k–l) also gave desired products in excellent yield. The ortho-substituted example (4m–n) were also tolerated however in reduced yield for sterically hindered siteselectivity. Furthermore, the formation of sterically demanding per-methylated and per-fluorinated product was smoothly in this process (4o-4p). The high selectivity of  $Ag(i)$ / persulfate-catalyzed system across a range of sterically and

electronically diverse a-oxocarboxylates and organotrifluoroborate precursors is a particularly noteworthy feature of this reaction manifold.

To further explore the potential of this decarboxylative coupling method, several  $(E)$ -styryl potassium trifluoroborate derivatives were examined (Scheme 2). (E)-styryl potassium tri fluoroborate underwent decarboxylative coupling with potassium 4-methoxy-oxophenylacetate in moderate yield. Decarboxylative coupling of  $(E)$ -styryl potassium trifluoroborate with potassium 4-chloro-oxophenylacetate also provided chalcone in good to excellent yields. It is worth noting that when the hydroxyl group was introduced to ortho-position of potassium oxophenylacetate, the decarboxylative coupling with vinyl potassium trifluoroborate gave acceptable conversion to afford chromone, showing the robustness of our protocol and demonstrating potential for practical applications. This synthetic method could further be extended to a large-scale decarboxylative acylation. The desired products (3a, 3i, 3k, 4f, 4i) were obtained with slightly decreased yields on 10 mmol



<sup>a</sup> Reaction conditions: *a*-oxocarboxylates (1.0 mmol), phenyltrifluoroborate (1.05 mmol), AgNO<sub>3</sub> (0.1 mmol), K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> (1.0 mmol), H<sub>2</sub>O (2 mL), 25 °C, 1 h.  $<sup>b</sup>$  Isolated yields.</sup>



Scheme 2 Applications of Ag(I)/persulfate-catalyzed decarboxylative coupling

scale (78% for 3a, 72% for 3i, 86% for 3k, 77% for 4f, and 81% for 4i).

With regard to our results and literature reports,<sup>10</sup> we propose the following possible reaction mechanism (Scheme 3). Monovalent silver catalyst was oxidized by  $K_2S_2O_8$  to persulfate anion radical and divalent silver ion (S3-1†) (the persulfate anion radical may also oxidize Ag(I) to Ag(II) (S3-2 $\dagger$ )). Then the Ag(II) generated in situ reacted with the  $\alpha$ -oxocarboxylates formed acyl radical complexes and  $CO<sub>2</sub>$  (S3-3†). Cross-coupling product obtained after the attack of acyl radical to

aryltrifluoroborate  $(S3-4\dagger)$ , which could be terminated by addition of radical scavengers such as hydroquinone or 2,6-di-tertbutylphenol, suggesting that the cross-coupling reactions proceed via a radical mechanism. Potassium trifluoroborate radical was finally quenched by oxidation of persulfate anion (S3-5†). Finally, 1.5 equiv. of TEMPO (2,2,6,6-tetramethyl-1 piperidinyloxy) was added to the reaction, and the acyl radical was trapped successfully by TEMPO, affording the corresponding adduct in 67% yields (Scheme 3, S3-6†).



Scheme 3 Possible mechanism.

### Conclusions

In conclusion, we have demonstrated a general method for highly selective decarboxylative coupling of  $\alpha$ -oxocarboxylates with aryltrifluoroborates enabled by versatile  $Ag(1)/persu$  Ifatecatalyzed system. The development of this coupling further highlights the beneficial use of inexpensive  $Ag(i)$  catalysts for decarboxylation, while exploiting  $K_2S_2O_8$  promoter. We anticipate that this strategy will find widespread use as a practical alternative to the classic ketone construction in organic synthesis. Most notably, this new protocol features the synthesis of chalcone and chromone derivatives. Further mechanistic studies and extension to a variety of coupling partners are currently underway in our laboratories.

## Conflicts of interest

There are no conflicts to declare.

### Acknowledgements

This work was supported by the State Key Laboratory of Medicinal Chemical Biology (2017001), National innovative undertaking plan of college students (201713743012), Young talent training plan Of Jilin Science and Technology Bureau (20156429), and NSFC (81773692).

### References

1 (a) H. Surburg and J. Panten, Common Fragrance and Flavor Materials, Wiley, Weinheim, Germany, 2006; (b) Y. Deng, Y.-W. Chin, H. Chai, W. J. Keller and A. D. Kinghorn, J. Nat. Prod., 2007, 70, 2049; (c) K. R. Romins, G. A. Freeman, L. T. Schaller, J. R. Cowan, S. S. Gonzales, J. H. Tidwell, C. W. Andrews, D. K. Stammers, R. J. Hazen, R. G. Ferris, S. A. Short, J. H. Chan and L. R. Boone, J. Med. Chem., 2006, 49, 727; (d) B. M. O'Keefe, N. Simmons and S. F. Martin, Org. Lett., 2008, 10, 5301; (e) S. Rahimipour, C. Palivan, F. Barbosa, I. Bilkis, Y. Koch, L. Weiner,

M. Fridkin, Y. Mazur and G. Gescheidt, J. Am. Chem. Soc., 2003, 125, 1376; (f) D. Ma, Y. Jiang, F. Chen, L.-K. Gong, K. Ding, Y. Xu, R. Wang, A. Ge, J. Ren, J. Li, J. Li and Q. Ye, J. Med. Chem., 2006, 49, 456; (g) Y. Nishizuka, Nature, 1988, 334, 661; (h) M. R. Hansen and L. H. Hurley, Acc. Chem. Res., 1996, 29, 249.

- 2 (a) G. Sartori and R. Maggi, Chem. Rev., 2006, 106, 1077; (b) G. A. Olah, Friedel-Crafts Chemistry., Wiley, New York, 1973.
- 3 (a) J. E. Bäckvall, Modern Oxidation Methods., Wiley-VCH, Weinheim, 2010; (b) G. E. Dobereiner and R. H. Crabtree, Chem. Rev., 2010, 110, 681.
- 4 For reviews see: (a) C. Pan, X. Jia and J. Cheng, Synthesis, 2012, 44, 677; (b) X.-F. Wu, H. Neumann and M. Beller, Chem. Soc. Rev., 2011, 40, 4986; (c) J. Ruan and J. Xiao, Acc. Chem. Res., 2011, 44, 614; (d) L. Kollár, Modern Carbonylation Methods., Wiley, Weinheim, 2008; (e) A. Brennfuhrer, H. Neumann and M. Beller, Angew. Chem., Int. Ed., 2009, 48, 4114; (f) P. Anbarasan, T. Schareina and M. Beller, Chem. Soc. Rev., 2011, 40, 5049.
- 5 (a) P. Hu, M. Zhang, X. Jie and W. Su, Angew. Chem., Int. Ed., 2012, 51, 227; (b) P. Hu, Y. Shang and W. Su, Angew. Chem., Int. Ed., 2012, 51, 5945; (c) J. Cornella and I. Larrosa, Synthesis, 2012, 653; (d) J. D. Weaver, A. Recio, A. J. Grenning and J. A. Tunge, Chem. Rev., 2011, 111, 1846; (e) O. Baudoin, Angew. Chem., Int. Ed., 2007, 46, 1373; (f) N. Rodriguez and L. J. Goossen, Chem. Soc. Rev., 2011, 40, 5030; (g) T. Satoh and M. Miura, Synthesis, 2010, 3395; (h) L. J. Goosen, N. Rodríguez and K. Goosen, Angew. Chem., Int. Ed., 2008, 47, 3100; (i) D.-X. Liu, F.-L. Li, H.-X. Li, J. Gao and J.-P. Lang, Tetrahedron, 2014, 70, 2416; (j) W.-J. Gong, D.-X. Liu, F.-L. Li, J. Gao, H.-X. Li and J.-P. Lang, Tetrahedron, 2015, 71, 1269; (k) D.-X. Liu, F.-L. Li, H.-X. Li, W.-J. Gong, J. Gao and J.-P. Lang, Eur. J. Org. Chem., 2014, 4817; (l) R.-T. He, J.-F. Wang, H.-F. Wang, Z.-G. Ren and J.-P. Lang, Dalton Trans., 2014, 43, 9786. Open Access Article. Published on 08 November 2017. Downloaded on 6/9/2025 8:16:03 AM. This article is licensed under a [Creative Commons Attribution-NonCommercial 3.0 Unported Licence.](http://creativecommons.org/licenses/by-nc/3.0/) **[View Article Online](https://doi.org/10.1039/c7ra10924e)**
	- 6 (a) Q. Jiang, J. Jia, B. Xu, A. Zhao and C.-C. Guo, J. Org. Chem., 2015, 80, 3586; (b) N. Zhang, D. Yang, W. Wei, L. Yuan, F. Nie, L. Tian and H. Wang, J. Org. Chem., 2015, 80, 3258; (c) W.-P. Mai, G.-C. Sun, J.-T. Wang, G. Song, P. Mao, L.-R. Yang, J.-W. Yuan, Y.-M. Xiao and L.-B. Qu, J. Org. Chem., 2014, 79, 8094; (d) K. Yan, D. Yang, W. Wei, F. Wang, Y. Shuai, Q. Li and H. Wang, J. Org. Chem., 2015, 80, 1550.
	- 7 C. Chatgilialoglu, D. Crich, M. Komatsu and I. Ryu, Chem. Rev., 1999, 99, 1991.
	- 8 (a) L. J. Gooßen, F. Rudolphi, C. Oppel and N. Rodríguez, Angew. Chem., Int. Ed., 2008, 47, 3043; (b) L. J. Gooßen, B. Zimmermann, C. Linder, N. Rodríguez, P. P. Lange and J. Hartung, Adv. Synth. Catal., 2009, 351, 2667; (c) L. J. Gooßen, B. Zimmermann and T. Knauber, Angew. Chem., Int. Ed., 2008, 47, 7103; (d) F. Rudolphi, B. Song and L. J. Gooßen, Adv. Synth. Catal., 2011, 353, 337; (e) Y. Ji, X. Yang and W. Mao, Appl. Organomet. Chem., 2014, 28, 678.
	- 9 (a) Z. He, X. Qi, S. Li, Y. Zhao, G. Gao, Y. Lan, Y. Wu, J. Lan and J. You, Angew. Chem., Int. Ed., 2015, 54, 855; (b) Z. He,

X. Qi, Z. She, Y. Zhao, S. Li, J. Tang, G. Gao, Y. Lan and J. You, J. Org. Chem., 2017, 82, 1403.

- 10 (a) J. Liu, C. Fan, H. Yin, C. Qin, G. Zhang, X. Zhang, H. Yi and A. Lei, Chem. Commun., 2014, 50, 2145; (b) Y. Ma, Z. Yan, C. Bian, K. Li, X. Zhang, M. Wang, X. Gao, H. Zhang and A. Lei, Chem. Commun., 2015, 51, 10524.
- 11 (a) K. Cheng, B. Zhao and C. Qi, RSC Adv., 2014, 4, 48698; (b) N. Zhang, D. Yang, W. Wei, L. Yuan, F. Nie, L. Tian and H. Wang, J. Org. Chem., 2015, 80, 3258; (c) L.-N. Guo, H. Wang and X.-H. Duan, Org. Biomol. Chem., 2016, 14, 7380; (d) S. Zhang, L.-N. Guo, H. Wang and X.-H. Duan, Org. Biomol. Chem., 2013, 11, 4308; (e) H. Wang, L.-N. Guo and X.-H. Duan, Org. Biomol. Chem., 2013, 11, 4573; (f) H. Wang, H. Yang, Y. Li and X.-H. Duan, RSC Adv., 2014, 4, 8720; (g) H. Wang, L.-N. Guo and X.-H. Duan, Org. Lett., 2012, 14, 4358. NSC Advances Wewell-Comes Articles. Published on 2017. Downloaded on 1972. The Common Access Articles. Published on 6. Zhang, 2. Zha
	- 12 (a) W. P. Mai, B. Sun, L. Q. You, L. R. Yang, P. Mao, J. W. Yuan, Y. M. Xiao and L. B. Qu, Org. Biomol. Chem.,

2015, 13, 2750; (b) R. Suresh, R. S. Kumaran, V. Senthilkumar and S. Muthusubramanian, RSC Adv., 2014, 4, 31685; (c) H. Wang, S. L. Zhou, L. N. Guo and X. H. Duan, Tetrahedron, 2015, 71, 630.

- 13 (a) Z. Y. Lin, Y. L. Chen, C. S. Lee and C.-P. Chuang, Eur. J. Org. Chem., 2010, 3876; (b) T. Liu, Q. Ding, Q. Zong and G. Qiu, Org. Chem. Front., 2015, 2, 670; (c) H. Wang, L. N. Guo and X. H. Duan, Adv. Synth. Catal., 2013, 355, 2222; (d) H. Yang, L. N. Guo and X. H. Duan, RSC Adv., 2014, 4, 52986; (e) W. P. Mai, G. C. Sun, J. T. Wang, G. Song, P. Mao, L. R. Yang, J. W. Yuan, Y. M. Xiao and L. B. Qu, J. Org. Chem., 2014, 79, 8094.
- 14 (a) C. Y. Hao, D. Wang, Y. W. Li, L. L. Dong, Y. Jin, X. R. Zhang, H. Y. Zhu and S. Chang, RSC Adv., 2016, 6, 86502; (b) S. Chang, Y. Jin, X. R. Zhang and Y. B. Sun, Tetrahedron Lett., 2016, 57, 2017; (c) S. Chang, Y. B. Sun, X. R. Zhang, L. L. Dong, H. Y. Zhu, H. W Lai and D. Wang, Appl. Organomet. Chem., 2017, DOI: 10.1002/aoc.3970.