



Cite this: *RSC Adv.*, 2017, 7, 29925

Correction: Microwave-assisted synthesis of polypyridyl ruthenium(II) complexes as potential tumor-targeting inhibitors against the migration and invasion of Hela cells through G2/M phase arrest

Jieqiong Cao,^a Qiong Wu,^b Wenjie Zheng,^{*abd} Li Li^c and Wenjie Mei^{*c}

DOI: 10.1039/c7ra90067h

www.rsc.org/advances

Correction for 'Microwave-assisted synthesis of polypyridyl ruthenium(II) complexes as potential tumor-targeting inhibitors against the migration and invasion of Hela cells through G2/M phase arrest' by Jieqiong Cao *et al.*, *RSC Adv.*, 2017, 7, 26625–26632.

The authors regret that Fig. 1 is incorrect in the original manuscript. The transwell result of 5 μM was the same image as the image used for 10 μM . The correct figure is displayed below.




Fig. 1 (A) The wound healing assay of Hela cells after treatment with 4 (0, 2, 4 and 8 mM) and [NAMI-A] = 10 mM. (B) The healing rate of Hela cells treated with 4 and NAMI-A. (C) The transwell assay of Hela cells after treatment with 4 (0, 2, 4 and 8 mM) and (D) [NAMI-A] = (0, 5, 10 and 20 mM).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aCollege of Pharmacy, Jinan University, Guangzhou, China. E-mail: tzhwj@jnu.edu.cn

^bIntegrated Chinese and Western Medicine Postdoctoral Research Station, Jinan University, Guangzhou, China

^cSchool of Pharmacy, Guangdong Pharmaceutical University, Guangzhou, China. E-mail: wenjiemei@126.com

^dDepartment of Chemistry, Jinan University, Guangzhou, China

