CrossMark
click for updatesCite this: *Chem. Sci.*, 2017, 8, 616Received 17th June 2016
Accepted 1st September 2016

DOI: 10.1039/c6sc02660e

www.rsc.org/chemicalscience

Highly selective olefin-assisted palladium-catalyzed oxidative carbocyclization *via* remote olefin insertion†

Youai Qiu, Bin Yang, Can Zhu and Jan-E. Bäckvall*

A highly selective olefin-assisted palladium-catalyzed oxidative carbocyclization *via* remote olefin insertion to afford cyclohexenes has been developed. It was shown that the assisting olefin moiety was indispensable for the formation of the cyclohexene product. Furthermore, preliminary studies on chiral anion-induced asymmetrical carbocyclization–borylation of enallenes have been carried out.

The development of modern methodologies for the efficient synthesis of carbocycles is of central importance in modern organic chemistry,¹ given the fact that carbocycles are basic units in pharmacologically active skeletons, as well as in natural products.² Among these methodologies, transition metal-catalyzed carbocyclizations with the involvement of π -bonds have emerged as an effective strategy for the preparation of carbocycles,^{3,4} considering that the π -bond moiety would not only perform as the assisting group for the formation of the carbon–metal (C–M) bond, but also as the building block for the subsequent carbocyclization. In this way, an atom-economical transformation can be achieved.

We recently reported on an olefin-directed palladium-catalyzed oxidative carbocyclization–borylation of allenes to cyclobutenes (Scheme 1a).⁵ In this reaction, the coordination of the olefin in

Int-A triggers the allene attack on palladium,⁶ which results in the formation of *Int-B*. Subsequent olefin insertion to form a cyclobutene intermediate *Int-C*, followed by transmetalation and reductive elimination afforded the borylated cyclobutene derivatives **A**.

On the basis of these observations, we were particularly interested in the involvement of an additional double bond in a carbocyclization (Scheme 1b). We envisioned that the olefin insertion of intermediate *Int-1* could lead to intermediate *Int-2*, and that coordination of the additional olefin to palladium would lead to a second carbocyclization to form spirocyclic intermediate *Int-3*, which on reaction with B_2pin_2 would give **B**. Alternatively, *Int-1* may undergo ligand exchange and olefin insertion to give *Int-5*, which can be quenched by either B_2pin_2 or $ArB(OH)_2$ to give either **2** or **3**, respectively (Scheme 1b).

Based on this concept, we initially chose a readily accessible 3,4-dienoate **1a** as the standard substrate. When **1a** bearing an extra olefin was treated with $Pd(OAc)_2$ (5 mol%), B_2pin_2 (1.3 equiv.), and BQ (*p*-benzoquinone) (1.1 equiv.) in THF at room temperature for 12 h, the envisioned spirocyclic product **B** was not observed (Scheme 2). Interestingly, the cyclohexene product **2a** was obtained in 76% yield instead. It is obvious that intermediate *Int-2* was not formed from intermediate *Int-1* *via* olefin insertion as we had envisioned, but rather the ligand exchange, from proximal olefin to remote olefin occurred in *Int-1* to produce *Int-4*. Subsequent olefin insertion⁷ to give cyclic intermediate *Int-5* followed by B_2pin_2 quenching would produce **2a** (Scheme 1b). During this transformation, the

a) Previous work:

Scheme 1 Previous reports and this approach.

Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, SE-106 91, Stockholm, Sweden. E-mail: jeb@organ.su.se

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c6sc02660e

Scheme 2 Palladium-catalyzed oxidative carbocyclization–borylation of enallene **1a**.

proximal olefin is acting as the assisting group for the generation of palladium intermediate **Int-1**, while the remote olefin participates in the carbocyclization (Scheme 1c). To the best of our knowledge,^{8–10} the formation of six-membered rings in palladium-catalyzed oxidative carbocyclization of enallenes *via* olefin exchange has been rarely reported.¹¹

To demonstrate the necessity of the assisting olefin group, we first investigated comparative experiments with enallenes lacking the additional olefin (Scheme 3). When substrate **1f** with an assisting olefin was allowed to react under the same reaction conditions as those in Scheme 2, the cyclohexene product **2f** was formed in 70% yield (Scheme 3a). However, when substrate **1fb**, lacking the additional double bond, was subjected to the reaction conditions of Scheme 2 the corresponding six-membered ring product **2fb** was not formed; instead **1fb** was recovered in 88% (Scheme 3b). Importantly, we also observed the exclusive formation of **2h** in 68% yield from substrate **1h** (Scheme 3c). We also examined the reaction of a malonate-tethered substrate **1i**, but the envisioned product **2i** was not observed (Scheme 3d). These comparative experiments indicate that the assisting olefin of the substrate is an indispensable group for the formation of the palladium intermediate **Int-4**.

With these inspiring results in hand, we began to optimize the reaction conditions (for details, see the ESI, Table S1†). Solvent screening showed that 1,2-dichloroethane was the best solvent, in which the yield of **2a** was 91% (Table S1,† entry 7). Other solvents such as THF, MeCN, and 1,4-dioxane also gave good yields (Table S1,† entries 1, 5, and 6). When the amount of BQ was increased to 1.5 equivalents, the yield of **2a** was decreased to 78% (Table S1,† entry 8). The use of 2,6-dimethyl-BQ instead of BQ, gave a lower yield (Table S1,† entry 9). Catalyst screening showed that Pd(TFA)₂ (TFA = trifluoroacetate) produced the corresponding cyclohexene derivative in only 36% yield together with 34% starting material recovered (Table S1,† entry 10).

The substrate scope for the formation of cyclohexene boron compounds **2**^{12,13} was then studied under the optimized reaction conditions (Scheme 4): in addition to methyl substituents on the

Scheme 4 Scope of palladium-catalyzed oxidative borylating carbocyclization. The reaction was conducted in DCE at room temperature using **1** (0.2 mmol), B₂pin₂ (1.3 equiv.), and BQ (1.1 equiv.) in the presence of Pd(OAc)₂ (5 mol%).

enallene moiety, cyclobutylidene, cyclopentylidene, and cyclohexylidene enallenes (**1b**, **1c**, and **1d**) also gave the corresponding products **2b**, **2c**, and **2d** in good yields. To our delight, enallenes with functional groups, such as free hydroxyl in **1e** and imide in **1k**, furnished cyclohexene derivatives **2e** and **2k** in 83% and 76% yield. Furthermore, the reaction tolerates R to be different alkyl groups in this reaction, *e.g.* *n*-butyl (**1f**), or benzyl (**1g**).¹⁴ It is worth noting that the product **2h** was exclusively obtained in 84% yield. Finally, the reaction of a dissymmetric allene **1i**, bearing Me and phenyl, or **1j**, bearing Me and *i*-Pr, afforded **2i** in 86% yield, and **2j** and **2j'** in 60% yield, respectively. The ratio of **2j** and **2j'** was 1 : 2 due to the selective C–H bond cleavage, which occurred during allene attack forming **Int-1** (see Scheme 1b). Notably, the reaction could be easily extended to a scale of 4.5 mmol of **1a** (1.053 g) to afford the corresponding cyclohexene compound **2a** (1.551 g, 90% yield).

After realization of the borylative carbocyclization, we next turned our attention to the arylating carbocyclization of enallenes. We were pleased to find that the arylative products¹⁵ could be obtained in good yields with a catalyst loading of 2 mol% Pd(OAc)₂ (Scheme 5). The reaction of substrates with two methyls, cyclopentylidene, and cyclohexylidene, afforded the corresponding product **3a**, **3c**, and **3d** in good yields. Interestingly, the substrate containing a free hydroxyl group could also be employed. Different alkyl substituents on the starting materials, such as *n*-butyl, benzyl, and 4-pentenyl groups were tolerated (**3f–h**). We also examined the scope of arylboronic acids, and electron-donating substituents such as 3-Me, and 3-MeO reacted smoothly under the standard conditions in good yields. Notably, the procedure tolerates a range of additional functional groups on the arylboronic acid, including bromo (**3ad**), vinyl (**3ae**), formyl (**3af**), and acetyl (**3ag**) groups, which is useful for further functionalization. Finally, it is worth noting that 2-naphthylboronic acid and 1-naphthylboronic acid

Scheme 3 Comparative experiments.

Scheme 5 Scope of the palladium-catalyzed oxidative arylating carbocyclization. ^a The reaction was conducted in DCE at room temperature using **1** (0.2 mmol), $\text{ArB}(\text{OH})_2$ (1.3 equiv.), and BQ (1.1 equiv.) in the presence of $\text{Pd}(\text{OAc})_2$ (2 mol%). ^b The reaction was conducted at 50°C .

also worked well, affording **3ah** and **3ai** in 84% and 55% yield, respectively.

Interestingly, this new olefin-assisting strategy could also be applied to an oxidative carbonylating carbocyclization^{9c,f,g} for the preparation of cyclohexene esters (Scheme 6).¹⁶ When the substrate **1a** was treated with $\text{Pd}(\text{OAc})_2$ (2 mol%), and BQ (1.1 equiv.) under carbon monoxide (1 atm) in methanol at room temperature for 12 h the carbonylation product **4a** was formed in

Scheme 6 Scope of palladium-catalyzed oxidative carbonylating carbocyclization. The reaction was conducted in $\text{R}'\text{OH}$ using **1** (0.2 mmol), DMSO (20 mol%), and BQ (1.1 equiv.) in the presence of $\text{Pd}(\text{OAc})_2$ (2 mol%).

82% yield (Scheme 6). Under the optimal reaction conditions, cyclopentylidene and cyclohexylidene substrates **1c** and **1d** afforded **4c** and **4d**, respectively in good yields. The substrate bearing the free hydroxyl group (**1e**) also worked well. Different allenes with alkyl substituents such as *n*-butyl, benzyl, and 4-pentenyl group, were also tested and worked well in this reaction. The reaction of a dissymmetric allene **1i**, bearing Me and phenyl, afforded **4i** in 70% yield. Finally, the scope of the alcohol partners in the carbocyclization carbonylation reaction was explored, and in addition to MeOH, ethanol and isopropanol were shown to react smoothly to provide the desired esters in good yield.

The biomimetic approach with the use of electron-transfer mediators (ETMs) is known to decrease the kinetic barrier for the reoxidation.¹⁷ In this aerobic approach the high kinetic barrier will be divided into several smaller units, and catalytic amounts of oxidant (BQ) would be enough to realize these transformations. When the reaction of **1a** was treated with B_2pin_2 (1.3 equiv.), BQ (20 mol%), $\text{Pd}(\text{OAc})_2$ (5 mol%), and cobalt(salophen) (5 mol%) in the presence of O_2 (1 atm), borylated product **2a** was obtained in 89% yield. Phenylated product **3a** was provided in 86% yield when $\text{PhB}(\text{OH})_2$ was used in place of B_2pin_2 (Scheme 7).

Preliminary attempt to develop an enantioselective carbocyclization–borylation of enallenes revealed that a reasonably good *er* value (83 : 17) was observed in the presence of catalytic amounts of $\text{Pd}(\text{OAc})_2$ and biphenol-type phosphoric acid **C**^{9d} while poor enantiocontrol (55 : 45 *er*) was obtained with VAPOL phosphoric acid (Scheme 8).^{18,19}

Based on the experiments in Scheme 3 and the reaction outcome, a possible mechanism for the olefin-assisted palladium-catalyzed oxidative carbocyclization of enallenes *via* remote olefin insertion is given in Scheme 9. The reaction of palladium with enallene **1** bearing the assisting olefin forms vinylpalladium intermediate **Int-1** *via* allene attack involving allenic C–H bond cleavage, which is promoted by the coordination of allene and the assisting olefin to $\text{Pd}(\text{II})$.^{5,6} Then, the vinylpalladium intermediate **Int-4** would be generated from **Int-1** *via* ligand exchange (from proximal olefin to remote olefin), instead of a direct olefin insertion to form cyclobutene complex **Int-2**.⁵ Intermediate **Int-4** would undergo a remote olefin insertion to give cyclic intermediate **Int-5**. Subsequent transmetalation of **Int-5** with B_2pin_2 or arylboronic acid, followed by reductive elimination would give the target cyclohexene derivatives **2** or **3**. Under CO pressure in alcohol, **Int-5** can undergo an alkoxy-carbonylation to provide product **4**.

Scheme 7 Palladium-catalyzed oxidative carbocyclization under biomimetic conditions.

In conclusion, we have developed a highly selective olefin-assisted palladium-catalyzed oxidative carbocyclization of enallenes *via* remote olefin insertion for the selective formation of the cyclohexene skeleton. It was demonstrated that the assisting olefin moiety is essential for the formation of the cyclohexene derivatives. These reactions all show a broad substrate scope and good tolerance for various functional groups, and the catalyst loading could be decreased to 2 mol% in the arylation and carbonylation reactions with good to excellent yields. The biphenol-type chiral phosphoric acid was used in preliminary experiments of enantioselective carbocyclization–borylation of enallene. Further studies on the scope, synthetic application, and asymmetric variants of these reactions are currently carried out in our laboratory.

Financial support from the European Research Council (ERC AdG 247014), The Swedish Research Council (621-2013-4653), The Berzelii Center EXSELENT, and the Knut and Alice Wallenberg Foundation is gratefully acknowledged.

1 For selected reviews involving carbocyclization reaction, see:
(a) E. I. Negishi, C. Copéret, S. Ma, S. Y. Liou and F. Liu,
Chem. Rev., 1996, **96**, 365; (b) M. Méndez and
A. M. Echavarren, *Eur. J. Org. Chem.*, 2002, 15; (c)

B. M. Trost, F. D. Toste and A. B. Pinkerton, *Chem. Rev.*, 2001, **101**, 2067; (d) C. Aubert, L. Fensterbank, V. Gandon and M. Malacria, *Top. Organomet. Chem.*, 2006, **19**, 259; (e) D. H. Zhang, Z. Zhang and M. Shi, *Chem. Commun.*, 2012, **48**, 10271; (f) C. Aubert, L. Fensterbank, P. Garcia, M. Malacria and A. Simonneau, *Chem. Rev.*, 2011, **111**, 1954; (g) R. I. McDonald, G. Liu and S. S. Stahl, *Chem. Rev.*, 2011, **111**, 2981.

- Chem. Sci., 2017, 8, 616–620 | 619

- Lett.*, 2006, **8**, 3801; (b) M. A. Tarselli, A. R. Chianese, S. J. Lee and M. R. Gagné, *Angew. Chem., Int. Ed.*, 2007, **46**, 6670; (c) F. Grau, A. Heumann and E. Duñach, *Angew. Chem., Int. Ed.*, 2006, **45**, 7285; (d) Y. Qiu, J. Zhou, J. Li, C. Fu, Y. Guo, H. Wang and S. Ma, *Chem.-Eur. J.*, 2015, **21**, 15939; (e) J. W. Park, Z. Chen and V. M. Dong, *J. Am. Chem. Soc.*, 2016, **138**, 3310.
- 11 Pd-catalyzed oxidative carbocyclization of bisallenes to cyclohexene derivatives have been realized in our laboratory: C. M. R. Volla and J.-E. Bäckvall, *ACS Catal.*, 2016, **6**, 6398.
- 12 For selected reviews involving the organoboronates, see: (a) J. Tsuji, *Palladium Reagents and Catalysts*, John Wiley & Sons, Chichester, England, 2004, pp. 289–310; (b) D. S. Matteson, *Chem. Rev.*, 1989, **89**, 1535; (c) D. S. Matteson, *Acc. Chem. Res.*, 1988, **21**, 294; (d) G. C. Fu, *Acc. Chem. Res.*, 2008, **41**, 1555.
- 13 (a) H. C. Brown, R. Liotta and G. W. Kramer, *J. Am. Chem. Soc.*, 1979, **101**, 2966; (b) J. Kister, A. C. DeBaillie, R. Lira and W. R. Roush, *J. Am. Chem. Soc.*, 2009, **131**, 14174; (c) W. Yuan and S. Ma, *Adv. Synth. Catal.*, 2012, **354**, 1867; (d) S. B. Thorpe, X. Guo and W. L. Santos, *Chem. Commun.*, 2011, **47**, 424; (e) F. Meng, B. Jung, F. Haeffner and A. H. Hoveyda, *Org. Lett.*, 2013, **15**, 1414; (f) B. Jung and A. H. Hoveyda, *J. Am. Chem. Soc.*, 2012, **134**, 1490; (g) K. Semba, T. Fujihara, J. Terao and Y. Tsuji, *Angew. Chem., Int. Ed.*, 2013, **52**, 12400; (h) K. Semba, N. Bessho, T. Fujihara, J. Terao and Y. Tsuji, *Angew. Chem., Int. Ed.*, 2014, **53**, 9007; (i) C. Zhu, B. Yang, Y. Qiu and J.-E. Bäckvall, *Chem.-Eur. J.*, 2016, **22**, 2939.
- 14 Allenes **1f**, **1g**, and **1h** were prepared by iron-catalyzed cross coupling of propargyl acetate **5** and the appropriate Grignard reagent: S. N. Kessler and J.-E. Bäckvall, *Angew. Chem., Int. Ed.*, 2016, **55**, 3734.
- 15 (a) Y. Deng, T. Bartholomeyzik, A. K. Å. Persson, J. Sun and J.-E. Bäckvall, *Angew. Chem., Int. Ed.*, 2012, **51**, 2703; (b) T. Bartholomeyzik, J. Mazuela, R. Pendrill, Y. Deng and J.-E. Bäckvall, *Angew. Chem., Int. Ed.*, 2014, **53**, 8696.
- 16 For selected reviews on palladium catalyzed carbonylation, see: (a) T. Morimoto and K. Kakiuchi, *Angew. Chem., Int. Ed.*, 2004, **43**, 5580; (b) A. Brennfürer, H. NeuMann and M. Beller, *Angew. Chem., Int. Ed.*, 2009, **48**, 4114; (c) R. Grigg and S. P. Mutton, *Tetrahedron*, 2010, **66**, 5515; (d) X. F. Wu, H. Neumann and M. Beller, *Chem. Rev.*, 2013, **113**, 1.
- 17 (a) J. Piera and J.-E. Bäckvall, *Angew. Chem., Int. Ed.*, 2008, **47**, 3506; (b) J.-E. Bäckvall, R. B. Hopkins, H. Grennberg, M. Mader and A. K. Awasthi, *J. Am. Chem. Soc.*, 1990, **112**, 5160; (c) C. M. R. Volla and J.-E. Bäckvall, *Angew. Chem., Int. Ed.*, 2013, **52**, 14209.
- 18 For selected examples, see: (a) G. Jiang and B. List, *Angew. Chem., Int. Ed.*, 2011, **50**, 9471; (b) K. Ohmatsu, N. Imagawa and T. Ooi, *Nat. Chem.*, 2014, **6**, 47; (c) Z. Chai and T. Rainey, *J. Am. Chem. Soc.*, 2012, **134**, 3615; (d) P. Wang, H. Lin, Y. Zhai, Z. Han and L. Gong, *Angew. Chem., Int. Ed.*, 2014, **53**, 12218.
- 19 For selected reviews, see: (a) M. Mahlau and B. List, *Angew. Chem., Int. Ed.*, 2013, **52**, 518; (b) R. J. Phipps, G. L. Hamilton and F. D. Toste, *Nat. Chem.*, 2012, **4**, 603; (c) F. Parmar, E. Sugiono, S. Raja and M. Rueping, *Chem. Rev.*, 2014, **114**, 9047; (d) D. Chen, Z. Han, X. Zhou and L. Gong, *Acc. Chem. Res.*, 2014, **47**, 2365.