# Chemical Science



## **EDGE ARTICLE**

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2017, 8, 2457

Received 6th October 2016 Accepted 26th December 2016

DOI: 10.1039/c6sc04459j

www.rsc.org/chemicalscience

# CO/CO and NO/NO coupling at a hidden frustrated Lewis pair template†

Tongdao Wang,<sup>a</sup> Long Wang,<sup>a</sup> Constantin G. Daniliuc,<sup>a</sup> Kamil Samigullin,<sup>b</sup> Matthias Wagner,<sup>b</sup> Gerald Kehr<sup>a</sup> and Gerhard Erker<sup>\*a</sup>


N-Allyltetramethylpiperidine is readily isomerized to the corresponding enamine by treatment with catalytic amounts of  $B(C_6F_5)_3$ . It adds  $HB(C_6F_5)_2$  at the nucleophilic enamine carbon atom to form a C/B Lewis adduct. This reacts with two molar equivalents of carbon monoxide by selective head to tail coupling to give a five-membered  $C_2O_2B$  heterocycle. In contrast the enamine/ $HB(C_6F_5)_2$  Lewis pair reacts with two molar equiv. of nitric oxide by head to head coupling. This reaction probably proceeds *via* equilibrium with the corresponding vicinal N/B Lewis pair. Most products were characterized by X-ray diffraction.

### Introduction

Frustrated Lewis pairs (FLPs) have become well known for their ability of binding and/or activating a variety of small molecules. The reactivity of *e.g.* the ethylene-bridged P/B FLP 1<sup>3</sup> toward carbon monoxide is remarkable. Compound 1 adds both the phosphane Lewis base and the borane Lewis acid to the CO carbon atom to yield the five-membered heterocyclic carbonyl compound 2. This behavior of FLP 1 remotely resembles CO/metal coordination chemistry in contrast to the conventional role of such P/B pairs as ambiphilic ligands. Compound 1 reacts analogously with nitric oxide (NO), again serving in a "pseudometal like" role, giving the persistent nitroxide radical 3 (see Scheme 1).

There are very few first cases where frustrated Lewis pairs induce coupling of two such element oxides. Stephan  $et\ al.$  had found the formation of 4 upon treatment of a 2:1 mixture of  $B(C_6F_5)_3$  and  $P^tBu_3$  with  $CO/H_2$  (ref. 8) and we recently reported the first example of a CO/NO coupling reaction at the FLP 1 and elucidated the pathway of the unprecedented formation of product 5.9 It is also well known that a variety of alkyl or aryl transition metal and main group metal complexes, respectively, undergo reactions involving the head to head coupling of two nitric oxide (NO) molecules to give the respective O-metallated N-hydrocarbyl-N-nitroso-hydroxylaminato metal complexes. The tungsten and aluminum complexes 6 and 7 are representative examples. There have also been early reports about the

We had previously shown that enamines can react with Piers' borane  $^{17}$  by addition of the Lewis acidic borane at the nucleophilic  $\beta$ -carbon atom. Nevertheless some of these systems were  $H_2$ -activators, probably via dissociation of the Lewis pair adduct followed by in situ generation of the respective vicinal N/B FLP. We have now conveniently prepared an enamine/HB( $C_6F_5$ )<sub>2</sub> Lewis pair adduct containing the very bulky tetramethylpiperidinyl (TMP) moiety by an isomerization route starting from the respective allylamine precursor. The TMP-enamine/HB( $C_6F_5$ )<sub>2</sub> C/B adduct was shown to undergo CO/CO and NO/NO coupling reactions in rather differently oriented pathways. This opposing set of reactions will be described and discussed in this account.



Scheme 1

reaction of Et<sub>3</sub>B with NO to apparently give a Et<sub>2</sub>B-ON(Et)NO type product.<sup>16</sup>

<sup>&</sup>quot;Organisch-Chemisches Institut, Westfälische Wilhelms-Universität Münster, Corrensstraße 40, 48149 Münster, Germany. E-mail: erker@uni-muenster.de

<sup>&</sup>lt;sup>b</sup>Institut für Anorganische und Analytische Chemie, Goethe-Universität Frankfurt, Maxvon-Laue-Straße 7, 60438 Frankfurt am Main, Germany

<sup>†</sup> Electronic supplementary information (ESI) available: Experimental and analytical details, spectral data and crystallographic data. CCDC 1506872–1506879. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c6sc04459j

### Results and discussions

## CO/CO coupling with the enamine/HB( $C_6F_5$ ) $_2$ C/B addition product

For this part of our study we started from *N*-allyl tetramethylpiperidine (**8a**) which was isomerized to the enamine **10** (*trans/cis* mixture of ~10:8) by treatment with a catalytic quantity (10 mol%) of the strong boron Lewis acid B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> (toluene, r.t., 2 d).<sup>21</sup> The *trans-/cis*-enamine mixture was isolated by distillation. Compound *trans-10* shows an olefinic [N]–CH =  $^1$ H NMR resonance at  $\delta$  5.75 ppm (dq,  $^3J_{\rm HH}$  = 13.4 Hz,  $^4J_{\rm HH}$  = 1.6 Hz), whereas *cis-***10** features the respective olefinic resonance at  $\delta$  5.68 ppm (dq,  $^3J_{\rm HH}$  = 7.8 Hz,  $^4J_{\rm HH}$  = 1.8 Hz). We assume a reaction pathway (see Scheme 2) that involves hydride abstraction<sup>18,22</sup> in the  $\alpha$ -position to the amine to give the conjugated iminium/hydridoborate salt **9** followed by H<sup>-</sup> addition to the terminal =CH<sub>2</sub> group.

The enamine isomers **10** rapidly add B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> to the enamine carbon atom<sup>18</sup> (n-pentane, r.t., 30 min) to give the zwitterionic adduct **11** (isolated in 93% after workup). Compound **11** can also be obtained directly from the allylamine **8a** by treatment with a stoichiometric quantity of B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> (toluene, r.t., 30 min); in this case *in situ* generation of the enamine is assumed. We isolated compound **11** in 81% from this stoichiometric reaction. Compound **11** shows a <sup>11</sup>B NMR resonance at  $\delta$  –11.9 ppm and a typical set of <sup>19</sup>F NMR signals of  $o_{ij}p_{,m}$ -fluorines of the three C<sub>6</sub>F<sub>5</sub>-groups at boron with  $\Delta \delta$  <sup>19</sup>F<sub>m,p</sub> = 4.5 ppm. The <sup>1</sup>H/<sup>13</sup>C NMR iminium [N]=CH- resonances were located at  $\delta$  8.74/ $\delta$  189.3 ppm, respectively. Compound **11** was also characterized by X-ray diffraction. The structure is depicted in the ESI.†

The reaction of the enamine isomer mixture **10** with Piers' borane  $[HB(C_6F_5)_2]^{17}$  was carried out analogously (n-pentane, r.t., 30 min) and gave the zwitterionic addition product **12a** as a white solid, isolated in 87% yield. Compound **12a** was characterized by an X-ray crystal structure analysis (suitable single crystals were obtained at -35 °C from a dichloromethane solution covered with a n-pentane layer). In the structure (see Fig. 1) the bulky tetramethylpiperidide unit is part of the iminium functional group. It has the remaining saturated  $C_2$ -unit

$$(C_6F_5)_3B-H$$

$$B(C_6F_5)_3$$

$$1 \text{ equiv.} \qquad B(C_6F_5)_3$$

$$2 \text{ equiv.} \qquad B(C_6F_5)_3$$

$$3 \text{ equiv.} \qquad B(C_6F_5)_3$$

$$4 \text{ equiv.} \qquad B(C_6F_5)_3$$

$$5 \text{ equiv.} \qquad C_6F_5$$




Fig. 1 The molecular structure of the enamine/HB( $C_6F_5$ )<sub>2</sub> addition product 12a (thermal ellipsoids are shown at the 50% probability level). Selected bond lengths (Å) and angles (°): N1–C1 1.299(2), C1–C2 1.463(2), C2–C3 1.549(2), C2–B1 1.682(3), B1–C21 1.636(3), B1–C31 1.649(3); N1–C1–C2 131.8(2), C1–C2–C3 108.6(2), C1–C2–B1 106.2(1), N1–C1–C2–B1 125.4(2).

attached, which features the  $-B(H)(C_6F_5)_2$  substituent bonded to carbon atom C2. The boron atom shows a pseudo-tetrahedral coordination geometry with a sum of heavy atom bond angles of  $\sum B1^{CCC} = 335.6^{\circ}$ . The structure of the  $B(C_6F_5)_3$ /enamine adduct 11 is very similar (see the ESI†).

In solution (CD<sub>2</sub>Cl<sub>2</sub>) compound **12a** shows the  $^1\text{H}/^{13}\text{C}$  NMR resonances of the [N]=CH- iminium functionality at  $\delta$  8.12 (d,  $^3J_{\text{HH}}=13.1$  Hz) and  $\delta$  189.2 ppm, respectively. The  $^{11}\text{B}$  NMR signal occurs at  $\delta$  –20.0 ppm as a doublet ( $^1J_{\text{BH}}\sim93$  Hz). The corresponding  $^1\text{H}$  NMR [B]-H resonance was located at 2.74 ppm as a broad 1:1:1:1 intensity quartet. Due to the chiral center (C2) we observe a 1:1 pair of o,p,m- $^{19}\text{F}$  NMR signals and a total of four tetramethylpiperidino  $^1\text{H}$  NMR methyl group signals.

We reacted the enamine/HB( $C_6F_5$ )<sub>2</sub> adduct **12a** with carbon monoxide ( $CH_2Cl_2$ , r.t., 2.0 bar). The reaction was followed *in situ* by NMR spectroscopy. This showed the formation of the new product **13a** (which contained two B( $C_6F_5$ )<sub>2</sub> units) and the boron free enamine **10**. After 3 h reaction time workup eventually furnished the product **13a**, isolated as a white solid in 43% yield. Since we needed two molar equiv. of HB( $C_6F_5$ )<sub>2</sub> for a complete conversion of **12a** to the carbonylation product **13a**, we reacted the precursor **12a** with CO under similar conditions in the presence of an additional molar equivalent of Piers' borane. Workup after 3 h reaction time involving washing with  $CH_2Cl_2$  gave the product **13a** in 41% yield (for details see the ESI†).

The X-ray crystal structure analysis of **13a** (Fig. 2, single crystals were obtained at -35 °C from a THF solution covered with a layer of n-pentane) revealed that a central five-membered heterocycle had been formed by head to tail coupling of two CO-molecules.<sup>8</sup> The carbon atom of one of them (C4) now bears a hydrogen originating from a HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> equivalent and the enamine derived substituent. The other CO equivalent is found C-O bonded between a pair of boron atoms. Both the boron atoms feature pseudotetrahedral coordination geometries ( $\sum B1^{CCC} = 344.4^{\circ}$ ,  $\sum B2^{CCC} = 337.5^{\circ}$ ). The enamine derived residue shows an iminium functionality.

Edge Article Chemical Science




Fig. 2 A projection of the carbonylation product **13a** (thermal ellipsoids are shown at the 50% probability level). Selected bond lengths (Å) and angles (°): N1–C1 1.286(3), C1–C2 1.498(3), C4–B1 1.658(3), B1–O2 1.551(3), O2–C5 1.272(2), B2–C5 1.604(3), C5–O1 1.320(2), O1–C4 1.494(3); C4–O1–C5 111.5(2), B1–O2–C5 113.4(2).

In solution (THF-d<sub>8</sub>) compound **13a** features  $^1\text{H}/^{13}\text{C}$  NMR resonances of the iminium moiety at  $\delta$  8.68 ( $^3J_{\text{HH}} = 10.4$  Hz) and 184.5 ppm ([N]=CH-), respectively. The central CO derived -O-C[B]-O- carbon atom shows a  $^{13}\text{C}$  NMR signal at  $\delta$  213.7 ppm (C5), whereas the [B]CH carbon atom (C4 in Fig. 2) shows a broad  $^{13}\text{C}$  NMR resonance  $\delta$  86.2 ppm.

Compound **13a** shows a pair of <sup>11</sup>B NMR signals at  $\delta$  2.6 and -26.1 ppm ( ${}^1\!J_{\rm BH}=85.8$  Hz). Due to the C2–chirality center both the B(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> units show the <sup>19</sup>F NMR signals of pairs of diastereotopic C<sub>6</sub>F<sub>5</sub> substituents.

We assume that the enamine/HB( $C_6F_5$ )<sub>2</sub> C/B-adduct becomes reversible under the applied carbonylation conditions and we, consequently, observe a reaction under frustrated Lewis pair

Scheme 3

(FLP) conditions.  $^{18,20,23,24}$  CO activation can occur by formation of Piers' borane carbonyl $^{25,26}$  [OC–B(H)(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>] **14** as we had recently shown (we had actually isolated **14** at low temperature and characterized this borane carbonyl by X-ray diffraction). FLP addition of the enamine nucleophile followed by hydridoborate reduction of the acyl group in **15** would then lead to the intermediate **16** which could add to a second equivalent of the borane carbonyl to eventually yield the observed CO coupling product **13a** (see Scheme 3).

The reaction of the enamine/HB( $C_6F_5$ )<sub>2</sub> FLP **12a** with CO proceeds similarly in the presence of one molar equiv. of the strong boron Lewis acid B( $C_6F_5$ )<sub>3</sub>. The reaction was carried out at r.t. in dichloromethane with 2.0 bar carbon monoxide pressure. Workup after 3 h reaction time eventually gave the CO coupling product **13b** as a white solid, isolated in 76% yield. In THF-d<sub>8</sub> solution it shows a pair of <sup>11</sup>B NMR signals at  $\delta$  2.6 and  $\delta$  –17.7 ppm. We monitor a single set of  $o_1p_1$ m-C<sub>6</sub>F<sub>5</sub> <sup>19</sup>F NMR resonances of the exocyclic B( $C_6F_5$ )<sub>3</sub> substituent and the corresponding resonances of a pair of diastereotopic  $C_6F_5$  groups of the endocyclic B( $C_6F_5$ )<sub>2</sub> moiety. Compound **13b** was also characterized by X-ray diffraction. Its structure is analogous to that of **13a** (for details see the ESI†). We assume a similar reaction scheme for the formation of **13b** as we had discussed it for **13a** (see above) (Scheme 4).

#### NO/NO coupling with the enamine/HB( $C_6F_5$ )<sub>2</sub> C/B Lewis pair

The enamine/HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> adduct 12a reacted equally well with nitric oxide (NO). A solution of 12a in dichloromethane was exposed to a NO atmosphere (1.0 bar) for 3 h. Workup by removing the solvent in vacuo and washing of the residue with npentane gave the NO coupling product 17a as a pale yellow solid. It was isolated in 76% yield (see Scheme 5). Single crystals of compound 17a suited for the X-ray crystal structure analysis were obtained from a *n*-pentane solution at -35 °C (see Fig. 3). It shows that two NO molecules had been head to head coupled. This has formed the observed central planar five-membered N<sub>2</sub>O<sub>2</sub>B containing heterocycle. It contains a short N2=N3 linkage and pairs of N-O and B-O bonds in the σ-bond length range. Nitrogen atom N2 has the amino-alkyl substituent attached to it. We note that this does not contain the iminium moiety of the starting material (12a) any more but instead features a saturated tert amine [N]-CH2- unit.

In solution compound 17a shows the  $^1\text{H}$  NMR signals of the [N]–CH<sub>2</sub>–CH– section of the substituent at  $\delta$  2.72, 3.05 (CH<sub>2</sub>,  $^2J_{\text{HH}}=16.4$  Hz,  $^3J_{\text{HH}}=9.5$  Hz,  $^{13}\text{C}$ :  $\delta$  47.9) and  $\delta$  4.80 ppm (CH,  $^{13}\text{C}$ :  $\delta$  69.3), respectively. It features a  $^{11}\text{B}$  NMR resonance at  $\delta$  +12.2 ppm and two sets of  $o.p.m.^{19}\text{F}$  NMR signals of the pair of diastereotopic  $C_6F_5$  substituents at boron.

Scheme 4

Scheme 5

Fig. 3 Molecular structure of the NO coupling product 17a (thermal ellipsoids are shown at the 30% probability level). Selected bond lengths (Å) and angles (°): N1-C1 1.459(3), N2-C2 1.473(3), N2-N3 1.256(3), N2-O1 1.344(2), N3-O2 1.333(3), O1-B1 1.527(3), O2-B1 1.538(3); N1-C1-C2 114.5(2), C2-N2-O1 119.2(2), O1-N2-N3 117.3(2), N2-O1-B1 105.6(2), N3-O2-B1 110.3(2), O1-B1-O2 97.6(2).

From the structure of compound 17a (see Fig. 3), which contains the saturated [N]-CH<sub>2</sub>-CH- moiety, we must assume that the NO reaction started from the N/B FLP isomer 18a (see Scheme 5 and below). 20,23,24 NO addition to boron would give the radical 19 which might have rearranged to its isomer 20.20,27 Trapping with a second equivalent of NO would then straightforwardly lead to 17a. We must stress that this pathway which is depicted in Scheme 5 provides a mere possible rationalization for the observed formation of the NO/NO coupling product; so far none of the alleged intermediates has been observed directly.

We have, however, obtained additional indirect evidence for the formation of the N/B FLP 18a, which is apparently in an endothermal equilibrium situation with 12a (a situation that had previously been observed with other enamine/RB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> borane addition products as well).18 In this case we added a slight excess of phenylacetylene to a solution of the enamine/

 $HB(C_6F_5)_2$  adduct 12a in dichloromethane solution. The mixture was kept at r.t. for 1 h. Workup then gave the reaction product 22, which we isolated in 91% yield (see Scheme 6).

The X-ray crystal structure analysis of 22 (see Fig. 4) revealed that we had apparently trapped the typical N/B FLP reaction product with the terminal acetylene.28 The alkyne had become deprotonated to give the ammonium cation section of 22 and the resulting alkynyl carbanion had become attached to the boron atom. In the zwitterionic compound a saturated -CH<sub>2</sub>-CH(CH<sub>3</sub>)- bridge is formed connecting the boron and nitrogen atoms. Consequently, we observe the 13C NMR acetylide signals at  $\delta$  111.9 and 98.4 ppm, respectively (<sup>11</sup>B:  $\delta$  –17.3 ppm) and a broad NH <sup>1</sup>H NMR resonance at  $\delta$  6.08 ppm. We assume that the adduct formation of the enamine 10 with Piers' borane to give 12a is reversible in solution. From this equilibrium there might be an (endothermic) pathway of hydroboration of the enamine double bond to generate the N/B FLP 18a in situ in an equilibrium situation. This we have not observed as such,18 but we assume that its typical N/B FLP reaction with the added




Fig. 4 A projection of the molecular structure of compound 22 (thermal ellipsoids are shown at the 30% probability level). Selected bond lengths (Å) and angles (°): B1-C4 1.593(3), C4-C5 1.206(3), C5-C41 1.439(3), B1-C2 1.677(3), C1-C2 1.532(3), C1-N1 1.541(3); B1-C4-C5 174.3(2), C4-C5-C41 175.7(2), N1-C1-C2-B1 167.9(2), ∑N1<sup>CCC</sup> 343.5.

Edge Article Chemical Science

$$\begin{array}{c|c} & & & \\ & & &$$

phenylacetylene trapping reagent eventually led to the formation of the observed product 22 (see Scheme 6 and Fig. 4).

This set the scene for the formation of the isopropyl substituted enamine/HB( $C_6F_5$ )<sub>2</sub> adduct 12b and its reaction with nitric oxide. We treated the substituted allyl amine precursor  $8b^{20,29}$  with one molar equiv. of HB( $C_6F_5$ )<sub>2</sub> in n-pentane. This gave a yellow solution within 10 min. Workup after 30 min reaction time eventually gave the zwitterionic iminium/hydridoborate product 12b, which we isolated in 89% yield (see Scheme 7). The NMR spectra showed the presence of the iminium ion moiety ( $^{13}C$ :  $\delta$  187.1 ppm), and a  $^{11}B$  NMR resonance at  $\delta$  –21.1 ppm (d,  $^1J_{\rm BH}$   $\sim$  92 Hz). We observed the  $^1H$  NMR signals of the [N]=CH– proton at  $\delta$  8.40 ppm, the –CHMe<sub>2</sub> isopropyl–H at  $\delta$  2.06 and the [B]H at  $\sim$ 2.84 ppm (broad 1:1:1 quartet).

We also did the analogous reaction of 8b with the isotopically labelled  $\mathrm{DB}(C_6F_5)_2$  reagent. In the product we now find the deuterium atom scrambled over these two CH and the BH position listed above, which indicates that in this case the isomerization reaction of 8b to the respective enamine is probably taking place by means of hydroboration/retro-hydroboration (see Scheme 7, for further details see the ESI†).




Fig. 5 A view of the molecular structure of the NO coupling product 17b (thermal ellipsoids are shown at the 50% probability level). Selected bond lengths (Å) and angles (°): N1–C1 1.455(5), N2–C2 1.482(5), N2–N3 1.252(5), N2–O1 1.334(4), N3–O2 1.323(5), O1–B1 1.508(6), O2–B1 1.536(6); N1–C1–C2 113.5(4), C2–N2–O1 118.8(3), O1–N2–N3 117.4(4), N2–O1–B1 105.4(3), N3–O2–B1 109.4(3), O1–B1–O2 98.3(3).

Compound **12b** reacted rapidly with nitric oxide. A solution of **12b** in dichloromethane was stirred for 3 h at r.t. in a NO atmosphere (1.0 bar). Workup involving extraction with *n*-pentane eventually gave the NO coupling product **17b**, which we isolated as a pale yellow solid in 69% yield (see Scheme 7). It was characterized by X-ray diffraction (see Fig. 5) and by NMR spectroscopy. The <sup>1</sup>H NMR spectrum shows a total of six methyl group signals and an ABX type pattern of the [N]–CH<sub>2</sub>–CH–moiety [CH<sub>2</sub>:  $\delta$  2.95, 3.08 ( ${}^2J_{\rm HH}$  = 16.3 Hz,  ${}^3J_{\rm HH}$  = 9.8 Hz), 4.44 (CH);  ${}^{13}{\rm C}$ :  $\delta$  44.2, 78.9 ppm]. The <sup>11</sup>B NMR resonance of compound **17b** occurs at  $\delta$  +12.6 ppm and we have observed two sets of o,p,m- ${}^{19}{\rm F}$  NMR signals of the pair of diastereotopic C<sub>6</sub>F<sub>5</sub> substituents at boron.

### Conclusions

The HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> adduct of the very bulky enamine 8a reacts very differently with the element oxides CO and NO.30 Its facile reaction with carbon monoxide results in a selective head to tail coupling of the CO molecule. Formally, the initial sequence might well be regarded as an addition reaction of a C/B frustrated Lewis pair<sup>31</sup> to carbon monoxide. We assume a reaction pathway initiated by reversible HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> cleavage from the adduct. It may then combine via nucleophilic enamine addition to Piers' borane carbonyl  $[(C_6F_5)_2B(H)-CO 14]$ , a reactive borane carbonyl that we had recently prepared and characterized. Hydridoborate carbonyl reduction and B/O addition to a second borane Lewis acid CO equivalent might then close the reaction cycle as it was shown in Scheme 3. It seems an essential consequence of this reaction that the enamine/HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> adduct formation is reversible and that, consequently, the 10/  $HB(C_6F_5)_2 \rightleftharpoons 12a$  system may function as a reactive C/B frustrated Lewis pair.

The reaction of 12a with NO revealed another reaction type of the HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>/enamine C/B Lewis pair adduct. We here observe the formation of the product of selective NO/NO coupling. This is a typical reaction mode observed for many metal alkyls. 10-15 Here we have probably found the analogous reaction of an alkyl- $B(C_6F_5)_2$  functional group. Therefore, we assume that this reaction is initiated by the reaction of NO with the -CHR-B(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> function of the in situ generated vicinal N/B Lewis pair, which may be formed by cleavage of the enamine/HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> Lewis adduct followed by anti-Markovnikov hydroboration. The reaction then seems to follow the usual pathway, as it is often observed in the selective NO/NO coupling of metal hydrocarbyls,15 here to eventually give the respective boron based-ON(R)NO products 17. The NO coupling products<sup>16</sup> 17 are topologically related to cupferron (24), a reagent that had frequently be used for chelate metal complexation.32-34 It is formed by nitrosation of N-phenylhydroxylamine (23) (Scheme 8). We will find out whether the

Chemical Science Ed

boron based *N*-nitrosohydroxylaminato groups could potentially be used directly as reagents for metal complex formation.

The dual reaction pathway of the enamine/ $HB(C_6F_5)_2$  Lewis pair 12 with CO and with NO indicates a remarkable variability of bulky Lewis acid/Lewis base combinations in small molecule chemistry.

### Acknowledgements

Financial support from the European Research Council is gratefully acknowledged.

### Notes and references

- 1 D. W. Stephan and G. Erker, *Angew. Chem., Int. Ed.*, 2015, **54**, 6400.
- 2 (a) Frustrated Lewis Pairs I, Uncovering and Understanding, Top. Curr. Chem., ed. G. Erker and D. W. Stephan, Heidelberg, 2013, vol. 332; (b) Frustrated Lewis Pairs II, Expanding the Scope, Top. Curr. Chem., ed. G. Erker and D. W. Stephan, Heidelberg, 2013, vol. 334.
- 3 P. Spies, G. Erker, G. Kehr, K. Bergander, R. Fröhlich, S. Grimme and D. W. Stephan, *Chem. Commun.*, 2007, 5072.
- 4 M. Sajid, A. Lawzer, W. Dong, C. Rosorius, W. Sander, B. Schirmer, S. Grimme, C. G. Daniliuc, G. Kehr and G. Erker, *J. Am. Chem. Soc.*, 2013, **135**, 18567.
- 5 O. Ekkert, G. G. Miera, T. Wiegand, H. Eckert, B. Schirmer, J. L. Petersen, C. G. Daniliuc, R. Fröhlich, S. Grimme, G. Kehr and G. Erker, *Chem. Sci.*, 2013, 4, 2657.
- 6 (a) S. Bontemps, G. Bouhadir, K. Miqueu and D. Bourissou, J. Am. Chem. Soc., 2006, 128, 12056; (b) J. Grobe, K. Lütke-Brochtrup, B. Krebs, M. Läge, H.-H. Niemeyer and E.-U. Würthwein, Z. Naturforsch., B: J. Chem. Sci., 2006, 61, 882.
- 7 (a) M. Sajid, A. Stute, A. J. P. Cardenas, B. J. Culotta, J. A. M. Hepperle, T. H. Warren, B. Schirmer, S. Grimme, A. Studer, C. G. Daniliuc, R. Fröhlich, J. L. Petersen, G. Kehr and G. Erker, J. Am. Chem. Soc., 2012, 134, 10156; (b) M. Sajid, G. Kehr, T. Wiegand, H. Eckert, C. Schwickert, R. Pöttgen, A. J. P. Cardenas, T. H. Warren, R. Fröhlich, C. G. Daniliuc and G. Erker, J. Am. Chem. Soc., 2013, 135, 8882; (c) J. C. M. Pereira, M. Sajid, G. Kehr, A. M. Wright, B. Schirmer, Z.-W. Qu, S. Grimme, G. Erker and P. C. Ford, J. Am. Chem. Soc., 2014, 136, 513; (d) R. Liedtke, F. Scheidt, J. Ren, B. Schirmer, A. J. P. Cardenas, C. G. Daniliuc, H. Eckert, T. H. Warren, S. Grimme, G. Kehr and G. Erker, J. Am. Chem. Soc., 2014, 136, 9014.
- 8 R. Dobrovetsky and D. W. Stephan, *J. Am. Chem. Soc.*, 2013, 135, 4974.
- 9 (*a*) K.-Y. Ye, G. Kehr, C. G. Daniliuc, L. Liu, S. Grimme and G. Erker, *Angew. Chem., Int. Ed.*, 2016, 55, 9216; (*b*) A description of an additional new example of this CO/NO coupling reaction was added to the ESI† of this paper.
- 10 Tungsten: (a) S. R. Fletcher, A. Shortland, A. C. Skapski and G. Wilkinson, *J. Chem. Soc., Dalton Trans.*, 1972, 922; (b) S. R. Fletcher and A. C. Skapski, *J. Organomet. Chem.*, 1973, 59, 299; (c) A. Shortland and G. Wilkinson, *J. Chem. Soc.*,

- Dalton Trans., 1973, 872; (d) P. Legzdins, S. J. Rettig and L. Sánchez, Organometallics, 1988, 7, 2394; (e) S. V. Evans, P. Legzdins, S. J. Rettig, L. Sánchez and J. Trotter, Acta Crystallogr., Sect. C: Cryst. Struct. Commun., 1995, C51, 584.
- 11 Zirconium: (a) P. C. Wailes, H. Weigold and A. P. Bell, J. Organomet. Chem., 1972, 34, 155; (b) A. R. Middleton and G. Wilkinson, J. Chem. Soc., Dalton Trans., 1980, 1888; (c) C. J. Jones, J. A. McCleverty and A. S. Rothin, J. Chem. Soc., Dalton Trans., 1985, 405; (d) G. Fochi, C. Floriani, A. Chiesi-Villa and C. Guastini, J. Chem. Soc., Dalton Trans., 1986, 445.
- 12 Other transition metals: (a) J. D. Wilkins and M. G. B. Drew, J. Organomet. Chem., 1974, 69, 111; (b) A. R. Middleton and G. Wilkinson, J. Chem. Soc., Dalton Trans., 1981, 1898; (c) S. C. Puiu and T. H. Warren, Organometallics, 2003, 22, 3974; (d) H. Lei, B. D. Ellis, C. Ni, F. Grandjean, G. J. Long and P. P. Power, Inorg. Chem., 2008, 47, 10250; (e) I. J. Casely, Y. Suh, J. W. Ziller and W. J. Evans, Organometallics, 2010, 29, 5209; (f) S.-C. So, W.-M. Cheung, G.-C. Wang, E. K. Huang, M.-K. Lau, Q.-F. Zhang, H. H.-Y. Sung, I. D. Williams and W.-H. Leung, Organometallics, 2014, 33, 4497.
- 13 Main group metals: (a) E. Müller and H. Metzger, Chem. Ber.,
  1956, 89, 396; (b) I. S. Butler and M. L. Newbury, J. Coord.
  Chem., 1976, 6, 195; (c) S. Amirkhalili, A. J. Conway and
  J. D. Smith, J. Organomet. Chem., 1978, 149, 407; (d)
  S. Amirkhalili, P. B. Hitchcock, J. D. Smith and
  J. G. Stamper, J. Chem. Soc., Dalton Trans., 1980, 2493.
- 14 Carbanion equivalents: (a) M. J. Danzig, R. F. Martel and S. R. Riccitiello, J. Org. Chem., 1961, 65, 3327; (b) E. V. Arnold, L. K. Keefer and J. A. Hrabie, Tetrahedron Lett., 2000, 41, 8421; (c) J. A. Hrabie, E. V. Arnold, M. L. Citro, C. George and L. K. Keefer, J. Org. Chem., 2000, 65, 5745.
- Reviews: (a) J. A. McCleverty, Chem. Rev., 1979, 79, 53; (b)
   J. A. Hrabie and L. K. Keefer, Chem. Rev., 2002, 102, 1135.
- 16 (a) S. J. Brois, Tetrahedron Lett., 1964, 345; (b) See also:M. Inatome and L. Kuhn, Adv. Chem. Ser., 1964, 42, 183.
- 17 (a) D. J. Parks, R. E. von H. Spence and W. E. Piers, Angew.
   Chem., Int. Ed. Engl., 1995, 34, 809; (b) D. J. Parks,
   W. E. Piers and G. P. A. Yap, Organometallics, 1998, 17, 5492.
- 18 (a) S. Schwendemann, R. Fröhlich, G. Kehr and G. Erker, Chem. Sci., 2011, 2, 1842; (b) S. Schwendemann, S. Oishi, S. Saito, R. Fröhlich, G. Kehr and G. Erker, Chem.-Asian J., 2013, 8, 212.
- (a) D. M. Hodgson, C. D. Bray and N. D. Kindon, *J. Am. Chem. Soc.*, 2004, **126**, 6870; (b) D. M. Hodgson, C. D. Bray, N. D. Kindon, N. J. Reynolds, S. J. Coote, J. M. Um and K. N. Houk, *J. Org. Chem.*, 2009, 74, 1019.
- 20 T. Wang, G. Kehr, L. Liu, S. Grimme, C. D. Daniliuc and G. Erker, *J. Am. Chem. Soc.*, 2016, **138**, 4302.
- 21 (a) A. G. Massey and A. J. Park, J. Organomet. Chem., 1964, 2, 245; (b) A. G. Massey, A. J. Park and F. G. A. Stone, Proc. Chem. Soc., London, 1963, 212.
- 22 (a) F. Focante, P. Mercandelli, A. Sironi and L. Resconi, Coord. Chem. Rev., 2006, 250, 170; (b) See also: H. Volz and H.-H. Kiltz, Liebigs Ann. Chem., 1971, 752, 86.

**Edge Article** 

23 See for a comparison: (a) M. A. Dureen and D. W. Stephan, *J. Am. Chem. Soc.*, 2010, 132, 13559; (b) K. Chernichenko, M. Nieger, M. Leskelä and T. Repo, *Dalton Trans.*, 2012, 41, 9029; (c) K. Chernichenko, Á. Madarász, I. Pápai, M. Nieger, M. Leskelä and T. Repo, *Nat. Chem.*, 2013, 5, 718; (d) K. Chernichenko, B. Kótai, I. Pápai, V. Zhivonitko, M. Nieger, M. Leskelä and T. Repo, *Angew. Chem., Int. Ed.*, 2015, 54, 1749; (e) M.-A. Légaré, M.-A. Courtemanche, É. Rochette and F.-G. Fontaine, *Science*, 2015, 349, 513; (f) M.-A. Courtemanche, A. P. Pulis, É. Rochette, M.-A. Légaré, D. W. Stephan and F.-G. Fontaine, *Chem. Commun.*, 2015, 51, 9797; (g) K. Chernichenko, M. Lindqvist, B. Kótai, M. Nieger, K. Sorochkina, I. Pápai and T. Repo, *J. Am. Chem. Soc.*, 2016, 138, 4860.

- 24 (a) D. Winkelhaus, B. Neumann, H.-G. Stammler, R. J. F. Berger, Y. V. Vishnevskiy and N. W. Mitzel, *Chem. Eur. J.*, 2012, 18, 9312; (b) L. A. Körte, S. Blomeyer, S. Heidemeyer, A. Mix, B. Neumann and N. W. Mitzel, *Chem. Commun.*, 2016, 52, 9949.
- 25 M. Sajid, G. Kehr, C. G. Daniliuc and G. Erker, Angew. Chem., Int. Ed., 2014, 53, 1118.
- 26 See also: (a) A. G. Burg and H. I. Schlesinger, J. Am. Chem. Soc., 1937, 59, 780; (b) T. P. Fehlner and W. S. Koski, J. Am. Chem. Soc., 1965, 87, 409; (c) J. C. Carter, A. L. Moyé and G. W. Luther III, J. Am. Chem. Soc., 1974, 96, 3071.
- 27 See for a comparison: C. Ollivier and P. Renaud, *Chem. Rev.*, 2001, **101**, 3415.
- 28 (a) M. A. Dureen and D. W. Stephan, J. Am. Chem. Soc., 2009,
  131, 8396; (b) C. M. Mömming, S. Frömel, G. Kehr,
  R. Fröhlich, S. Grimme and G. Erker, J. Am. Chem. Soc.,
  2009, 131, 12280; (c) M. A. Dureen, C. C. Brown and
  D. W. Stephan, Organometallics, 2010, 29, 6594; (d) T. Voss,
  T. Mahdi, E. Otten, R. Fröhlich, G. Kehr, D. W. Stephan
  and G. Erker, Organometallics, 2012, 31, 2367; (e) B.-H. Xu,
  K. Bussmann, R. Fröhlich, C. D. Daniliue,

- J. G. Brandenburg, S. Grimme, G. Kehr and G. Erker, *Organometallics*, 2013, **32**, 6745.
- 29 V. Sumerin, F. Schulz, M. Atsumi, C. Wang, M. Nieger, M. Leskelä, T. Repo, P. Pyykkö and B. Rieger, *J. Am. Chem. Soc.*, 2008, 130, 14117.
- 30 D. W. Stephan and G. Erker, Chem. Sci., 2014, 5, 2625.
- 31 (a) D. Holschumacher, T. Bannenberg, C. G. Hrib, P. G. Jones and M. Tamm, Angew. Chem., Int. Ed., 2008, 47, 7428; (b) P. A. Chase and D. W. Stephan, Angew. Chem., Int. Ed., 2008, 47, 7433; (c) P. A. Chase, A. L. Gille, T. M. Gilbert and D. W. Stephan, Dalton Trans., 2009, 7179; (d) D. Holschumacher, C. Taouss, T. Bannenberg, C. G. Hrib, C. G. Daniliuc, P. G. Jones and M. Tamm, Dalton Trans., 2009, 6927; (e) M. Alcarazo, C. Gomez, S. Holle and R. Goddard, Angew. Chem., Int. Ed., 2010, 49, 5788; (f) B. Ines, S. Holle, R. Goddard and M. Alcarazo, Angew. Chem., Int. Ed., 2010, 49, 8389; (g) D. Holschumacher, T. Bannenberg, K. Ibrom, C. G. Daniliuc, P. G. Jones and M. Tamm, Dalton Trans., 2010, 39, 10590; (h) M. A. Dureen, C. C. Brown and D. W. Stephan, Organometallics, 2010, 29, 6422; (i) E. L. Kolychev, T. Bannenberg, M. Freytag, C. G. Daniliuc, P. G. Jones and M. Tamm, Chem.-Eur. J., 2012, 18, 16938.
- 32 (a) C. S. Marvel and O. Kamm, *J. Am. Chem. Soc.*, 1919, 41, 276; (b) C. S. Marvel, *Org. Synth.*, 1925, 4, 19; (c) D. van der Helm, L. L. Merritt Jr, R. Degeilh and C. H. MacGillavry, *Acta Crystallogr.*, 1965, 18, 355.
- 33 See for a comparison: (a) K. G. Orrell, V. Šik and D. Stephenson, *Magn. Reson. Chem.*, 1987, 25, 1007; (b)
   D. Beaudoin and J. D. Wuest, *Chem. Rev.*, 2016, 116, 258.
- 34 See also: (a) N. Xu, J. H. Christian, N. S. Dalal, E. G. Abucayon, C. Lingafelt, D. R. Powell and G. B. Richter-Addo, *Dalton Trans.*, 2015, 44, 20121; (b) S. Kundu, S. C. E. Stieber, M. G. Ferrier, S. A. Kozimor, J. A. Bertke and T. H. Warren, *Angew. Chem., Int. Ed.*, 2016, 55, 10321.