# Chemical Science



### **EDGE ARTICLE**

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2017, 8, 4511

# Enantioselective $\gamma$ -borylation of unsaturated amides and stereoretentive Suzuki-Miyaura cross-coupling†

Gia L. Hoang and James M. Takacs \*\* \*\*

Received 9th March 2017 Accepted 19th April 2017

DOI: 10.1039/c7sc01093a

rsc.li/chemical-science

The rhodium-catalyzed, directed catalytic asymmetric hydroboration of  $\gamma$ , $\delta$ -unsaturated amides affords a direct route to chiral acyclic secondary  $\gamma$ -borylated carbonyl derivatives in high enantiomeric purity. In contrast to a similar  $\beta$ -borylated amide derivative, the  $\gamma$ -borylated amide undergoes Suzuki–Miyaura cross-coupling with stereoretention. The utility of the boronic ester products is further illustrated by other stereospecific C–B bond transformations leading to  $\gamma$ -amino acid derivatives, 1,4-amino alcohols, and 5-substituted- $\gamma$ -lactone and  $\gamma$ -lactam ring systems.

#### Introduction

Chiral boronic esters are valuable synthetic intermediates for a variety of stereospecific transformations.<sup>1-3</sup> Consequently, an assortment of enantioselective routes for their preparation are under active development.<sup>4-19</sup> We have focused on the rhodium-catalyzed catalytic asymmetric hydroboration (CAHB) of  $\beta$ , $\gamma$ -unsaturated amides,<sup>20</sup> esters,<sup>20 $\alpha$ </sup> and oxime ethers<sup>21</sup> for the preparation of functionalized chiral boronic esters. For example, disubstituted alkenes such as (E)-1 undergo highly enantioselective  $\beta$ -borylation by pinacolborane (pinBH) when catalyzed by the combination of (R)-L1 with [Rh(nbd)<sub>2</sub>BF<sub>4</sub>] (*i.e.*, 2:1 L1:Rh). CAHB followed by oxidation of the C–B bond affords  $\beta$ -hydroxy amide (S)-2 in an enantiomeric ratio (er) greater than 99:1 (Fig. 1).<sup>20 $\alpha$ </sup>

In an effort to expand the scope of CAHB,  $\gamma$ , $\delta$ -unsaturated amides such as 3 were explored. Such substrates differ from 1 in a number of important ways including, (i) the alkene is more remote than in the  $\beta$ , $\gamma$ -unsaturated amides, (ii) substrate 3 contains two distinctly different alkene moieties, although each is ostensibly positioned  $\gamma$ , $\delta$ - with respect to the carbonyl directing group, and (iii) enantioselective borylation of the endocyclic alkene requires controlling the stereochemical course by desymmetrization rather than  $\pi$ -facial discrimination, a fundamentally different set of requirements. In the event, substrate 3 undergoes CAHB/oxidation to give the monounsaturated  $\gamma$ -hydroxy amide (1S,3S)-4 in high yield (80%), regioselectivity (>20:1) and enantioselectivity (99:1 er). Only the endocyclic double bond in 3 undergoes reaction; the

pendant acyclic alkene is untouched. We now report the efficient, regioselective  $\gamma$ -borylation of a similarly disposed alkene in  $\gamma$ , $\delta$ -unsaturated amide **5a** (Fig. 1). CAHB by pinBH (0.5% [Rh(nbd)<sub>2</sub>BF<sub>4</sub>/2 (*R*)-L1]) followed by oxidation affords chiral  $\gamma$ -hydroxy amide (*S*)-**7a** in 79% overall yield (97 : 3 er). While conjugate addition<sup>11</sup> and C–H activation<sup>10α</sup> methodologies provide efficient alternatives to CAHB for enantioselective  $\beta$ -borylation of carbonyl compounds, direct  $\gamma$ -borylation is unique to CAHB.




Fig. 1 CAHB/oxidation of  $\beta$ , $\gamma$ - versus  $\gamma$ , $\delta$ -unsaturated amides.

Department of Chemistry, University of Nebraska-Lincoln, Lincoln, NE 68588-0304, USA. E-mail: jtakacs1@unl.edu

† Electronic supplementary information (ESI) available: Catalyst optimization data; experimental procedures; compound characterization data; spectra. See DOI: 10.1039/c7sc01093a

Results and discussion

**Chemical Science** 

The  $\gamma$ , $\delta$ -unsaturated amides (E)-5b-n shown in Fig. 2, along with several related structures, were treated with the catalyst system for  $\gamma$ -borylation (*i.e.*, 2 : 1 combinations of **L1** or **L2** with [Rh(nbd)<sub>2</sub>BF<sub>4</sub>], 0.5 mol% unless otherwise noted). In addition to phenyl amide 5a described above, the corresponding Weinreb amide 5b, the morpholine amide<sup>24</sup> 5c and benzyl amide 5d undergo CAHB to afford the intermediate γ-borylated amides 6b-d with high levels of enantioselectivity. Within this series of amides, the secondary amides (i.e., N-phenyl and N-benzyl) give the highest  $\gamma$ -selectivity (>20 : 1). Oxidations of **6b-d** afford the corresponding chiral  $\gamma$ -hydroxy amides 7b-d ( $\geq 94:6$  er). CAHBs of 5c and 5d were carried out on gram scale giving chiral boronic esters 6c (82%, 94.5 : 5.5) and 6d (81%, 96.5 : 3.5 er) in good yield and without loss of enantioselectivity. CAHB/ oxidation of an isomeric (Z)-alkene, benzyl amide (Z)-5 $\mathbf{d}$ , affords results similar to those obtained with the (*E*)-isomer.

In contrast to amides 5a–d, the analogous ester 8 is largely recovered unchanged upon attempted CAHB; only trace amounts of borylated products are identified along with some evidence for alkene isomerization.  $\delta$ , $\epsilon$ -Unsaturated amide 9, a one-carbon homologue of benzyl amide (E)-5d, is found to be

considerably less reactive and less selective. Complete consumption of starting material requires 2% catalyst loading to afford 56% yield of a mixture of borylated products.<sup>25</sup>

Benzyl amides **5e-g** containing heteroaromatic ring systems are nonetheless good substrates under the standard CAHB/ oxidation conditions yielding 7e-g, respectively. Certain branched alkyl substituents (i.e., 5h-j) are also well tolerated. In particular, the chiral substrate (E)-5i demonstrates that (i) the proximal disubstituted alkene with respect to the amide directing group undergoes CAHB while the more distal trisubstituted alkene is untouched and (ii) the stereochemical course of γ-borylation is efficiently catalyst controlled. CAHB/oxidation with (R)-L2 affords (4S,7S)-7 $\mathbf{j}$ ; (S)-L2 affords (4R,7S)-7 $\mathbf{j}$ . The silyl ether moiety in 5k is tolerated and affords 7k (78%, 97:3 er). Chiral acetal 5l again undergoes catalyst controlled γ-borylation with high diastereoselectivity; (R)-L1 affords (S,S)-7l (72%, 95:5 dr); (S)-L1 affords (R,S)-7l in the same yield and diastereomer ratio. However, substrate 5m, in which the chiral acetal moiety is in closer proximity to the site of hydroboration, shows a strong matched/mismatched effect. While (R)-L1 affords (R,S)-7m (70%, 92 : 8 dr), the catalyst employing (S)-L1 gives rise to a complex mixture of regioisomers 10. Substrate  $5n (R^1 = Me)$ also exhibits only modest regioselectivity (3:1), perhaps due to




Fig. 2 Substrate scope for CAHB of  $\gamma$ ,δ-unsaturated amides. <sup>a</sup>Unless otherwise noted all reaction use 0.5% [Rh(nbd)<sub>2</sub>BF<sub>4</sub>/2 (R)-L1], 1.5 equiv. pinBH, THF, 40 °C (12 h) followed by oxidation using H<sub>2</sub>O<sub>2</sub>/aq. NaOH. Unless otherwise noted, the isolated yield is that of the major regioisomer and reflects the average of three experiments generally exhibiting a spread of ±2%; regioselectivity is determined from the crude <sup>1</sup>H NMR of 7. Enantiomer ratios (er) are determined by chiral HPLC analysis; diastereomer ratios (dr) are determined for the purified mixture of diastereomers by integrating major and minor <sup>13</sup>C NMR resonances. <sup>b</sup>2.0% [Rh(nbd)<sub>2</sub>BF<sub>4</sub>/2 (R)-L1]. <sup>c</sup>Oxidation conditions: NaBO<sub>3</sub>/H<sub>2</sub>O. <sup>d</sup>1.0% [Rh(nbd)<sub>2</sub>BF<sub>4</sub>/2 L1]; (R)-L1 is used unless noted otherwise in the figure. <sup>e</sup>1.0% [Rh(nbd)<sub>2</sub>BF<sub>4</sub>/2 L2]. <sup>f</sup>er is determined by <sup>19</sup>F NMR of the corresponding Mosher ester (see ESI for details†).

**Edge Article** 

the size of the vinyl substituent compared to other derivatives described above; however, CAHB proceeds in good yield and high enantioselectivity (61%, 95:5 er).

Having developed an efficient method for the  $\gamma$ -borylation of γ,δ-unsaturated amides, Suzuki-Miyaura cross-coupling of 11 was examined (Fig. 3). Stereochemical aspects of the palladiumcatalyzed cross-coupling of chiral secondary organoboron derivatives have recently attracted a great deal attention. Molander,26 Suginome,27 and Hall28 reported that β-borylated carbonyl derivatives 12-14, whether as the boronic ester or the trifluoroborate salt, undergo cross-coupling with stereoinversion. The stereochemical course is rationalized by intramolecular coordination between the carbonyl oxygen and the boron atom of the boronic ester or the partially hydrolyzed trifluoroborate. The intramolecular coordination promotes invertive transmetallation resulting in overall stereoinversion for cross-coupling. Biscoe<sup>19a,29</sup> also found stereoinversion for simple substrates lacking functionality needed for coordination to boron during the course of transmetallation (e.g., 15). On the other hand, Suginome<sup>30</sup> reported that boracyclic intermediate 16 undergoes cross-coupling with stereoretention. Similarly, Morken<sup>31</sup> reported that 17 undergoes hydroxyl-directed, innersphere, retentive transmetallation and overall cross-coupling with stereoretention. However, when the hydroxyl is onecarbon further removed, 18 fails to undergo cross-coupling under the otherwise same conditions. We have previously shown that 19 undergoes cross-coupling with overall stereoretention.22

Chiral boronic ester **6c** (*i.e.*, the morpholine amide) was converted to its corresponding trifluoroborate salt  $\mathbf{11c^{32}}$  and subjected to palladium-catalyzed cross-coupling using the Buchwald cataCXium® A Pd G3 (**20**) precatalyst.<sup>22,33</sup> Cross-coupling with chlorobenzene yields amide (*S*)-**21c** (63%); 4-chloroanisole yields (*S*)-**22c** (52%). The products are obtained with essentially complete overall stereoretention.<sup>34</sup> We find that




Fig. 3 Stereochemical course of Suzuki–Miyaura cross-coupling with chiral secondary boronic esters or trifluoroborate salts.




Fig. 4 Stereospecific transformations of organoboranes **6b–d**. Conditions: (a) NaOH/H<sub>2</sub>O<sub>2</sub>; (b) (i) n-BuLi, thiophene, (ii) NBS; (c) BCl<sub>3</sub>, BnN<sub>3</sub>; (d) BCl<sub>3</sub>, PhN<sub>3</sub>; (e) 6 M HCl; (f) LiAlH<sub>4</sub>; (g) (i) vinylMgBr, (ii) I<sub>2</sub>/NaOMe.

the nature of the amide is important to the success of the cross-coupling. In contrast to the tertiary morpholine amide, the analogous secondary amide **11d** does not undergo cross-coupling under the conditions employed for **11c**. Hall *et al.*<sup>28a</sup> reported that  $\beta$ -boronic esters of secondary amides failed to cross-couple in cases where the corresponding tertiary amide coupled smoothly.

Chiral organoboronates are useful for a variety of other stereospecific transformations. Fig. 4 illustrates several examples starting from chiral boronic esters 6b-d; the latter are isolated in 69-82% yield from the corresponding alkenes. Treating 6b with H<sub>2</sub>O<sub>2</sub>/aq. NaOH affords the known chiral 5substituted-γ-lactone 23 (95%).35 As an alternative to palladiumcatalyzed cross-coupling, the morpholine amide derivative 6c undergoes stereoretentive cross-coupling with 2-lithiothiofuran under the conditions developed by Aggarwal<sup>2e</sup> to give 24c (84%). Compound 6c also undergoes BCl3-assisted amination with benzyl azide under the conditions reported by Knochel36 to form the  $\gamma$ -amino acid derivative 25c (65%). Phenyl azide also serves as a good nucleophile in such amination reactions, and 6c is converted to the corresponding N-phenyl γ-amino acid en route to the 5-substituted-γ-lactam 26 (68%) by acid catalyzed cyclization. Benzyl amide derivative 6d is efficiently converted to 1,4aminoalcohol 27d after oxidation of the C-B bond followed by amide reduction with LAH (94%). While the secondary N-benzyl amide 11d failed in the attempted palladium-catalyzed crosscoupling described above, 6d undergoes efficient vinylation in a sequence initiated the by treatment with excess vinyl Grignard;<sup>2h</sup> amide **28d** is formed in high yield (93%).

#### Conclusions

 $\gamma$ , $\delta$ -Unsaturated secondary (*i.e.*, *N*-phenyl and *N*-benzyl) and tertiary (*i.e.*, Weinreb and morpholine) amides undergo efficient rhodium-catalyzed CAHB to afford  $\gamma$ -borylated derivatives in good yield and with high levels of asymmetric induction; enantioselectivity as high as 97:3 er is observed. While two

good alternative methods are available to prepare chiral secondary  $\beta$ -borylated carbonyl compounds, the present method of directed-CAHB provides to our knowledge the first direct route to chiral acyclic secondary  $\gamma$ -borylated carbonyl compounds with high regio- and enantioselectivity.

**Chemical Science** 

A previous study found β- and γ-borylation of related substrates differ in the sense of stereoinduction, *i.e.*,  $\pi$ -facial discrimination. However, it is not the case in the present study; β-borylation of β,γ-unsaturated amide 1 and γ-borylation of the one-carbon homologue γ,δ-unsaturated amide 5 add to the same face of the alkene. In the present study, CAHB of a substrate bearing both di- and trisubstituted alkene moieties (*i.e.*, 5j) occurs only on the disubstituted double bond proximal to the carbonyl group. Chiral substrates 5j and 5l undergo highly diastereoselective CAHB with catalyst control; however, substrate 5m, in which the resident oxygen-bearing stereocenter resides adjacent to the alkene, exhibits a strong matched and mismatched effect with enantiomeric catalysts.

The  $\gamma$ -borylated products are found to undergo stereoretentive palladium-catalyzed Suzuki–Miyaura cross-coupling, presumably via amide-directed inner-sphere stereoretentive transmetallation, as well as stereoretentive C–B to C–C transformations using main group organometallic reagents (e.g., lithium and magnesium). In addition, a variety of other stereospecific transformations are highlighted by the conversions of chiral, secondary  $\gamma$ -boronic esters **6b–d** to 1,4-amino alcohols,  $\gamma$ -amino acid derivatives, and 5-substituted- $\gamma$ -lactone and  $\gamma$ -lactam ring systems. Further studies are in progress.

## Acknowledgements

Funding from the NIH National Institutes of General Medical Sciences (R01-GM100101) is gratefully acknowledged. We thank S. Zhang (UNL) for some preliminary experiments.

#### Notes and references

- 1 For selected recent reviews, see: (a) C.-Y. Wang, J. Derosa and M. R. Biscoe, *Chem. Sci.*, 2015, **6**, 5105; (b) L. Xu, S. Zhang and P. Li, *Chem. Soc. Rev.*, 2015, **44**, 8848; (c) A. H. Cherney, N. T. Kadunce and S. E. Reisman, *Chem. Rev.*, 2015, **115**, 9587; (d) D. Leonori and V. K. Aggarwal, *Angew. Chem., Int. Ed.*, 2015, **54**, 1082.
- 2 (a) Y. Wang, A. Noble, E. L. Myers and V. K. Aggarwal, Angew. Chem., Int. Ed., 2016, 55, 4270; (b) A. Noble, S. Roesner and V. K. Aggarwal, Angew. Chem., Int. Ed., 2016, 55, 15920; (c) A. Fawcett, D. Nitsch, M. Ali, J. M. Bateman, E. L. Myers and V. K. Aggarwal, Angew. Chem., Int. Ed., 2016, 55, 14663; (d) A. Millán, J. R. Smith, J. L.-Y. Chen and V. K. Aggarwal, Angew. Chem., Int. Ed., 2016, 55, 2498; (e) M. Odachowski, A. Bonet, S. Essafi, P. Conti-Ramsden, J. N. Harvey, D. Leonori and V. K. Aggarwal, J. Am. Chem. Soc., 2016, 138, 9521; (f) J. Llaveria, D. Leonori and V. K. Aggarwal, J. Am. Chem. Soc., 2015, 137, 10958; (g) C. Sandford, R. Rasappan and V. K. Aggarwal, J. Am. Chem. Soc., 2015, 137, 10100; (h) R. P. Sonawane, V. Jheengut, C. Rabalakos,

- R. Larouche-Gauthier, H. K. Scott and V. K. Aggarwal, *Angew. Chem., Int. Ed.*, 2011, **50**, 3760.
- 3 (a) J. Schmidt, J. Choi, A. T. Liu, M. Slusarczyk and G. C. Fu, Science, 2016, 354, 1265; (b) B. Potter, E. K. Edelstein and J. P. Morken, Org. Lett., 2016, 18, 3286; (c) T. Rybak and D. G. Hall, Org. Lett., 2015, 17, 4156; (d) S. C. Matthew, B. W. Glasspoole, P. Eisenberger and C. M. Crudden, J. Am. Chem. Soc., 2014, 136, 5828; (e) A. W. Buesking and J. A. Ellman, Chem. Sci., 2014, 5, 1983; (f) D. Imao, B. W. Glasspoole, V. S. Laberge and C. M. Crudden, J. Am. Chem. Soc., 2009, 131, 5024.
- 4 Stoichiometric reagents: J. M. Brown and B. N. Nguyen, Stereoselective hydroboration and diboration of carboncarbon double bonds, Georg Thieme Verlag, 2011, vol. 1, p.
- 5 For a review on recent advances in the preparation and application of allylboron species, see: C. Diner and K. J. Szabó, *J. Am. Chem. Soc.*, 2017, **139**, 2.
- 6 Enantioselective allyl- and arylborylation: (a) H. M. Nelson, B. D. Williams, J. Miró and F. D. Toste, J. Am. Chem. Soc., 2015, 137, 3213; (b) T. Jia, P. Cao, B. Wang, Y. Lou, X. Yin, M. Wang and J. Liao, J. Am. Chem. Soc., 2015, 137, 13760.
- 7 Enantioselective allylic substitution with diborylmethane: (a) Y. Shi and A. H. Hoveyda, *Angew. Chem., Int. Ed.*, 2016, 55, 3455; (b) M. Zhan, R.-Z. Li, Z.-D. Mou, C.-G. Cao, J. Liu, Y.-W. Chen and D. Niu, *ACS Catal.*, 2016, 6, 3381.
- 8 Enantioselective aminoborylation: (a) D. Nishikawa, K. Hirano and M. Miura, *Org. Lett.*, 2016, **18**, 4856; (b) N. Matsuda, K. Hirano, T. Satoh and M. Miura, *J. Am. Chem. Soc.*, 2013, **135**, 4934; (c) K. Hong and J. P. Morken, *J. Am. Chem. Soc.*, 2013, **135**, 9252.
- Enantioselective B-H bond insertion: (a) D. Chen, X. Zhang,
   W.-Y. Qi, B. Xu and M.-H. Xu, J. Am. Chem. Soc., 2015, 137,
   5268; (b) Q.-Q. Cheng, S.-F. Zhu, Y.-Z. Zhang, X.-L. Xie and
   Q.-L. Zhou, J. Am. Chem. Soc., 2013, 135, 14094.
- 10 Enantioselective C-H borylation: (a) J. He, Q. Shao, Q. Wu and J.-Q. Yu, J. Am. Chem. Soc., 2017, 139, 3344; (b) L.-L. Li, Z.-L. Tao, Z.-Y. Han and L.-Z. Gong, Org. Lett., 2017, 19, 102.
- 11 Enantioselective conjugate borylation: (a) Q. Jiang, T. Guo and Z. Yu, J. Org. Chem., 2017, 82, 1951; (b) C. Jarava-Barrera, A. Parra, A. López, F. Cruz-Acosta, D. Collado-Sanz, D. J. Cárdenas and M. Tortosa, ACS Catal., 2016, 6, 442; (c) J.-B. Xie, S. Lin, S. Qiao and G. Li, Org. Lett., 2016, 18, 3926; (d) Z. Niu, J. Chen, Z. Chen, M. Ma, C. Song and Y. Ma, J. Org. Chem., 2015, 80, 602; (e) H. Wu, J. M. Garcia, F. Haeffner, S. Radomkit, A. R. Zhugralin and A. H. Hoveyda, J. Am. Chem. Soc., 2015, 137, 10585; (f) S. Radomkit and A. H. Hoveyda, Angew. Chem., Int. Ed., 2014, 53, 3387; (g) Y. Luo, I. D. Roy, A. G. E. Madec and H. W. Lam, Angew. Chem., Int. Ed., 2014, 53, 4186.
- 12 Enantioselective conjunctive cross-coupling: (a)
  G. J. Lovinger, M. D. Aparece and J. P. Morken, J. Am. Chem. Soc., 2017, 139, 3153; (b) L. Zhang, G. J. Lovinger, E. K. Edelstein, A. A. Szymaniak, M. P. Chierchia and J. P. Morken, Science, 2016, 351, 70.
- 13 Enantioselective diboration: (a) L. Fang, L. Yan, F. Haeffner and J. P. Morken, *J. Am. Chem. Soc.*, 2016, **138**, 2508; (b) Z. Yu,

**Edge Article** 

R. J. Ely and J. P. Morken, *Angew. Chem., Int. Ed.*, 2014, 53, 9632; (c) J. R. Coombs, F. Haeffner, L. T. Kliman and J. P. Morken, *J. Am. Chem. Soc.*, 2013, 135, 11222; (d) G. E. Ferris, K. Hong, I. A. Roundtree and J. P. Morken, *J. Am. Chem. Soc.*, 2013, 135, 2501.

- 14 Stereospecific catalytic Miyaura borylation: (a) Q. Zhou, H. D. Srinivas, S. Zhang and M. P. Watson, J. Am. Chem. Soc., 2016, 138, 11989; (b) C. H. Basch, K. M. Cobb and M. P. Watson, Org. Lett., 2016, 18, 136.
- 15 Enantioselective nucleophilic addition to aldehyde: (a) M. V. Joannou, B. S. Moyer and S. J. Meek, J. Am. Chem. Soc., 2015, 137, 6176; (b) K. Kubota, E. Yamamoto and H. Ito, J. Am. Chem. Soc., 2015, 137, 420.
- 16 Group-selective cross-coupling of diboranes: (*a*) S. N. Mlynarski, C. H. Schuster and J. P. Morken, *Nature*, 2014, **505**, 386; (*b*) C. Sun, B. Potter and J. P. Morken, *J. Am. Chem. Soc.*, 2014, **136**, 6534; (*c*) R. D. Dewhurst and T. B. Marder, *Nat. Chem.*, 2014, **6**, 279.
- 17 Enantioselective homologation of organoboron *via* lithiation-borylation: (a) A. Varela, L. K. B. Garve, D. Leonori and V. K. Aggarwal, *Angew. Chem., Int. Ed.*, 2017, **56**, 2127; (b) D. J. Blair, D. Tanini, J. M. Bateman, H. K. Scott, E. L. Myers and V. K. Aggarwal, *Chem. Sci.*, 2017, **8**, 2898; (c) For a review on lithiation-borylation methodology and its application in synthesis, see: D. Leonori and V. K. Aggarwal, *Acc. Chem. Res.*, 2014, 47, 3174; (d) For related computational studies, see: S. Essafi, S. Tomasi, V. K. Aggarwal and J. N. Harvey, *J. Org. Chem.*, 2014, **79**, 12148.
- 18 From vinylboronates: (a) J. T. Han, W. J. Jang, N. Kim and J. Yun, J. Am. Chem. Soc., 2016, 138, 15146; (b)
  D. Nishikawa, K. Hirano and M. Miura, J. Am. Chem. Soc., 2015, 137, 15620; (c) J. J. Verendel, O. Pamies, M. Dieguez and P. G. Andersson, Chem. Rev., 2014, 114, 2130.
- 19 Examples of CAHB by copper catalysts (a-i), cobalt catalysts (k-m), iron catalysts (n), iridium catalysts (o-p), and rhodium catalysts (q-r): (a) M. G. Ceinos, A. Parra, V. M. Heras and M. Tortosa, Angew. Chem., Int. Ed., 2016, 55, 6969; (b) Y. Xi and J. F. Hartwig, J. Am. Chem. Soc., 2016, 138, 6703; (c) D. Wang, P. Cao, B. Wang, T. Jia, Y. Lou, M. Wang and J. Liao, Org. Lett., 2015, 17, 2420; (d) A. Parra, L. Amenós, M. Guisán-Ceinos, A. López, J. L. G. Ruano and M. Tortosa, J. Am. Chem. Soc., 2014, 136, 15833; (e) Z.-T. He, Y.-S. Zhao, P. Tian, C.-C. Wang, H.-Q. Dong and G.-Q. Lin, Org. Lett., 2014, 16, 1426; (f) X. Feng, H. Jeon and J. Yun, Angew. Chem., Int. Ed., 2013, 52, 3989; (g) D. Noh, S. K. Yoon, J. Won, J. Y. Lee and J. Yun, Chem.-Asian J., 2011, 6, 1967; (h) Y. Sasaki, C. Zhong, M. Sawamura and H. Ito, J. Am. Chem. Soc., 2010, 132, 1226; (i) D. Noh, H. Chea, J. Ju and J. Yun, Angew. Chem., Int. Ed., 2009, 48, 6062; (j) R. Corberán, N. W. Mszar and A. H. Hoveyda, Angew. Chem., Int. Ed., 2011, 50, 7079; (k) H. Zhang and Z. Lu, ACS Catal., 2016, 6, 6596; (l) L. Zhang, Z. Zuo, X. Wan and Z. Huang, J. Am. Chem. Soc., 2014, 136, 15501; (m) J. Chen, T. Xi, X. Ren, B. Cheng, J. Guo and Z. Lu, Org. Chem. Front., 2014, 1, 1306; (n) J. Chen, T. Xi and Z. Lu, Org. Lett., 2014, 16, 6452; (o) M. Magre, M. Biosca, O. Pàmies and

- M. Diéguez, ChemCatChem, 2015, 7, 114; (p) C. Mazet and D. Gérard, Chem. Commun., 2011, 47, 298; (q) N. Hu, G. Zhao, Y. Zhang, X. Liu, G. Li and W. Tang, J. Am. Chem. Soc., 2015, 137, 6746; (r) S. A. Moteki, K. Toyama, Z. Liu, J. Ma, A. E. Holmes and J. M. Takacs, Chem. Commun., 2012, 48, 263.
- (a) S. M. Smith, G. L. Hoang, R. Pal, M. O. B. Khaled, L. S. W. Pelter, X. C. Zeng and J. M. Takacs, *Chem. Commun.*, 2012, 48, 12180; (b) S. M. Smith, M. Uteuliyev and J. M. Takacs, *Chem. Commun.*, 2011, 47, 7812; (c) S. M. Smith and J. M. Takacs, *Org. Lett.*, 2010, 12, 4612; (d) S. M. Smith and J. M. Takacs, *J. Am. Chem. Soc.*, 2010, 132, 1740; (e) S. M. Smith, N. C. Thacker and J. M. Takacs, *J. Am. Chem. Soc.*, 2008, 130, 3734.
- 21 V. M. Shoba, N. C. Thacker, A. J. Bochat and J. M. Takacs, *Angew. Chem., Int. Ed.*, 2016, 55, 1465.
- 22 G. L. Hoang, Z.-D. Yang, S. M. Smith, R. Pal, J. L. Miska, D. E. Pérez, L. S. W. Pelter, X. C. Zeng and J. M. Takacs, *Org. Lett.*, 2015, 17, 940.
- 23 The symmetry in substrate 3 was also exploited in a DFT study of potential catalytic cycle for the reaction; see: Z.-D. Yang, R. Pal, G. L. Hoang, X. C. Zeng and J. M. Takacs, *ACS Catal.*, 2014, 4, 763.
- 24 R. Peters, P. Waldmeier and A. Joncour, *Org. Process Res. Dev.*, 2005, **9**, 508.
- 25 When reacted separately substrate **9** requires higher catalyst loading for complete consumption as stated above (*i.e.*, 2.0% Rh for **9** *versus* 0.5% for **5d**). However, the direct competition of **9** and **5d** for limiting pinBH (equivalent amounts of each) affords only a modest excess of recovered **9** (74% recovered) relative to recovered **5d** (61% recovered). The results suggest that **5d** is consumed only slightly faster than **9** by active catalyst. A relatively slow conversion of the rhodium precatalyst to active catalyst by **9** in the absence of **5d** could account of the conflicting observations; see ref. 22 for a similar example.
- 26 D. L. Sandrock, L. Jean-Gerard, C. Chen, S. D. Dreher and G. A. Molander, J. Am. Chem. Soc., 2010, 132, 17108.
- 27 (a) T. Awano, T. Ohmura and M. Suginome, J. Am. Chem. Soc., 2011, 133, 20738; (b) T. Ohmura, T. Awano and M. Suginome, J. Am. Chem. Soc., 2010, 132, 13191.
- 28 (a) J. C. H. Lee, H.-Y. Sun and D. G. Hall, J. Org. Chem., 2015, 80, 7134; (b) J. C. Lee, R. McDonald and D. G. Hall, Nat. Chem., 2011, 3, 894.
- 29 L. Li, S. Zhao, A. Joshi-Pangu, M. Diane and M. R. Biscoe, J. Am. Chem. Soc., 2014, 136, 14027.
- 30 M. Daini and M. Suginome, J. Am. Chem. Soc., 2011, 133, 4758.
- 31 T. P. Blaisdell and J. P. Morken, *J. Am. Chem. Soc.*, 2015, **137**, 8712.
- 32 A. J. J. Lennox and G. Lloyd-Jones, *Angew. Chem., Int. Ed.*, 2012, **51**, 9385.
- 33 N. C. Bruno, M. T. Tudge and S. L. Buchwald, *Chem. Sci.*, 2013, 4, 916.
- 34 Authentic (*R*)-21c was prepared *via* enolate allylation using the Evans chiral auxiliary: *in situ* half reduction to aldehyde followed by Wadsworth-Horner-Emmons

olefination, Pd/C hydrogenation and transamidation with

morpholine is described in the accompanying ESI.†

**Chemical Science** 

- 35 (a) C. H. Larsen, B. H. Ridway, J. T. Shaw, D. M. Smith and K. A. Woerpel, J. Am. Chem. Soc., 2005, 127, 10879; (b)
  M. Aquino, S. Cardani, G. Fronza, C. Fuganti, R. P. Fernandez and A. Tagliani, Tetrahedron, 1991, 47, 7887.
- 36 (a) E. Hupe, I. Marek and P. Knochel, *Org. Lett.*, 2002, **4**, 2681, see also; (b) P. Y. Chavant, F. Lhermitte and M. Vaultier, *Synlett*, 1993, 519; (c) D. S. Matteson and G. Y. Kim, *Org. Lett.*, 2002, **4**, 2153; (d) V. Bagutski, T. G. Elford and V. K. Aggarwal, *Angew. Chem., Int. Ed.*, 2011, **50**, 1080.