Chemical Science

EDGE ARTICLE

Check for updates

Cite this: Chem. Sci., 2017, 8, 6852

Received 4th May 2017 Accepted 26th July 2017 DOI: 10.1039/c7sc02018j

rsc.li/chemical-science

Formation and ligand-based reductive chemistry of bridged bis-alkylidene scandium(III) complexes†

Wangyang Ma, ២^a Chao Yu, ២^a Yue Chi, ២^a Tianyang Chen, ២^a Lianjun Wang, ២^{ac} Jianhao Yin, ២^a Baosheng Wei, ២^a Ling Xu, ២^a Wen-Xiong Zhang ២^{*ab} and Zhenfeng Xi ២^a

The chemistry of rare-earth carbene and alkylidene complexes including their synthesis, structure and reaction is a challenging issue because of their high reactivity (or instability) and the lack of synthetic methods. In this work, we report the first synthesis of the bridged bis-alkylidene complexes which feature a 2-butene-1,1,4,4-tetraanion and four $Sc-C(sp^3)$ bonds by the reaction of 1,4-dilithio-1,3-butadienes with $ScCl_3$. This reaction proceeds *via* two key intermediates: an isolable scandacyclopentadiene and a proposed scandacyclopropene. The scandacyclopentadiene undergoes β , $\beta'-C-C$ bond cleavage to generate the scandacyclopropene, which then dimerizes to afford the bridged bis-alkylidene complex *via* a cooperative double metathesis reaction. Reaction chemistry study of the bridged bis-alkylidene complex reveals their ligand-based reduction reactivity towards different oxidants such as hexachloroethane, disulfide and cyclooctatetraene.

Transition metal carbene and alkylidene complexes have been extensively studied because of their importance in organometallic chemistry, coordination chemistry and synthetic organic chemistry.1 In contrast, rare-earth metal carbene and alkylidene complexes are very limited mainly due to the energy mismatch between the rare-earth metals and ligand orbitals.²⁻¹² Since the rare-earth alkylidene complex was first postulated in 1979,3 pioneering works have been made to isolate and characterize it. Some pincer-like rare-earth alkylidene complexes have been reported independently by Cavell,4 Liddle,5 and Mézailles.6 Very recently, Cui et al. reported the lutetium methanediide-alkyl complexes,⁷ and Chen *et al.* reported the non-pincer-type mononuclear scandium alkylidene complexes.8 Furthermore, rare-earth methylidene complexes were also stabilized by chloride bridges9 or Lewis-acids such as AlMe₃.10 Interestingly, mixed methyl/methylidene complexes11 and cubane-like methylidene complexes¹² have been reported. Despite these recent advances, the chemistry of rare-earth alkylidene complexes is still in its infancy, and the bridged bis-alkylidene complex remains scarce.

Reductive reaction of rare-earth organometallic compounds is a fundamental process in organometallic chemistry and coordination chemistry.¹³ Rare earth metal complexes (Ce, Sm, Eu and Yb) supported by redox-inert ligands tend to perform a single electron redox process. The utilization of redox-active ligands at the rare earth metal centers is an alternative strategy for affording multi-electron redox reactivity.¹⁴ Ligandbased reductive chemistry of trivalent rare-earth organometallic compounds has received much attention. Evans and coworkers have made great progress in studying the reductive reactivity of $(C_5Me_5)_3Ln$ (Ln = La, Nd, Sm, *etc.*) and provided a wide variety of new reductive chemistry for rare earth metals.^{13a,15}

Herein, we report the first synthesis of the bridged bisalkylidene complex featuring a 2-butene-1,1,4,4-tetraanion and four Sc–C(sp³) bonds from 1,4-dilithio-1,3-butadienes and ScCl₃. This reaction proceeds *via* two key intermediates: scandacyclopentadiene^{16,17} and scandacyclopropene.^{18,19} DFT calculations indicate that the dimerization of scandacyclopropenes *via* the cooperative double metathesis is the key factor for the formation of the bridged bis-alkylidene complex. Interestingly, the bridged bis-alkylidene scandium(III) complex shows unexpected ligand-based two-electron or four-electron reduction reactivity towards different oxidants such as hexachloroethane, disulfide and cyclooctatetraene.

Silyl-substituted 1,4-dilithio-1,3-butadienes **1a–c** were readily prepared according to our previous procedure.²⁰ When the 1 : 1 reaction of **1a** and solvated ScCl₃ in THF was conducted at -20 °C, the light yellow crystalline complex **2a** could be isolated exclusively in 65% yield (Scheme 1). An X-ray analysis of **2a**



View Article Online

View Journal | View Issue

[&]quot;Beijing National Laboratory for Molecular Sciences, MOE Key Laboratory of Bioorganic Chemistry and Molecular Engineering, College of Chemistry, Peking University, Beijing 100871, China. E-mail: wx_zhang@pku.edu.cn

^bState Key Laboratory of Elemento-Organic Chemistry, Nankai University, Tianjin, 300071, China

^cSchool of Chemistry and Chemical Engineering, Hunan Institute of Engineering, Xiangtan, 411104, China

[†] Electronic supplementary information (ESI) available. CCDC 1555495, 1504242, 1519012 and 1539530. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7sc02018j



Scheme 1 Synthesis of scandacyclopentadiene 2a and bridged bisalkylidene scandium(III) complexes 3a-c.



Fig. 1 Molecular structure of complex 2a with thermal ellipsoids at 30% probability. H atoms are omitted for clarity.

revealed that it is a LiCl-ligated scandacyclopentadiene (Fig. 1). The Sc(m) center adopts a distorted octahedral fashion bonded with two C(sp²) atoms, two chlorides and two THF molecules. The C1–C2 (1.348(4) Å) and C3–C4 (1.376(4) Å) bond lengths are within the range of standard C=C bond lengths, and the C2–C3 bond length (1.520(3) Å) indicates a typical C–C single bond. These data of bond lengths clearly show the butadienyl dianionic structure in **2a**.

Complex 2a is sensitive to air and moisture but stable under dry N₂ atmosphere. In the ¹H NMR spectrum in THF- d_8 , a singlet at -0.38 ppm was observed and assigned to the proton resonance of TMS groups. Two β -C(sp²) atoms (C2 and C3) displayed a singlet at 167.6 ppm in the ¹³C NMR spectrum, while two α -C(sp²) atoms (C1 and C4) showed a broad peak at 203.8 ppm, probably due to the coupling with scandium (nuclear spin quantum number I = 7/2). The ¹H NMR spectrum of 2a in THF- d_8 showed no obvious change for 2 weeks at room temperature. However, when the THF- d_8 solution of 2a was heated at 45 °C for 3 h or 80 °C for 10 min, the TMS proton resonance at -0.38 ppm completely disappeared in the ¹H NMR spectrum, and two new singlets integrated to the same number of protons appeared at -0.23 ppm and 0.20 ppm (see ESI[†] for more details). The singlet at 0.20 ppm was assigned to the TMS proton resonance of PhC=CTMS by comparison with its standard spectrum. The GC retention time and molecular ion peak (m/z = 174) detected by GC-MS are also consistent with those of the standard sample of PhC=CTMS. The other new singlet at -0.23 ppm was assigned to the TMS groups of a new complex **3a**, which was obtained in almost quantitative yield by thermolysis of **2a**. Furthermore, we found that the synthesis of **3a** does not require isolation of **2a** as the starting material. **3a** could be conveniently prepared by the reaction of **1a** with solvated ScCl₃ in THF solution at 80 °C for 3 h. Similarly, **3b** and **3c** could be prepared from the corresponding 1,4-dilithio-1,3-butadienes and ScCl₃ (Scheme 1).

An X-ray analysis of 3a reveals it is a bridged bis-alkylidene complex and adopts a dimeric ate complex via µ2-chloride bridges (Fig. 2). One scandium center (e.g. Sc1) is bonded with two carbon atoms and two terminal chlorides, while the other one (e.g. Sc2) is bonded with two carbon atoms, two bridged chlorides and one THF. The Sc1-Sc2 distance (3.1366(9) Å) is the shortest length found in the literature, which is notably shorter than those in dinuclear scandium hydride complexes (3.20-3.40 Å).²¹ Two lithium atoms act as counterions, and each lithium atom forms a distorted tetrahedron surrounded by four THF molecules. The bond lengths of C1-C2 (1.468(4) Å) and C3-C4 (1.465(5) Å) in 3a are significantly longer than those in 2a [C1-C2, 1.348(4) Å; C(3)-C(4), 1.376(4) Å]. The bond length of C2–C3 (1.430(4) Å) in **3a** is significantly shorter than the corresponding C2-C3 (1.520(3) Å) in 2a. Thus, the bond lengths in the C1-C2-C3-C4 moiety in 3a are averaged and are not the classical bond lengths of C-C single and double bonds. These results show that 3a has a highly delocalized structure with a tetraanionic ligand. Most importantly, these results are in striking contrast with what was observed previously for the transmetalation reactions of 1,4-dilithio-1,3-butadienes with metal salts which gave 1,3-butadiene-1,4-dianion complexes.20

The formation of the asymmetric unit in **3a** from two molecules of **2a** along with elimination of two alkynes is a very interesting process and intrigued us to explore the reaction mechanism. The crossover reaction between **2a** and **2a-D**₁₀ was carried out. When the reaction mixture was quenched with H₂O, **4a**, **4a-D**₅, and **4a-D**₁₀ could all be detected by HRMS (Scheme 2). This result unambiguously reveals that the 2-butene-1,1,4,4-tetraanion moiety in **3a** should be originated from two



Fig. 2 Molecular structure of complex 3a with thermal ellipsoids at 30% probability. H atoms and two $[Li(THF)_4]^+$ counterions are omitted for clarity.



molecules of scandacyclopentadienes instead of a simple reduction of a diene moiety in one scandacyclopentadiene. Thus, the crossover experiment excludes two possible pathways involving two eliminated alkynes from the same scandacyclopentadienes: (i) the cooperative intermolecular redox process, and (ii) stepwise intermolecular redox *via* the scandacyclopropene process (see ESI† for more details).

Based on the above information, we proposed a mechanism involving the scandacyclopropene intermediate. For a better understanding of the formation of **3a**, DFT calculations were carried out using Gaussian 09 (Fig. 3).²² We chose the LiCl-free scandacyclopentadiene **IM1** as a starting model compound and the THF-ligated monomer **3a-M** as a targeted compound for simplicity.²³ The structures of all of the minima and transition states were optimized at the B3LYP²⁴/LANL2DZ (for Sc)/6-31+G* (for other elements) level in the gas phase. The effect of the solvent was examined by performing single-point self-consistent reaction field (SCRF) calculations based on the polarizable continuum model (PCM) for gas-phase optimized

structures. Scandacyclopentadiene IM1 will undergo β , β' -C-C bond cleavage to generate scandacyclopropene IM2 by release of one equiv. of alkyne. The β , β' -C-C bond cleavage from **IM1** to IM2 is the critical step with the highest energy barrier of 13.3 kcal mol^{-1} in the solution phase, which means that IM1 is isolable. Metallacyclopropenes, as an important class of reactive intermediate, have been isolated and characterized in transition and main group organometallic chemistry.^{18,19} The metallacyclopropene, e.g. aluminacyclopropene, can undergo dimerization to give a 1,4-dialuminacyclohexadiene.25 In contrast, rare-earth metallacyclopropenes are unknown. IM2 is the first optimized structure of a rare-earth metallacyclopropene by DFT calculations. Next, we tried to optimize the dimeric structure of IM3' which is similar to 1,4-dialuminacyclohexadiene. However, the optimization of the structure of IM3' to a local energy minimum failed, probably because of its high energy and instability. Rather than giving IM3', a new intermediate, IM3, resulting from two IM2 species approaching each other via the weak Sc-C interaction, was optimized to a local minimal energy, 2.7 kcal mol^{-1} lower than IM2. Surprisingly, a cooperative double metathesis of IM3 gives 3a-M via the transition state TS2. In TS2, two scandacyclopropene rings adopt a triangular prism geometry, in which each Sc atom is coordinated to another carbon neighbouring TMS group. This geometry of TS2 could also explain the selectivity of C(Ph)-C(Ph) coupling.

The structure of **3a** features the 2-butene-1,1,4,4-tetraanion moiety and thus we thought it could be oxidized to generate the diene moiety in **2a**, as illustrated in Scheme 3. As we expected, **2a** was generated by treatment with two equivalents of hexachloroethane as an oxidant (Scheme 3a). This reaction resulted in the formation of ScCl₃ which can be characterized as a ScCl₃(THF)₃ adduct by X-ray analysis, along with two equivalents of tetrachloroethylene which were identified using the ¹³C NMR spectrum and GC-MS. When four equivalents of



Fig. 3 DFT calculated energy profiles of related intermediates and transition-states in the generation of **3a-M** (red lines: broken bonds; blue lines: newly formed bonds).



Scheme 3 Ligand-based reduction reactivity of 3a towards different oxidants.

hexachloroethane were used and the reaction mixture was heated at 80 °C, **3a** was transformed to PhC=CTMS and ScCl₃ (Scheme 3b). Furthermore, when disulfide **5** served as an oxidant,²⁶ the reaction of **3a** with **5** provided complex **6** (see ESI† for the X-ray structure of **6**, Scheme 3c) along with the formation of PhC=CTMS. When **3a** was treated with cyclooctatetraene at 80 °C, cyclooctatetraene was reduced to the cyclooctatetraene dianion. The corresponding complex **7** (see ESI† for the X-ray structure of **7**, Scheme 3d) could be isolated after being recrystallized in DME (DME = 1,2-dimethoxyethane) in high yields along with the formation of PhC=CTMS. These results clearly show that the bridged bis-alkylidene scandium(m) complex **3a** can act as an efficient two-electron or four-electron reductant.

Conclusions

In summary, we have developed a simple and efficient synthetic method for the first series of well-defined bridged bis-alkylidene scandium(m) complexes from 1,4-dilithio-1,3-butadienes and

ScCl₃. This reaction proceeds *via* two key intermediates: an isolable scandacyclopentadiene and a proposed scandacyclopropene. A mechanistic pathway of C–C bond recombination through the dimerization of scandacyclopropene intermediates is elucidated well by DFT calculations. Bridged bis-alkylidene scandium(III) complexes are found to show ligand-based reduction reactivity towards different kinds of oxidant. Further reaction chemistry of bis-alkylidene scandium(III) complexes and characterization of scandacyclopropenes are in progress.

Conflicts of interest

There are no conflicts of interest to declare.

Acknowledgements

This work was supported by the Natural Science Foundation of China (no. 21572005, 21372014).

Notes and references

- (a) R. R. Schrock, Angew. Chem., Int. Ed., 2006, 45, 3748–3759;
 (b) R. H. Grubbs, Angew. Chem., Int. Ed., 2006, 45, 3760–3765;
 (c) D. Bourissou, O. Guerret, F. P. Gabbai and G. Bertrand, Chem. Rev., 2000, 100, 39–91;
 (d) R. R. Schrock, J. Am. Chem. Soc., 1974, 96, 6796–6797;
 (e) E. O. Fischer and A. Maasböl, Angew. Chem., Int. Ed., 1964, 3, 580–581.
- 2 (a) P. L. Arnold and I. J. Casely, Chem. Rev., 2009, 109, 3599–3611; (b) P. L. Arnold and S. Pearson, Coord. Chem. Rev., 2007, 251, 596–609; (c) S. T. Liddle, I. S. Edworthy and P. L. Arnold, Chem. Soc. Rev., 2007, 36, 1732–1744; (d) P. L. Arnold and S. T. Liddle, Chem. Commun., 2006, 3959–3971; (e) G. R. Giesbrecht and J. C. Gordon, Dalton Trans., 2004, 2387–2393; (f) R. G. Cavell, R. P. K. Babu and K. Aparna, J. Organomet. Chem., 2001, 617–618, 158–169.
- 3 H. Schumann and J. Müller, *J. Organomet. Chem.*, 1979, **169**, C1–C4.
- 4 K. Aparna, M. Ferguson and R. G. Cavell, *J. Am. Chem. Soc.*, 2000, **122**, 726–727.
- 5 D. P. Mills, L. Soutar, W. Lewis, A. J. Blake and S. T. Liddle, *J. Am. Chem. Soc.*, 2010, **132**, 14379–14381.
- 6 M. Fustier, X. F. Le Goff, P. Le Floch and N. Mézailles, J. Am. Chem. Soc., 2010, 132, 13108–13110.
- 7 S. Li, M. Wang, B. Liu, L. Li, J. Cheng, C. Wu, D. Liu, J. Liu and D. Cui, *Chem.-Eur. J.*, 2014, **20**, 15493–15498.
- 8 (a) W. Mao, L. Xiang, C. A. Lamsfus, L. Maron, X. Leng and Y. Chen, J. Am. Chem. Soc., 2017, 139, 1081–1084; (b) C. Wang, J. Zhou, X. Zhao, L. Maron, X. Leng and Y. Chen, *Chem.-Eur. J.*, 2016, 22, 1258–1261; (c) J. Zhou, T. Li, L. Maron, X. Leng and Y. Chen, *Organometallics*, 2015, 34, 470–476.
- 9 H. M. Dietrich, K. W. Törnroos and R. Anwander, J. Am. Chem. Soc., 2006, **128**, 9298–9299.
- 10 (a) W. Huang, C. T. Carver and P. L. Diaconescu, *Inorg. Chem.*, 2011, 50, 978–984; (b) R. Litlabø, M. Zimmermann, K. Saliu, J. Takats, K. W. Törnroos and R. Anwander,

Angew. Chem., Int. Ed., 2008, 47, 9560–9564; (c) M. Zimmermann, J. Takats, G. Kiel, K. W. Törnroos and R. Anwander, Chem. Commun., 2008, 612–614; (d) J. Scott, H. Fan, B. F. Wicker, A. R. Fout, M.-H. Baik and D. J. Mindiola, J. Am. Chem. Soc., 2008, 130, 14438–14439.

- 11 J. Hong, L. Zhang, X. Yu, M. Li, Z. Zhang, P. Zheng, M. Nishiura, Z. Hou and X. Zhou, *Chem.-Eur. J.*, 2011, 17, 2130–2137.
- 12 (a) W.-X. Zhang, Z. Wang, M. Nishiura, Z. Xi and Z. Hou, J. Am. Chem. Soc., 2011, 133, 5712–5715; (b) T. Li, M. Nishiura, J. Cheng, Y. Li and Z. Hou, Chem.-Eur. J., 2012, 18, 15079–15085.
- 13 For a selected review and book see: (a) W. J. Evans, *Inorg. Chem.*, 2007, **46**, 3435–3449; (b) R. H. Crabtree, *The Organometallic Chemistry of the Transition Metals*, Wiley Interscience, New York, 2014.
- 14 Selected examples of multielectron redox reactions of rareearth metal complexes: (a) Y. Lv, C. E. Kefalidis, J. Zhou, L. Maron, X. Leng and Y. Chen, J. Am. Chem. Soc., 2013, 135, 14784–14796; (b) C. Camp, V. Guidal, B. Biswas, J. Pécaut, L. Dubois and M. Mazzanti, Chem. Sci., 2012, 3, 2433–2448; (c) W. Huang, S. I. Khan and P. L. Diaconescu, J. Am. Chem. Soc., 2011, 133, 10410–10413.
- 15 Selected reviews on sterically induced reduction (SIR) of lanthanides: (a) W. J. Evans and B. L. Davis, Chem. Rev., 2002, 102, 2119–2136; (b) W. J. Evans, J. Organomet. Chem., 2002, 647, 2–11; (c) W. J. Evans, Coord. Chem. Rev., 2000, 206–207, 263–283.
- 16 Selected reviews of metallacyclopentadienes: (a) W. Ma, C. Yu, T. Chen, L. Xu, W.-X. Zhang and Z. Xi, Chem. Soc. Rev., 2017, 46, 1160–1192; (b) X. Yan and C. Xi, Acc. Chem. Res., 2015, 48, 935–946; (c) T. Takahashi and Y. Li, in Titanium and Zirconium in Organic Synthesis, ed. I. Marek, Wiley-VCH, Weinheim, 2002, ch. 2.
- 17 Examples of rare-earth metallacyclopentadienes: (a) L. Xu,
 Y. Wang, Y.-C. Wang, Z. Wang, W.-X. Zhang and Z. Xi,
 Organometallics, 2016, 35, 5–8; (b) L. Xu, J. Wei,
 W.-X. Zhang and Z. Xi, Chem.-Eur. J., 2015, 21, 1586015866; (c) L. Xu, Y.-C. Wang, J. Wei, Y. Wang, Z. Wang,
 W.-X. Zhang and Z. Xi, Chem.-Eur. J., 2015, 21, 6686–6689.

- 18 Selected reviews on metallacyclopropenes: (a) K. D. J. Parker and M. D. Fryzuk, Organometallics, 2015, 34, 2037–2047; (b) U. Rosenthal, V. V. Burlakov, P. Arndt, W. Baumann and A. Spannenberg, Organometallics, 2003, 22, 884–900.
- 19 Selected examples of metallacyclopropenes: (a) L. Zhang,
 G. Hou, G. Zi, W. Ding and M. D. Walter, J. Am. Chem. Soc., 2016, 138, 5130–5142; (b) E. D. Jemmis, S. Roy,
 V. V. Burlakov, H. Jiao, M. Klahn, S. Hansen and
 U. Rosenthal, Organometallics, 2010, 29, 76–81.
- 20 (a) Z. Xi, Acc. Chem. Res., 2010, 43, 1342–1351; (b) W.-X. Zhang and Z. Xi, Org. Chem. Front., 2014, 1, 1132–1139.
- 21 (a) N. R. Halcovitch and M. D. Fryzuk, Organometallics, 2013,
 32, 5705–5708; (b) J. Oyamada, M. Nishiura and Z. Hou, Angew. Chem., Int. Ed., 2011, 50, 10720–10723; (c)
 J. R. Hagadorn and J. Arnold, Organometallics, 1996, 15, 984–991; (d) Y. Mu, W. E. Piers, D. C. MacQuarrie, M. J. Zaworotko and V. G. Young Jr, Organometallics, 1996, 15, 2720–2726.
- 22 M. J. Frisch, *et al.*, Gaussian 09 (Revision C.01), Gaussian, Inc., Wallingford CT, 2010, for full reference, see ESI.[†]
- 23 The Sc–O interaction is much stronger than the Sc–Cl interaction because of the oxophilicity of rare-earth elements. In THF, LiCl in 2a will be easily replaced by THF to yield a LiCl-free complex **IM1**. Based on the effects of the THF atmosphere, we excluded LiCl from the calculation. In THF, 3a tends to be a monomer due to the solvent coordination interaction. Furthermore, the bond lengths and angles of the calculated monomeric structure are similar to those of the crystal dimeric structure. Thus, we think the calculation of monomer **3a-M** is enough to describe the reaction pathway.
- 24 (a) A. D. Becke, J. Chem. Phys., 1993, 98, 5648-5652; (b)
 C. Lee, W. Yang and R. G. Parr, Phys. Rev. B: Condens. Matter Mater. Phys., 1988, 37, 785-789.
- 25 (a) C. Üffing, A. Ecker, R. Köppe, K. Merzweiler and H. Schnöckel, *Chem.-Eur. J.*, 1998, 4, 2142–2147; (b)
 H. Hoberg, V. Gotor, A. Milchereit, C. Krüger and J. C. Sekutowski, *Angew. Chem., Int. Ed.*, 1977, 16, 539.
- 26 I. V. Basalov, D. M. Lyubov, G. K. Fukin, A. S. Shavyrin and A. A. Trifonov, Angew. Chem., Int. Ed., 2012, 51, 3444–3447.