

Cite this: *Chem. Sci.*, 2017, 8, 6645

Received 24th June 2017

Accepted 20th July 2017

DOI: 10.1039/c7sc02809a

rsc.li/chemical-science

# Catalytic asymmetric hydroxylative dearomatization of 2-naphthols: synthesis of lacinilene derivatives†

Yu Zhang,<sup>ID</sup> Yuting Liao,<sup>ID</sup> Xiaohua Liu,<sup>ID</sup> \* Xi Xu, Lili Lin<sup>ID</sup> and Xiaoming Feng<sup>ID</sup> \*

An enantioselective hydroxylative dearomatization of 2-naphthols with oxaziridines has been accomplished using a *N,N'*-dioxide–scandium(III) complex catalyst. Various substituted *ortho*-quinols could be obtained in high yields (up to 99%) and enantioselectivities (up to 95 : 5 er). This methodology could be applied in the synthesis of bioactive lacinilenes in a gram-scale reaction. Based on the experimental investigations and previous work, a possible catalytic model was proposed.

## Introduction

Substituted *ortho*-quinols are essential structural motifs in a number of natural products and pharmaceuticals.<sup>1</sup> For instance, chiral lacinilene derivatives (Fig. 1), a series of phytoalexins isolated from cotton plants, have been utilized for inhibiting the growth of cotton bacterial pathogens, such as *Xanthomonas campestris* or *malvacearum*.<sup>2</sup> Studies have showed that the (*S*)-enantiomer of lacinilene C is more active than the (*R*)-enantiomer.<sup>2c</sup> While these biological activities provide a justification for the development of approaches to the synthesis of enantiomerically enriched lacinilene derivatives, novel catalytic enantioselective methods remain limited.<sup>2b,d</sup>

Optically active lacinilene derivatives in nature were proposed to be produced enzymically from the oxidation of dihydroxycadalenes, thus it is of practical interest to discover a catalytic asymmetric oxidative dearomatization route to the synthesis of these cadinanes.<sup>3</sup> Compared with other successful dearomatization events of phenols or naphthols,<sup>4,5</sup> controlling the chemo-, regio- and enantioselectivity of the asymmetric

hydroxylative dearomatization is more difficult,<sup>6</sup> as there might be serious side reactions in the presence of oxidants including overoxidation of alkene functions, competitive *para*-oxidation and homocoupling.<sup>6c,e</sup> Additionally, the *ortho*-quinol product could undergo an unexpected  $\alpha$ -ketol rearrangement, which enhances the difficulty of controlling the reactivity and selectivity.<sup>6a,7</sup> In this respect, only a few reports related to asymmetric hydroxylative dearomatization of phenols or naphthols have been reported. Asymmetric oxidative dearomatization of phenolate mediated by copper–sparteine–dioxide complexes followed a [4 + 2] dimerization cascade, giving bicyclo[2.2.2] octenones as the final products.<sup>6a</sup> Several chiral hypervalent organoiodine compounds were developed for the asymmetric hydroxylative dearomatization of phenols and 1-naphthols.<sup>6b–e</sup> Taking these examples into account, we want to engage in discovering new enantioselective strategies for the synthesis of *ortho*-quinol moieties with improved efficiency and selectivity. Here, we present an efficient asymmetric hydroxylative dearomatization of 2-naphthols catalyzed by a chiral *N,N'*-dioxide–scandium(III) complex catalyst.<sup>8</sup> The process could be applied to the synthesis of various 1-hydroxy-1-alkyl-naphthalen-2-one derivatives including lacinilene C methyl ether and lacinilene D, in high to excellent yields and good enantioselectivities under mild reaction conditions (Scheme 1).




Fig. 1 Representative active lacinilene derivatives bearing *ortho*-quinol structures.


Key Laboratory of Green Chemistry & Technology, Ministry of Education, College of Chemistry, Sichuan University, Chengdu 610064, China. E-mail: liuxh@scu.edu.cn; xmfeng@scu.edu.cn; Fax: +86 28 85418249; Tel: +86 28 85418249

† Electronic supplementary information (ESI) available. CCDC 1536822. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7sc02809a


## Results and discussion

We selected the hydroxylative dearomatization of 1-methylnaphthalen-2-ol **1a** as the model substrate using 3-phenyl-2-tosyl-1,2-oxaziridine **2a** as the oxidant which was proven to be chemoselective as a phase-transfer-catalyst under basic conditions (Table 1).<sup>7a</sup> Initially, the catalytic asymmetric reaction was performed with 10 mol% of chiral *N,N'*-dioxide **1-PiPr<sub>2</sub>**–Sc(OTf)<sub>3</sub> complex in DCM at 30 °C, and the desired product **3a** could be obtained dominantly with 80 : 20 er while the  $\alpha$ -ketol rearrangement byproduct **4a** was isolated in around one-fourth of

**Previous work:** Asymmetric hydroxylative dearomatization of phenols and 1-naphthols



**This work:** Asymmetric hydroxylative dearomatization of 2-naphthols



**Scheme 1** Catalytic asymmetric hydroxylative dearomatization of phenols and naphthols.

**Table 1** Optimization of the reaction conditions<sup>a</sup>

Entry	Metal salt	L*	Yield <sup>b</sup> (%)	Ratio (3a/4a) <sup>c</sup>	er (3a) <sup>c</sup>
1	Sc(OTf) <sub>3</sub>	L-PiPr <sub>2</sub>	96	73 : 27	80 : 20
2	Sc(OTf) <sub>3</sub>	L-PrPr <sub>2</sub>	99	79 : 21	63 : 37
3	Sc(OTf) <sub>3</sub>	L-RaPr <sub>2</sub>	99	75 : 25	53.5 : 46.5
4	Sc(OTf) <sub>3</sub>	L-PiMe <sub>2</sub>	90	>95 : 5	60 : 40
5	Sc(OTf) <sub>3</sub>	L-PiPr <sub>3</sub>	96	89 : 11	73 : 27
6	Sc(NTf <sub>2</sub> ) <sub>3</sub>	L-PiPr <sub>2</sub>	99	>95 : 5	92 : 8
7 <sup>d</sup>	Sc(NTf <sub>2</sub> ) <sub>3</sub>	L-PiPr <sub>2</sub>	99	>95 : 5	95 : 5
8 <sup>e</sup>	Sc(NTf <sub>2</sub> ) <sub>3</sub>	L-PiPr <sub>2</sub>	86	>95 : 5	93.5 : 6.5
9 <sup>d,f</sup>	Sc(NTf <sub>2</sub> ) <sub>3</sub>	L-PiPr <sub>2</sub>	99	>95 : 5	94.5 : 5.5


<sup>a</sup> Unless otherwise noted, the reactions were performed with L\*/Sc(III) (1 : 1, 10 mol%), **1a** (0.10 mmol) and **2a** (2.0 equiv.) in DCM (1.0 mL) under N<sub>2</sub> at 30 °C for 3 h. <sup>b</sup> Isolated yield by silica gel chromatography. <sup>c</sup> Determined by chiral HPLC analysis. <sup>d</sup> 5 mol% catalyst loading at 0 °C. <sup>e</sup> 1 mol% catalyst loading at 0 °C for 4 h. <sup>f</sup> **2a** (1.5 equiv.) was used.

a 96% total yield (Table 1, entry 1). The evaluation of the structure of the *N,N'*-dioxides showed that L-PiPr<sub>2</sub> was the optimal ligand in terms of the enantioselectivity albeit ligand L-PiMe<sub>2</sub> and L-PiPr<sub>3</sub> improved the yield of the desired product **3a** (entries 2–5). Fortunately, changing the counterion of the scandium salt from OTf to NTf<sub>2</sub> could suppress the α-ketol rearrangement, delivering the quinol **3a** in a 99% yield with

92 : 8 er (Table 1, entry 6). Further optimization of the reaction conditions, such as decreasing the temperature and the catalyst loading to 5 mol%, resulted in slightly improved enantioselectivity with maintained efficiency (entry 7). Lowering the catalyst loading to 1 mol% or the amount of the oxidant **2a** decreased either the yield or the selectivity a little (entries 8 and 9). We therefore chose the reaction conditions in Table 1, entry 7 for further studies.

We next explored the substrate scope of 2-naphthols (Table 2). The introduction of bromo or methoxyl groups at the C6-position of 2-naphthols had no obvious effect on the result. The 6-aryl substituted 2-naphthol derivatives **1d–1l** tethering various electron-donating and electron-withdrawing substituents could undergo the transformations smoothly, providing the products **3d–3l** in 95–99% yield and 93.5 : 6.5–95 : 5 er. It was noteworthy that 6-alkenyl and alkynyl substituted substrates **1m–1q** were compatible with the reaction conditions, and no aminohydroxylation of the unsaturated carbon–carbon bond occurred, giving the hydroxylative dearomatization products **3m–3q** in good to excellent yields and enantioselectivities.<sup>9</sup>

**Table 2** Substrate scope for 2-naphthols<sup>a</sup>



<sup>a</sup> Reaction conditions: the same as entry 7 in Table 1. <sup>b</sup> 10 mol% catalyst loading. <sup>c</sup> L-PiEt<sub>2</sub>-Sc(OTf)<sub>3</sub> (1 : 1, 5 mol%). <sup>d</sup> Total yield of **3ab** and **4ab**, **3ab/4ab** = 87 : 13.

Additionally, 6-alkyl substituted 2-naphthols **3r–3w** bearing methyl, ethyl, and butyl groups were well tolerated, accomplishing the asymmetric hydroxylative reaction with the outcomes of 95–99% yield and 94 : 6–95 : 5 er. The installation of substituents to the 5- and 7-positions did not influence the reaction efficiency (**3x–3z**). The MOM-protected substrate **1z** could deliver the desired product **3z** with good results without any deprotection process occurring under the reaction conditions. However, the increase of steric hindrance at the *ortho*-position of 2-naphthol was harmful as a consequence (**3aa** and **3ab**).


To show the synthetic utility of the current catalyst system, asymmetric synthesis of bioactive lacinilenes was carried out (Scheme 2). Initially, the direct deprotection of the product **3z** under acidic conditions formed the optically active lacinilene D, but an aromatization side product 1-ethyl-5-methylnaphthalene-2,6-diol was obtained.<sup>24,10</sup> It was anticipated that the TBS protecting group could be easily removed under neutral conditions, which might avoid the occurrence of the aromatization process. As expected, the TBS-substituted 2-naphthol **1ae** could be easily synthesized from **9** in 66% yield after 3 steps, which was further enantioselectively oxidized into the product **3ae** in quantitative yield and 95 : 5 er, even when it was performed at the gram scale. The absolute configuration of **3ae** from **L-PiPr<sub>2</sub>**-Sc(NTf<sub>2</sub>)<sub>3</sub> complex catalysis was determined to be (*R*) by X-ray crystal diffraction analysis.<sup>11</sup> For the benefit of the further differential biological activity study on

each enantiomer of the chiral lacinilenes,<sup>2c</sup> (*S*)-lacinilene D was synthesized using an *ent*-**L-PiPr<sub>2</sub>**-Sc(NTf<sub>2</sub>)<sub>3</sub> complex with a comparable result of 99% yield and 95 : 5 er. Next, the synthesis of optically active lacinilene C methyl ether was explored. The synthetic route began from 1,2-dihydronaphthalene **12**, which could be easily accessed from 2-methoxytoluene through a four-step protocol.<sup>2d</sup> Subsequent two-step oxidation could afford the 2-naphthol derivative **1y** in 49% yield, which underwent hydroxylative dearomatization catalyzed by 0.1 mol% of the Sc(NTf<sub>2</sub>)<sub>3</sub>/*rac*-**L-PiPr<sub>2</sub>** complex to produce racemic lacinilene **3y** in 90% yield.<sup>2d</sup> After trimethylsilylation and copper catalyzed 1,4-addition/aromatization, 2-naphthol **1af** could be attained in 45% yield after two steps. By treatment with oxaziridine **2a** in the presence of Sc(OTf)<sub>3</sub>/**L-RaPr<sub>2</sub>**, chiral lacinilene C methyl ether could be obtained in quantitative yield and 83 : 17 er, which could further transform to lacinilene C according to the literature.<sup>2b</sup>


To elucidate the stereochemical course of the oxidation process, some control experiments were conducted (Scheme 3). The optically pure oxaziridine (*S*)-**2a** reacted with 2-naphthol **1a** in the presence of the Sc(NTf<sub>2</sub>)<sub>3</sub>/**L-PiPr<sub>2</sub>** complex, affording the (*R*)-quinol **3a** in 49% yield and 95.5 : 4.5 er with the recovered oxaziridine (*S*)-**2a** in 45% yield.<sup>12d</sup> Using *ent*-**L-PiPr<sub>2</sub>** as the ligand, (*S*)-quinol **3a** was obtained in 46% yield and 90 : 10 er with the recovered oxaziridine (*S*)-**2a** in 52% yield. This indicates that the chiral matched and mis-matched effect between chiral ligand and chiral oxaziridine was not obvious in this case compared to previous reports,<sup>12</sup> and there might be negligible interaction between the chiral catalyst and oxaziridine.

To probe into the interaction between the catalysts and 2-naphthol, <sup>1</sup>H NMR analysis of the mixture of components was carried out (see ESI† for details). The chemical shift of 1-methyl-2-naphthol **1a** remained nearly unchanged after Sc(NTf<sub>2</sub>)<sub>3</sub> was added. There was an obvious high-field shift for most signals of **1a** after mixing with the Sc(NTf<sub>2</sub>)<sub>3</sub>/**L-PiPr<sub>2</sub>** catalyst. This indicates that the chiral catalyst makes the 2-naphthol reactive for hydroxylative reactions. Based on these results and our previous study on the chiral *N,N'*-dioxide-metal complex catalysts,<sup>8,13</sup> we suggested an enantioselective catalytic model as shown in Fig. 2. The ligand **L-PiPr<sub>2</sub>** binds to the scandium(III) center *via* four oxygens to form a polycyclic octahedral metal complex catalyst. The 2-naphthol coordinates to the metal center at one of the vacant sites, with its *Re*-face shielded by one amide unit of the ligand. Therefore, **2a** preferably attacked the  $\alpha$ -position of 2-naphthol from the *Si*-face to generate the corresponding *R*-configured product **3ae** and imine byproduct. If a substituent was introduced into the C3 or C4 positions of 2-naphthol, the steric hindrance discrimination between the two sides of the


#### Concise synthesis of (–)-lacinilene D and (+)-lacinilene D



#### Concise synthesis of chiral lacinilene C methyl ether



**Scheme 2** Concise synthesis of chiral lacinilene C methyl ether, (–)-lacinilene D and (+)-lacinilene D.



**Scheme 3** Control experiments.




Fig. 2 Proposed enantioselective catalytic model.

hydroxyl group decreases, thus it is difficult to control the face-selection. As a result, the enantioselectivity for the generation of product **3aa** and lacinilene C methyl ether is lower than that for the others.

## Conclusions

In summary, we have described a highly chemo- and enantioselective hydroxylative dearomatization of 2-naphthol derivatives with oxaziridine catalyzed by a chiral  $N,N'$ -dioxide- $\text{Sc}(\text{NTf}_2)_3$  complex catalyst. The desired substituted *ortho*-quinols with one quaternary carbon stereogenic center were afforded with high enantioselectivities and reactivity (up to 99% yield and 95 : 5 er). The  $\alpha$ -ketol rearrangement byproducts were efficiently suppressed. This new procedure has been successfully applied to the catalytic asymmetric synthesis of the phytoalexins lacinilenes. The application of the  $N,N'$ -dioxide/metal catalyst system in the synthesis of other bioactive molecules will be explored.

## Acknowledgements

The study was funded by the National Natural Science Foundation of China (No. 21290182, 21432006 and 21625205).


## Notes and references

- (a) D. Magdziak, S. J. Meek and T. R. R. Pettus, *Chem. Rev.*, 2004, **104**, 1383; (b) S. P. Roche and J. A. Porco Jr, *Angew. Chem., Int. Ed.*, 2011, **50**, 4068.
- (a) P. W. Jeffs and D. G. Lynn, *J. Org. Chem.*, 1975, **40**, 2958; (b) J. P. McCormick, T. Shinmyozu, J. P. Pachlatko, T. R. Schafer and J. W. Gardner, *J. Org. Chem.*, 1984, **49**, 34; (c) R. D. Stipanovic, J. P. McCormick, E. O. Schlemper, B. C. Hamper, T. Shinmyozu and W. H. Pirkle, *J. Org. Chem.*, 1986, **51**, 2500; (d) K. Krohn and G. Zimmermann, *J. Org. Chem.*, 1998, **63**, 4140.
- For selected examples of asymmetric oxidative dearomatization in the biosynthesis of natural products: (a) A. Bérubé, I. Drutu and J. L. Wood, *Org. Lett.*, 2006, **8**, 5421; (b) L. H. Mejorado and T. R. R. Pettus, *J. Am. Chem. Soc.*, 2006, **128**, 15625; (c) S. P. Cook, A. Polara and S. J. Danishefsky, *J. Am. Chem. Soc.*, 2006, **128**, 16440; (d) J. Gagnepain, F. Castet and S. Quideau, *Angew. Chem., Int. Ed.*, 2007, **46**, 1533; (e) A. Rudolph, P. H. Bos, A. Meetsma, A. J. Minnaard and B. L. Feringa, *Angew. Chem., Int. Ed.*, 2011, **50**, 5834.
- For selected reviews on dearomatization reactions of phenols and naphthols: (a) C.-X. Zhuo, C. Zheng and S.-L. You, *Acc. Chem. Res.*, 2014, **47**, 2558; (b) W.-T. Wu, L. Zhang and S.-L. You, *Chem. Soc. Rev.*, 2016, **45**, 1570; (c) W. Sun, G. Li, L. Hong and R. Wang, *Org. Biomol. Chem.*, 2016, **14**, 2164. For selected recent examples, see: (d) Q. Yin, S.-G. Wang, X.-W. Liang, D.-W. Gao, J. Zheng and S.-L. You, *Chem. Sci.*, 2015, **6**, 4179; (e) S.-G. Wang, Q. Yin, C.-X. Zhuo and S.-L. You, *Angew. Chem., Int. Ed.*, 2015, **54**, 647; (f) D. Yang, L. Wang, F. Han, D. Li, D. Zhao and R. Wang, *Angew. Chem., Int. Ed.*, 2015, **54**, 2185; (g) J. Nan, J. Liu, H. Zheng, Z. Zuo, L. Hou, H. Hu, Y. Wang and X. Luan, *Angew. Chem., Int. Ed.*, 2015, **54**, 2356; (h) D. Yang, L. Wang, M. Kai, D. Li, X. Yao and R. Wang, *Angew. Chem., Int. Ed.*, 2015, **54**, 9523; (i) S.-G. Wang, X.-J. Liu, Q.-C. Zhao, C. Zheng, S.-B. Wang and S.-L. You, *Angew. Chem., Int. Ed.*, 2015, **54**, 14929; (j) L. Yang, H. Zheng, L. Luo, J. Nan, J. Liu, Y. Wang and X. Luan, *J. Am. Chem. Soc.*, 2015, **137**, 4876; (k) J. Zheng, S.-B. Wang, C. Zheng and S.-L. You, *J. Am. Chem. Soc.*, 2015, **137**, 4880; (l) Q. Cheng, Y. Wang and S.-L. You, *Angew. Chem., Int. Ed.*, 2016, **55**, 3496; (m) H.-F. Tu, C. Zheng, R.-Q. Xu, X.-J. Liu and S.-L. You, *Angew. Chem., Int. Ed.*, 2017, **56**, 3237; (n) D. Shen, Q. Chen, P. Yan, X. Zeng and G. Zhong, *Angew. Chem., Int. Ed.*, 2017, **56**, 3242.
- For selected reviews on recent asymmetric oxidative dearomatization of phenols and naphthols: (a) M. Uyanik and K. Ishihara, *Asymmetric Oxidative Dearomatization Reaction*, in *Asymmetric Dearomatization Reactions*, ed. S.-L. You, Wiley-VCH, Weinheim, 2016, ch. 6, pp. 129–152. For examples: (b) T. Dohi, A. Maruyama, N. Takenaga, K. Senami, Y. Minamitsuji, H. Fujioka, S. B. Caemmerer and Y. Kita, *Angew. Chem., Int. Ed.*, 2008, **47**, 3787; (c) M. Uyanik, T. Yasui and K. Ishihara, *Angew. Chem., Int. Ed.*, 2010, **49**, 2175; (d) M. Uyanik, T. Yasui and K. Ishihara, *Tetrahedron*, 2010, **66**, 5841; (e) T. Oguma and T. Katsuki, *J. Am. Chem. Soc.*, 2012, **134**, 20017; (f) T. Dohi, N. Takenaga, T. Nakae, Y. Toyoda, M. Yamasaki, M. Shiro, H. Fujioka, A. Maruyama and Y. Kita, *J. Am. Chem. Soc.*, 2013, **135**, 4558; (g) M. Uyanik, T. Yasui and K. Ishihara, *Angew. Chem., Int. Ed.*, 2013, **52**, 9215; (h) T. Oguma and T. Katsuki, *Chem. Commun.*, 2014, **50**, 5053; (i) S. J. Murray and H. Ibrahim, *Chem. Commun.*, 2015, **51**, 2376; (j) D.-Y. Zhang, L. Xu, H. Wu and L.-Z. Gong, *Chem.-Eur. J.*, 2015, **21**, 10314; (k) N. Jain, S. Xu and M. A. Ciufolini, *Chem.-Eur. J.*, 2017, **23**, 4542.
- For selected examples of asymmetric hydroxylative dearomatization: (a) S. Dong, J. Zhu and J. A. Porco Jr, *J. Am. Chem. Soc.*, 2008, **130**, 2738; (b) J. K. Boppisetti and V. B. Birman, *Org. Lett.*, 2009, **11**, 1221; (c) S. Quideau, G. Lyvinec, M. Marguerit, K. Bathany, A. Ozanne-Beaudenon, T. Buffeteau, D. Cavagnat and A. Chénédé, *Angew. Chem., Int. Ed.*, 2009, **48**, 4605; (d) K. A. Volp and A. M. Harned, *Chem. Commun.*, 2013, **49**, 3001; (e) C. Bosset, R. Coffinier, P. A. Peixoto, M. El Assal,





- K. Miqueu, J.-M. Sotiropoulos, L. Pouységou and S. Quideau, *Angew. Chem., Int. Ed.*, 2014, **53**, 9860.
- 7 (a) C. Grandclaude and P. Y. Toullec, *Eur. J. Org. Chem.*, 2016, 260; (b) M. Uyanik, T. Mutsuga and K. Ishihara, *Angew. Chem., Int. Ed.*, 2017, **56**, 3956.
- 8 For reviews on chiral *N,N'*-dioxides: (a) X. H. Liu, L. L. Lin and X. M. Feng, *Acc. Chem. Res.*, 2011, **44**, 574; (b) X. H. Liu, L. L. Lin and X. M. Feng, *Org. Chem. Front.*, 2014, **1**, 298; for recent examples: (c) Q. Yao, Y. T. Liao, L. L. Lin, X. B. Lin, J. Ji, X. H. Liu and X. M. Feng, *Angew. Chem., Int. Ed.*, 2016, **55**, 1859; (d) Y. Xia, L. L. Lin, F. Z. Chang, Y. T. Liao, X. H. Liu and X. M. Feng, *Angew. Chem., Int. Ed.*, 2016, **55**, 12228; (e) Y. Zhang, Y. T. Liao, X. H. Liu, Q. Yao, Y. H. Zhou, L. L. Lin and X. M. Feng, *Chem.-Eur. J.*, 2016, **22**, 15119; (f) Y. T. Liao, X. H. Liu, Y. Zhang, Y. L. Xu, Y. Xia, L. L. Lin and X. M. Feng, *Chem. Sci.*, 2016, **7**, 3775.
- 9 (a) D. J. Michaelis, C. J. Shaffer and T. P. Yoon, *J. Am. Chem. Soc.*, 2007, **129**, 1866; (b) D. J. Michaelis, M. A. Ischay and T. P. Yoon, *J. Am. Chem. Soc.*, 2008, **130**, 6610.
- 10 We tested HCl, AlCl<sub>3</sub>/NaI, BBr<sub>3</sub>, and TMSBr to remove the MOM-protecting group, but the corresponding lacinilene D was not observed. The reactions proceeded as follows:



see ref. 2d for details.

- 11 CCDC 1536822 (**3ae**) contains the ESI crystallographic data for this paper.<sup>†</sup>
- 12 (a) P.-L. Shao, X.-Y. Chen and S. Ye, *Angew. Chem., Int. Ed.*, 2010, **49**, 8412; (b) S. X. Dong, X. H. Liu, Y. Zhu, P. He, L. L. Lin and X. M. Feng, *J. Am. Chem. Soc.*, 2013, **135**, 10026; (c) K. S. Williamson, J. W. Sawicki and T. P. Yoon, *Chem. Sci.*, 2014, **5**, 3524; (d) X. B. Lin, S. Ruan, Q. Yao, C. K. Yin, L. L. Lin, X. M. Feng and X. H. Liu, *Org. Lett.*, 2016, **18**, 3602.
- 13 X-ray single-crystal structure of *N,N'*-dioxide-Sc(III): Y. L. Liu, D. J. Shang, X. Zhou, X. H. Liu and X. M. Feng, *Chem.-Eur. J.*, 2009, **15**, 2055.

