

Cite this: *Chem. Sci.*, 2017, 8, 8094

## *Ex situ* generation of stoichiometric HCN and its application in the Pd-catalysed cyanation of aryl bromides: evidence for a transmetallation step between two oxidative addition Pd-complexes†

Steffan K. Kristensen,<sup>a</sup> Espen Z. Eikeland,<sup>b</sup> Esben Taarning,<sup>c</sup>  
Anders T. Lindhardt<sup>\*d</sup> and Troels Skrydstrup<sup>\*a</sup>

A protocol for the Pd-catalysed cyanation of aryl bromides using near stoichiometric and gaseous hydrogen cyanide is reported for the first time. A two-chamber reactor was adopted for the safe liberation of *ex situ* generated HCN in a closed environment, which proved highly efficient in the Ni-catalysed hydrocyanation as the test reaction. Subsequently, this setup was exploited for converting a range of aryl and heteroaryl bromides (28 examples) directly into the corresponding benzonitriles in high yields, without the need for cyanide salts. Cyanation was achieved employing the Pd(0) precatalyst, P(tBu)<sub>3</sub>-Pd-G3 and a weak base, potassium acetate, in a dioxane-water solvent mixture. The methodology was also suitable for the synthesis of <sup>13</sup>C-labelled benzonitriles with *ex situ* generated <sup>13</sup>C-hydrogen cyanide. Stoichiometric studies with the metal complexes were undertaken to delineate the mechanism for this catalytic transformation. Treatment of Pd(P(tBu)<sub>3</sub>)<sub>2</sub> with H<sup>13</sup>CN in THF provided two Pd-hydride complexes, (P(tBu)<sub>3</sub>)<sub>2</sub>Pd(H)(<sup>13</sup>CN), and [(P(tBu)<sub>3</sub>)Pd(H)]<sub>2</sub>Pd(<sup>13</sup>CN)<sub>4</sub>, both of which were isolated and characterised by NMR spectroscopy and X-ray crystal structure analysis. When the same reaction was performed in a THF : water mixture in the presence of KOAc, only (P(tBu)<sub>3</sub>)<sub>2</sub>Pd(H)(<sup>13</sup>CN) was formed. Subjection of this cyano hydride metal complex with the oxidative addition complex (P(tBu)<sub>3</sub>)Pd(Ph)(Br) in a 1 : 1 ratio in THF led to a transmetallation step with the formation of (P(tBu)<sub>3</sub>)Pd(H)(Br) and <sup>13</sup>C-benzonitrile from a reductive elimination step. These experiments suggest the possibility of a catalytic cycle involving initially the formation of two Pd(II)-species from the oxidative addition of L<sub>n</sub>Pd(0) into HCN and an aryl bromide followed by a transmetallation step to L<sub>n</sub>Pd(Ar)(CN) and L<sub>n</sub>Pd(H)(Br), which both reductively eliminate, the latter in the presence of KOAc, to generate the benzonitrile and L<sub>n</sub>Pd(0).

Received 6th September 2017  
Accepted 6th October 2017

DOI: 10.1039/c7sc03912c

rsc.li/chemical-science

## Introduction

Benzonitriles are a valuable class of compounds with a broad range of applications serving as important substructures in many agrochemicals, pharmaceuticals, dyes, polymers and other organic materials.<sup>1</sup> The nitrile group is also a precursor to a variety of important functionalities such as aldehydes,

ketones, carboxamides and carboxylates.<sup>2</sup> Conventional means for the instalment of a cyanide group onto an aromatic ring include the Rosenmund-von Braun or Sandmeyer reactions applying stoichiometric CuCN with aryl halides or diazonium salts.<sup>3</sup> Yet, in the last decade, transition metal catalysed cyanations of aromatic halides with metal complexes based on copper,<sup>4</sup> nickel<sup>5</sup> and palladium<sup>6,7</sup> have held a key position. In particular, the protocols based on Pd-catalysis through the extensive work by Beller,<sup>7</sup> Grushin<sup>8</sup> and Buchwald<sup>9</sup> are characterised by relatively mild reaction conditions and good functional group tolerance. A variety of different cyanide sources have been applied including salts or derivatives such as NaCN,<sup>8,10</sup> KCN,<sup>6a,c,d,11</sup> (Me)<sub>3</sub>SiCN,<sup>12</sup> acetone cyanohydrin,<sup>13</sup> K<sub>4</sub>Fe(CN)<sub>6</sub>,<sup>9a,14</sup> Zn(CN)<sub>2</sub>,<sup>9b,15</sup> CuCN,<sup>16</sup> etc. (Scheme 1). However, many of these sources also have their disadvantages. For example, metal contaminants can become an issue with the use of potassium ferrocyanide, whereas with the cyanohydrin of acetone, this chemical must be stored carefully and is approx. 20 times more costly than potassium cyanide.† Furthermore,

<sup>a</sup>Carbon Dioxide Activation Center (CADIAC), The Interdisciplinary Center (iNANO), Department of Chemistry, Aarhus University, Gustav Wieds Vej 14, 8000 Aarhus, Denmark. E-mail: ts@chem.au.dk

<sup>b</sup>Center for Materials Crystallography, The Interdisciplinary Center (iNANO), Department of Chemistry, Aarhus University, Langelandsgade 140, 8000 Aarhus, Denmark


<sup>c</sup>Haldor Topsøe A/S, New Business R&D, Nymøllevvej 55, 2800 Kgs. Lyngby, Denmark

<sup>d</sup>Carbon Dioxide Activation Center (CADIAC), The Interdisciplinary Center (iNANO), Biological and Chemical Engineering, Department of Engineering, Aarhus University, Finlandsgade 22, 8200 Aarhus N, Denmark

† Electronic supplementary information (ESI) available. CCDC 1504904–1504908. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7sc03912c




Established methods: Pd-catalysed cyanation of aryl halides with different cyanide sources



Cyanide source: NaCN, KCN, (Me)<sub>3</sub>SiCN, acetone cyanohydrin, CuCN, K<sub>4</sub>Fe(CN)<sub>6</sub>, Zn(CN)<sub>2</sub>

This work: Pd-catalysed cyanation of aryl bromides with gaseous HCN



Scheme 1 Previous developments in Pd-catalysed cyanation protocols and our new approach applying gaseous HCN.

considering the fact that all of these cyanide precursors originate from hydrogen cyanide (HCN), which is produced on a million-ton scale size *via* the Andrussov<sup>17</sup> and BMA-Degussa processes<sup>18</sup> from natural gas and ammonia or as a byproduct in acrylonitrile production,<sup>18,19</sup> there could be an interest in the development of a cyanation process of aryl halides exploiting this original source of cyanide, namely gaseous HCN.<sup>20</sup>

Only three reports are found in the literature applying HCN directly or by its *in situ* formation. In 2015, Buchwald and Hooker reported a <sup>11</sup>C-cyanation study for the synthesis of aromatic cyanides with applications for positron emission tomography (PET).<sup>21</sup> Here, submicromolar amounts of gaseous H<sup>11</sup>CN dissolved in THF were introduced to a 1000-fold excess of preformed oxidative addition palladium(II) complex, affording the desired <sup>11</sup>C-benzonitriles nearly instantaneously. A follow-up paper by the same two groups demonstrated the possibilities of this approach for the <sup>11</sup>C-labelling of unprotected peptides.<sup>22</sup> Alternatively, Beller and co-workers reported the *in situ* release of HCN from acetone cyanohydrin although this was carried out in the presence of sodium carbonate, which undoubtedly leads to the direct formation of the reactive cyanide anion species.<sup>13</sup>

Nevertheless, there is a reluctance to apply HCN (b.p. 27 °C) as a cyanation reagent in an academic setting or an R&D laboratory, which is undoubtedly linked to the high toxicity and explosive nature of this gaseous one-carbon reagent, thereby complicating its handling, storage and transportation.<sup>23</sup> Another challenge in the use of hydrogen cyanide as a cyanation reagent is how to control its dosage in stoichiometric amounts as Pd-catalysed cyanations of aryl halides are particularly sensitive to the cyanide concentrations. The strong binding of cyanide anion to transition metals can lead to catalyst deactivation if present in elevated concentrations.<sup>7,24</sup> With other cyanide reagents, this complication is normally addressed by their slow addition to the reaction mixture, slow mass transport

of the cyanide anion by specific water/organic solvents mixtures or by use of cyanide complexes with low solubilities.<sup>8,9b,13,14</sup> Grushin and co-workers have published impressive and detailed studies on the mechanism of cyanide induced catalyst deactivation in the Pd-catalysed cyanation of aromatic halides. In their work, they report that addition of an excess of <sup>13</sup>C-labelled potassium cyanide (K<sup>13</sup>CN) to different intermediates of the catalytic cycle in the Pd-catalysed cyanation of iodobenzene led to coordinatively saturated and catalytically inactive palladium complexes such as [Pd(CN)<sub>4</sub>]<sup>2-</sup>, [HPd(CN)<sub>3</sub>]<sup>2-</sup> or [ArPd(CN)<sub>3</sub>]<sup>2-</sup>.<sup>25</sup> More interestingly, the same group revealed that traces of water in the reaction mixture combined with K<sup>13</sup>CN forms H<sup>13</sup>CN that immediately undergoes oxidative addition with Pd(0) leading to the same coordinatively saturated and off-cycle palladium complexes with concurrent formation of hydrogen gas.<sup>26</sup>


In this paper, we report on the direct use of industrially important hydrogen cyanide in the palladium-catalysed cyanation of (hetero)aromatic bromides (Scheme 1). These results deviate from conventional wisdom, since Grushin's earlier work demonstrated the propensity of palladium(0) to undergo fast oxidative addition into the H-CN bond shutting down the catalytic cycle. And yet, our results demonstrate that conditions can be identified whereby the presence of HCN does not terminate the cyanation reaction, but enhances the reactivity *vis-à-vis* cyanide salts. Secondly and most important, we describe a simple and safe setup whereby HCN is delivered in stoichiometric amounts by *ex situ* generation in a two-chamber reactor, thereby providing a simple and safe setting for handling gaseous HCN in small-scale reactions. We demonstrate the usefulness of this setup not only for the Pd-catalysed cyanation of aryl bromides, but also for the Ni-mediated hydrocyanation of styrenes as a test reaction. With respect to the cyanation reactions good functional group tolerance was obtained, and the method proved amenable to scale-up, but also to carbon-13 isotope labelling applying H<sup>13</sup>CN. Surprisingly, the developed conditions proved to be dependent on water as the co-solvent and the presence of a weak base, KOAc. In their absence, catalytic shutdown was observed, thereby indicating the operation of an alternative mechanism to Pd-catalysed cyanations applying cyanide salts. Studies and isolation of the oxidative addition complexes suggest a mechanism involving the transmetalation between two palladium(II) species, both formed by the oxidative addition of palladium(0) into either HCN or the aryl bromide electrophile.

## Results and discussion

### Hydrogen cyanide releasing studies

In order to provide a system for the controlled dosage of hydrogen cyanide in stoichiometric quantities, we envisioned that the two-chamber system previously utilised for the *ex situ* liberation of carbon monoxide,<sup>27</sup> hydrogen<sup>28</sup> and ethylene<sup>29</sup> from specific nongaseous precursors could be exploited. Such an approach would provide a simple and safe setup without the direct handling of HCN gas. Hence, our focus was first directed to identifying a suitable system for the generation of HCN gas





**Scheme 2** Ni-catalysed hydrocyanation of styrenes employing a two-chamber system with *ex situ* generated HCN. <sup>a</sup>Chamber A: styrene (1.0 mmol), Ni(COD)<sub>2</sub> (5.0 mol%), XantPhos (7.5 mol%) and CPME (2 mL). Chamber B: KCN (1.5 mmol), ethylene glycol (1 mL), AcOH (9.0 mmol). Isolated yields are given as an average of 2 runs.

under a closed environment, and the well-established nickel-catalysed hydrocyanation of olefins was exploited as a test system for the optimisation.<sup>30–32</sup> After considerable experimentation on the hydrocyanation of styrene (see ESI<sup>†</sup>), we finally adopted the following conditions as illustrated in Scheme 2. A combination of KCN, ethylene glycol and AcOH provided the desired release of gaseous HCN. Ethylene glycol serves a dual purpose acting both as solvent while simultaneously ensuring separation of the AcOH from KCN until stirring is initiated. By utilising this setup for the HCN producing chamber in combination with the Ni-catalysed hydrocyanation conditions in the second chamber (Ni(COD)<sub>2</sub>, XantPhos), a range of different styrenes bearing both electron donating and withdrawing groups could successfully be hydrocyanated to products 1–7 in near quantitative yields. Full conversion of all styrene derivatives was attained applying only 1.5 equivalents of KCN for the production of HCN, indicating the effectiveness of this gas generator.

### Studies on the cyanation of (hetero)aryl bromides

With a simple and convenient system for the release of gaseous HCN in hand, we next turned our attention towards the Pd-catalysed cyanation of (hetero)aryl bromides. As depicted in the scheme of Table 1, initial optimisation results were carried out with 4-bromoanisole. This study revealed that reaction conditions consisting of a catalyst formed from Pd(dba)<sub>2</sub> with P(*t*Bu)<sub>3</sub>, in the presence of hydrogen cyanide (1.5 equiv.), a weak base such as KOAc, and a solvent mixture of water and dioxane could generate 4-methoxybenzonitrile (**8**) with a conversion of 78% according to the <sup>1</sup>H NMR analysis of the crude reaction mixture (Table 1, entry 1). P(*t*Bu)<sub>3</sub> as a ligand proved to be

**Table 1** Optimisation of the Pd-catalysed cyanation of aryl bromides using gaseous HCN<sup>a</sup>

Entry	Deviation from lead conditions	Conversion <sup>b</sup> (%)
1	None	78
2	Ligand: PCy <sub>3</sub>	0
3	Ligand: XPhos	60
4	Ligand: <i>t</i> Bu-XPhos	74
5	Ligand: DPPF	0
6	Ligand: XantPhos (2.5 mol%)	0
7	Base: Cy <sub>2</sub> NMe	38
8	Base: DBU	5
9	Base: Et <sub>3</sub> N	67
10	Base: KOAc (3.0 equiv.)	82
11	<b>Pd/ligand: P(<i>t</i>Bu)<sub>3</sub>-Pd-G3 KOAc (3.0 equiv.)</b>	<b>90 (90)<sup>c</sup></b>
12	Pd/ligand: P( <i>t</i> Bu) <sub>3</sub> -Pd-G3 solvent: THF	77
13	Pd/ligand: P( <i>t</i> Bu) <sub>3</sub> -Pd-G3 solvent: Toluene	61

<sup>a</sup> Chamber A: 4-bromoanisole (1.0 mmol), Pd(dba)<sub>2</sub> (2.5 mol%), P(*t*Bu)<sub>3</sub> (5.0 mol%), KOAc (2.0 mmol) in dioxane (1 mL) and H<sub>2</sub>O (1 mL). Chamber B: KCN (1.5 mmol), ethylene glycol (1 mL) and AcOH (9.0 mmol). <sup>b</sup> Determined by <sup>1</sup>H NMR using mesitylene as an internal standard. P(*t*Bu)<sub>3</sub>-Pd-G3 = third generation Buchwald precatalyst with the tri-*tert*-butylphosphine ligand. <sup>c</sup> Isolated yield.

superior to other monodentate and bidentate ligands (entries 2–6; see ESI<sup>†</sup> section for full optimisation studies), which is in accord with earlier results reported by Grushin and co-workers.<sup>8,26</sup> In general, weak bases such as KOAc and NaOAc provided better conversions to the desired 4-methoxybenzonitrile, whereas stronger bases including Cy<sub>2</sub>NMe, DBU and Et<sub>3</sub>N all proved inferior (entries 7–9). Increasing to three equivalents of KOAc provided a slight increase in the conversion to 82% (entry 10). The efficiency of the reaction could also be increased further exploiting the Buchwald pre-catalyst, P(*t*Bu)<sub>3</sub>-Pd-G3, providing a 90% isolated yield of benzonitrile **8** after column chromatography (entry 11). Control experiments with omission of either base or palladium precursors resulted in low or no conversion, resp.

Different water : solvent combinations were tested and dioxane proved to be the solvent of choice (entries 12 and 13, see also ESI<sup>†</sup>). Notably, the presence of water as the co-solvent is of key importance as similar experiments applying only dioxane as the solvent resulted in complete catalytic shutdown. In our scope studies, we later discovered that a 2 : 1 mixture of water and dioxane provided higher yields for certain substrates, and hence this ratio was used for all subsequent cyanations. Not surprisingly and in accord with Grushin's previous results, carrying out the Pd-catalysed cyanation in a single chamber reactor applying KCN directly with or without the addition of acetic acid under the conditions developed led to no formation of **8** (result not shown).




With the optimised reaction conditions in hand, we set out to explore the scope of the Pd-catalysed cyanation using gaseous HCN. All yields are reported as an average of two runs and the results are depicted in Scheme 3. Aryl bromides carrying electron donating substituents were initially examined. Methoxy-, hydroxy-, alkyl- and aryl-substituted aryl bromides provided the desired compounds in high yields ranging from 90% to 97% (compounds 8–11). Electron withdrawing substituents such as cyano, acyl and carboxylate afforded compounds 12, 13 and 14 in yields attaining quantitative. Even *p*-bromobenzoic acid underwent successful coupling affording *p*-cyanobenzoic acid (15) in an 86% yield, using 4 equiv. of KOAc combined with 5 mol% of Buchwald's precatalyst. The use of phenol and benzoic acid derivatives in the Pd-catalysed cyanation with NaCN were previously shown by Ushkov and Grushin to lead to catalyst deactivation due to the formation of HCN.<sup>8</sup> *p*-Bromobenzaldehyde also proved reactive with the isolation of 16 in a somewhat lower yield. This slight reduction in isolated yield

can possibly be explained by competing benzoin condensation of both starting material and product. The substitution pattern of bromoanilines interfered significantly with the developed conditions and installment of the cyano-functionality onto 2-, 3-, and 4-bromoaniline was achieved in yields of 95%, 77% and 47% respectively (compounds 17–19). Aryl bromides displaying *ortho*-substitutions were also effective for these substitution reactions as illustrated with compounds 17 and 20–22, all isolated in yields between 92% and 96%. However, 2-bromo-1,3-dimethylbenzene did require an increase of the catalyst loading to achieve the high 96% isolated yield of 21. Lowering the reaction temperature of the HCN consuming chamber, from 60 °C to 45 °C, ensured a chemoselective activation of the aromatic bromide in the presence of the *p*-chloride affording 23 in an 88% isolated yield. Subsequently, an aryl bromide possessing a benzylic alcohol, and five heteroaromatic bromides were tested under the developed conditions. All underwent successful cyanation to afford the desired target molecules in yields ranging from 75% to 95% (compounds 22, 24–28). It should be mentioned though that despite the successful coupling of these handful of heteroaryl bromides, two other heterocycles tested such as 2-acetyl-5-bromothiophene and 3-bromobenzothiophene failed to provide the corresponding nitrile product. The reasons for this catalytic shutdown are not completely understood (see Discussion in next section). Finally, attempts to perform catalytic cyanation on 4-biphenyl triflate, applying the developed protocol, were non-rewarding with no conversion observed for this electrophile.


Next, attention was directed to implementing this developed protocol for <sup>13</sup>C-labelling. By simply substituting KCN with its <sup>13</sup>C-labelled counterpart under otherwise unchanged conditions, direct access to isotopically labeled benzonitrile derivatives were achieved. Four entries from Scheme 3 were selected for labelling, and all compounds were isolated in near identical yields to the unlabeled compounds (compounds 10-<sup>13</sup>C, 11-<sup>13</sup>C, 12-<sup>13</sup>C and 28-<sup>13</sup>C).

Finally, the developed protocol was tested for the synthesis of three active pharmaceutical ingredients, namely dapivirine (reverse transcriptase inhibitor), citalopram (serotonin reuptake inhibitor) and letrozole (non-steroidal aromatase inhibitor), the results of which are depicted in Scheme 4.<sup>33</sup> Dapivirine (29) and citalopram (30) were obtained in satisfactory 96% and 86% isolated yields, respectively, from the corresponding bromide precursors applying 5.0 mol% of the Buchwald precatalyst. For the synthesis of letrozole (31) a double cyanation is required with the dibromide 31a. Doubling the loading of KCN for HCN release to 3 equivalents combined with an increase in KOAc to 4 equivalents, gratifyingly afforded the dicyanide 31 in an excellent 97% isolated yield again with 5.0 mol% precatalyst. The synthesis of letrozole was then attempted on a fivefold scale combined with a 100 mL two-chamber reactor to afford a near identical isolated yield of 95%. Finally, scaling up to 10 mmol in a 200 mL two-chamber reactor, with an HCN release from 30 mmol of KCN, corresponding to more than 700 mL of gaseous HCN, afforded the desired pharmaceutical in a good 85% isolated yield.



**Scheme 3** Pd-catalysed cyanation of aryl bromides. <sup>a</sup>Chamber A: aryl bromide (1.0 mmol), P(*t*Bu)<sub>3</sub>-Pd-G3 (2.5 mol%) and KOAc (3.0 mmol) in dioxane (1 mL) and H<sub>2</sub>O (2 mL). Chamber B: KCN (1.5 mmol) or K<sup>13</sup>CN (1.5 mmol) ethylene glycol (1 mL) and AcOH (9 mmol). Isolated yields are given as an average of 2 runs. <sup>b</sup>5.0 mol% catalyst used. <sup>c</sup>4.0 equiv. of KOAc used. <sup>d</sup>HCN consuming chamber only heated to 45 °C.





Scheme 4 Synthesis of pharmaceuticals by Pd-catalysed cyanation and scale up studies. <sup>a</sup>For reactions on 1.0 mmol scale, yields are an average of two runs.

### Mechanistic investigations


Having established the scope of the Pd-catalysed cyanation of (hetero)aryl bromides with hydrogen cyanide, some questions still remained. Given a pK<sub>a</sub> difference in water for acetic acid (4.76) and HCN (9.21) of almost 4.5 units, indicates that HCN is potentially the reactive species in solution and not the free cyanide anion. However, previous reports by Grushin and co-workers clearly demonstrated that the presence of free HCN formed from the protonation of KCN with trace amounts of water, quickly leads to shutdown of the catalytic activity.<sup>8,26</sup> Still this deactivation is a combined result of Pd(0) undergoing oxidative addition into the H–CN bond followed by trapping of this species with excess cyanide anion to generate off-cycle palladium(II) cyanide complexes. Furthermore, there was not a clear indication why certain heteroaryl bromides were successful coupling partners with HCN, whereas others were not. With the exception of the protocol developed by Buchwald and Hooker, in which H<sup>11</sup>CN is used to form PET tracers, no reports are found on palladium-catalysed cyanation of aryl halides using gaseous HCN as the reactant.<sup>21</sup> Although it should be noted under the conditions used for <sup>11</sup>C-isotope-labelling, the Pd(II)-aryl complex is in an approx. 1000-fold excess compared to the H<sup>11</sup>CN generated. With this in mind it was decided to take a closer look at the possible mechanism for the developed protocol.

The overall accepted mechanism for the Pd-catalysed cyanation using sodium or potassium cyanide is believed to go through an initial oxidative addition of the Pd(0) complex into the aryl halide, nucleophilic displacement by cyanide on the

formed palladium(II) centre, and lastly a reductive elimination furnishing the desired benzonitrile (Scheme 5).<sup>64,7,8,26</sup> When metal complexes such as Zn(CN)<sub>2</sub> are utilised, a change in mechanism occurs, whereby nucleophilic substitution is replaced with a transmetalation step.<sup>9b,34</sup>

During the optimisation of the catalytic system, different bases were tested (Table 1, entries 7–10). A clear trend can be abstracted from this base screening, where an increase in base strength leads to a decrease in conversion. This can be explained by the higher degree of deprotonation of HCN with strong bases leading to catalytic shutdown. The effect is highest for the strong base DBU affording a mere 5% conversion to product **8**. The optimum conditions found in this work applies KOAc as base, suggesting that HCN is the reactive species and not cyanide itself.

Initially, the oxidative addition of Pd(P(*t*Bu)<sub>3</sub>)<sub>2</sub> to H<sup>13</sup>CN in THF as the solvent was investigated using the two-chamber reactor (see ESI† for reaction details).§ This afforded two palladium-hydride species in roughly a 6 : 1 ratio, as observed by <sup>1</sup>H NMR analysis of the reaction mixture, with the hydride signals residing at –11.42 and 16.79 ppm (Fig. 1 and 3).<sup>35</sup> Careful manipulation of the reaction mixture allowed for the isolation and crystallisation of both hydride species **32** and **33**, the structures of which were confirmed by X-ray crystal structural analysis (Fig. 2 and 4).



Scheme 5 Accepted mechanism for the cyanation of aryl halides with cyanide salts.




Fig. 1 <sup>1</sup>H NMR ( $J_{\text{H-C}} = 53.0$  Hz,  $J_{\text{H-P}} = 4.7$  Hz) and <sup>31</sup>P NMR ( $J_{\text{P-C}} = 11.7$  Hz) of compound **32** in THF-*d*<sub>8</sub>.






Fig. 2 ORTEP representation of complex 32.



Fig. 3  $^1\text{H}$  NMR and  $^{31}\text{P}$  NMR of compound 33 in  $\text{CDCl}_3$ .

Fig. 4 ORTEP representation of complex 33.

Compound 32 was identified as the oxidative addition complex with  $\text{H}^{13}\text{CN}$ ,  $(\text{P}(\text{tBu})_3)_2\text{Pd}(\text{H})(^{13}\text{CN})$ . While others have reported the corresponding  $\text{HBr}$  and  $\text{HCl}$  adducts,<sup>36,37</sup> it is to the best of our knowledge the first time compound 32 has been isolated. The hydride signal is located at  $-11.42$  ppm as a double triplet in the  $^1\text{H}$  NMR spectrum (Fig. 1), whereby the multiplicity originates from a large *trans*-coupling with the 13-carbon of the cyanide, and small *cis*-coupling with the two

equivalent phosphine ligands.<sup>¶</sup> The  $^{31}\text{P}$  NMR analysis of 32 reveals a doublet at 93.1 ppm arising from  $\text{P}-\text{C}$  coupling with the  $^{13}\text{C}$ -labelled cyanide. Due to disturbance in the crystal structure, the hydride could not be located. However, in view of the combined data obtained from the NMR and the X-ray crystal structural analysis, we are confident that the structure of 32 is as indicated in Fig. 2.

To our surprise, the X-ray crystal structure analysis of the palladium-hydride complex 33 revealed it to be a tri-metallic species as illustrated in the ORTEP representation of Fig. 4. In the  $^1\text{H}$  NMR spectrum a hydride signal is located at  $-16.79$  ppm as a multiplet (Fig. 3), whereas for the  $^{31}\text{P}$  NMR spectrum, a singlet at 84.7 ppm can be found. A closer look at this  $\text{Pd}_3$ -complex, shows its resemblance to  $\text{K}_2[\text{Pd}(^{13}\text{CN})_4]$ , which can also be produced from the reaction of  $\text{Pd}(0)$  with  $\text{H}^{13}\text{CN}$  in turn formed from the hydrolysis of  $\text{K}^{13}\text{CN}$  with water.<sup>26</sup> In structure 33, the anionic core,  $[\text{Pd}(^{13}\text{CN})_4]^{2-}$ , carries two cationic counter ions in the form of  $[(\text{P}(\text{tBu})_3)\text{Pd}(\text{H})]^+$ .

To investigate the importance of the reaction conditions and influence on the formation of 32 and 33, a series of experiments were performed the results of which are shown in Table 2. As can be seen from entries 1–3, neither the amount of  $\text{H}^{13}\text{CN}$  nor the presence of water appears to influence the distribution between 32 and 33. However, the addition of  $\text{KOAc}$  and water resulted in the exclusive formation of the  $\text{Pd}$ -hydride complex 32 (entry 4).<sup>||</sup> Increasing the  $\text{H}^{13}\text{CN}$  loading from 1.0 to 3.0 equiv. provided a 92% conversion to 32. Full conversion to 32 was obtained with 3 equiv. of  $\text{H}^{13}\text{CN}$  in combination with 6 equiv. of  $\text{KOAc}$  (entry 6). Finally, direct formation of 33 from 32 can also be achieved by heating 32 in THF at  $60^\circ\text{C}$ , and after 3 h, a 3 : 1 ratio between 32 and 33 is achieved. This result could be explained by the slow transmetalation between two  $\text{Pd}$ -hydride species 32 leading to the formation of  $(\text{P}(\text{tBu})_3)_2\text{Pd}(\text{H})_2$  and

Table 2 Mechanistic evaluation of the reaction between  $\text{H}^{13}\text{CN}$  and  $\text{Pd}(\text{P}(\text{tBu})_3)_2^a$ 

Entry	Solvent	$\text{H}^{13}\text{CN}$ (equiv.)	$\text{KOAc}$ (equiv.)	Ratio 32 : 33 <sup>b</sup>
1	THF	1.0	—	85 : 15 (100%)
2	THF	3.0	—	84 : 16 (100%)
3	THF : $\text{H}_2\text{O}$	1.0	—	86 : 14 (100%)
4	THF : $\text{H}_2\text{O}$	1.0	2.0	>95 : 5 (29%)
5	THF : $\text{H}_2\text{O}$	3.0	2.0	>95 : 5 (92%)
6	THF : $\text{H}_2\text{O}$	3.0	6.0	>95 : 5 (97%)

<sup>a</sup> Reactions performed on a 0.1 mmol scale. Reactions stopped after 3 h.  
<sup>b</sup> Values in brackets are given as conversions.




$(P(tBu)_3)_2Pd(^{13}CN)_2$ .<sup>35,38</sup> Whereas, the former can reductively eliminate generating Pd(0) and dihydrogen, we speculate that  $(P(tBu)_3)_2Pd(^{13}CN)_2$  could abstract cyanide from two Pd-hydride complexes **32** ultimately leading to the formation of the Pd<sub>3</sub>-complex **33**.


To further probe the necessity of the bulky and electron rich  $P(tBu)_3$  ligand,  $Pd(PPh_3)_4$  was dissolved in THF and treated with  $H^{13}CN$  (Scheme 6). This resulted in the sole formation of  $(PPh_3)_2Pd(^{13}CN)_2$  (**34**), the structure of which was confirmed by both NMR and X-ray analysis (see ESI†).<sup>\*\*</sup> Possibly, the transmetallation event involving two  $(PPh_3)_2Pd(H)(^{13}CN)$  species is significantly faster than that for complex **32**, and therefore cyanide abstraction from  $(PPh_3)_2Pd(H)(^{13}CN)$  with complex **34** cannot compete as with similar palladium species bearing the  $P(tBu)_3$  ligand.

Next, we addressed the question whether complex **32** represents an active participant in the catalytic cycle. To this end, we examined the reaction between 4-bromobiphenyl and **32** in THF-*d*<sub>8</sub> at room temperature with mesitylene as an internal standard (Scheme 7). The reaction progress was followed by both <sup>1</sup>H- and <sup>31</sup>P-NMR. From the <sup>1</sup>H NMR spectrum it was clear that as the reaction progressed a build-up of the palladium-hydride species **35** in addition to the formation of the aromatic nitrile **11-<sup>13</sup>C** was observed. The reaction turned out to be relatively slow at 25 °C despite a 1 : 1 relationship between complex **32** and 4-bromobiphenyl. Nevertheless, after 12 h, the conversion into **11-<sup>13</sup>C** had reached 55%.

To verify that the structure of the new Pd-hydride species formed indeed is as proposed for compound **35**, a sample of this complex was prepared according to a known literature procedure involving the treatment of  $Pd(P(tBu)_3)_2$  with pyridinium bromide in toluene.<sup>39</sup> As can be seen in Fig. 5, complex **35** produced a triplet at -15.63 ppm in the <sup>1</sup>H NMR spectrum arising from the coupling to the two equivalent phosphorus atoms. In the <sup>31</sup>P NMR spectrum, **35** gives rise to a doublet located at 83.5 ppm. These spectroscopic data were in accord with those observed from the reaction of **32** with 4-bromobiphenyl. Finally, by dissolving **35** in a small amount of  $CH_2Cl_2$  layered with heptane resulted in crystals that were suitable for X-ray analysis (Fig. 6).



Scheme 6 Reaction between  $Pd(PPh_3)_4$  and  $H^{13}CN$ . <sup>a</sup>Reaction performed on a 0.1 mmol scale with a two-chamber system. The reaction was stopped after 3 h.



Scheme 7 Reactivity studies between complex **32** with 4-bromobiphenyl. <sup>a</sup>0.02 mmol of both **32** and 4-bromobiphenyl were added to a NMR-tube. Mesitylene was used as an internal standard.




Fig. 5 <sup>1</sup>H NMR ( $J_{H-P} = 7.2$  Hz) and <sup>31</sup>P NMR ( $J = 1.9$  Hz) of compound **35** in THF-*d*<sub>8</sub>.




Fig. 6 ORTEP representation of complex **35**.

The formation of aryl nitrile **11-<sup>13</sup>C** is only possible if complex **32** can reductively eliminate generating  $Pd(P(tBu)_3)_2$  and  $H^{13}CN$ , thereby implying that the addition of  $H^{13}CN$  to Pd(0) is a reversible process.<sup>40</sup> The Pd(0) species produced from this reductive elimination step can then either undergo an oxidative addition to  $H^{13}CN$  again or alternatively to the aryl bromide generating  $(P(tBu)_3)Pd(Ar)(Br)$ .<sup>41</sup> With the presence in solution of both this complex and **32**, a transmetallation event could take place between these two species leading to the formation of the Pd-hydride **35** and  $(P(tBu)_3)Pd(Ar)(^{13}CN)$ , which subsequently undergoes reductive elimination to the desired benzonitrile **11-<sup>13</sup>C** and Pd(0).<sup>††</sup>

The progress of the reaction between **32** and *p*-bromobiphenyl was monitored by <sup>31</sup>P NMR spectroscopy, the results of which are shown in Fig. 7. Within the first 10 minutes  $Pd(P(tBu)_3)_2$  (85.5 ppm) and free  $P(tBu)_3$  (63.4 ppm) were formed, which is consistent with the observations by Fu and coworkers when a  $P(tBu)_3/Pd(0)$  ratio of 2 : 1 is used.<sup>42</sup> The depletion of complex **32** then occurred with concurrent build-up of the Pd-hydride **35** residing at 83.5 ppm. No signal from the oxidative addition complex  $(P(tBu)_3)Pd(Ar)(Br)$  was observed, suggesting that the oxidative addition of  $Pd(P(tBu)_3)_2$  to 4-bromobiphenyl is the rate-determining step in this setup.







Fig. 7  $^{31}\text{P}$  NMR spectra of the reaction between complex 32 and *p*-bromobiphenyl.

From these initial experiments, a possible mechanistic scenario for the Pd-catalysed cyanation of aryl bromides with gaseous HCN is depicted in Scheme 8, which accounts for the above observations. After formation of the catalytically active palladium(0) species, two reversible oxidative addition events take place forming  $L_n\text{Pd}(\text{H})(\text{CN})$  and  $L_n\text{Pd}(\text{Ar})(\text{Br})$ . The necessity for the oxidative addition steps to be reversible at least for the latter complex may explain why aryl triflates are not competent electrophiles for this catalytic protocol. The two complexes can then undergo transmetalation providing  $L_n\text{Pd}(\text{Ar})(\text{CN})$  and  $L_n\text{Pd}(\text{H})(\text{Br})$ . Similar transmetalative events have previously been reported for two Pd(II)-aryl species.<sup>43</sup> Finally, the benzonitrile and the active  $L_n\text{Pd}(0)$  species are reformed through two reductive elimination events involving both  $L_n\text{Pd}(\text{Ar})(\text{CN})$  and  $L_n\text{Pd}(\text{H})(\text{Br})$ . In the latter case, base is required to initiate the regeneration of  $L_n\text{Pd}(0)$ , and the presence of aqueous KOAc should be sufficient for promoting this reduction step at the metal centre. Whereas stronger bases would be able to carry out this step as well, they would also lead to the deprotonation of HCN resulting in increased concentrations of cyanide anion in solution. Subsequently, this cyanide would trap the active species in the catalytic cycle as multi-cyano palladium complexes and as such lead to catalytic shutdown.



Scheme 8 A proposed mechanism for the Pd-catalysed cyanation of aryl bromides with HCN.


The mechanism depicted in Scheme 8 also supports the findings made by Buchwald and Hooker.<sup>21,22</sup> In their setup,  $\text{H}^{13}\text{C}\text{N}$  is introduced to a large excess of the oxidative addition complex of the same type as 36. Only a small portion of this Ar-Pd(II) complex would have to undergo reductive elimination to afford a Pd(0) species that could trap the added  $\text{H}^{13}\text{C}\text{N}$  leading to product formation.

In order to investigate the viability of such a transmetalation step between two Pd(II) species of the proposed catalytic cycle in Scheme 8, we investigated the reaction between Pd-hydride 32 and the preformed oxidative addition complex 36 (Scheme 9a). The aryl-Pd complex 36 was prepared in a 79% isolated yield *via* the oxidative addition of  $\text{Pd}(\text{P}(\text{tBu})_3)_2$  to bromobenzene according to a literature procedure.<sup>44</sup> Complexes 32 and 36 were dissolved in  $\text{THF-d}_8$ , and using mesitylene as the internal standard, their transformation was followed by  $^{31}\text{P}$  NMR spectroscopy. From this experiment, three species were formed as the reaction progressed, being  $^{13}\text{C}$ -benzonitrile (37),  $(\text{P}(\text{tBu})_3)_2\text{-Pd}(\text{H})(\text{Br})$  (35) and the  $\text{Pd}_3$ -complex 33. The transformation was completed in less than 30 min, and applying the internal standard, 37 was formed in a 53% yield while most of the remaining  $^{13}\text{C}$ -labelled cyanide could be accounted for from the formation of 33 (11% yield).

Given the undoubtedly strong binding mode of cyanide to palladium(II), the possibility of free cyanide in solution is most likely absent, thereby eliminating the possibility for the formation of 35 through a nucleophilic displacement pathway. This in turn indicates that 35 forms as a result of a transmetalation step between 36 and one or both of complexes 32 and 33.

As mentioned above, the  $\text{Pd}_3$ -complex 33 is presumably not formed under the optimised conditions. Nevertheless, we investigated whether 33 could also promote benzonitrile formation alone. The trinuclear complex 33 was mixed with 36 in a 1 : 1 relationship in  $\text{CDCl}_3$ , and followed by  $^1\text{H}$  NMR spectroscopy. Surprisingly, complete formation of  $^{13}\text{C}$ -benzonitrile (37) and Pd-hydride 35 was achieved after only 2 h (Scheme 9b). To evaluate the importance of the  $[(\text{P}(\text{tBu})_3)_2\text{-Pd}(\text{H})]^+$  cation in 33, the direct reaction between  $\text{K}_2[\text{Pd}(\text{CN})_4]$  and 36 was tested (Scheme 9c). However, following the reaction







**Scheme 9** Transmetalation studies with complex **36**. <sup>a</sup>0.02 mmol of either **32**, **33** or  $K_2[Pd(CN)_4]$  with **36** were added to a NMR-tube. Mesitylene was used as an internal standard.

by <sup>1</sup>H NMR spectroscopy, no conversion was observed between these two palladium complexes as seen from the absence of signals for benzonitrile and Pd-hydride **35**.


Next, we examined whether the different Pd-cyanide species are catalytically competent species and can promote the catalytic conversion of aryl bromides to aryl nitriles with HCN. By using **32** as the palladium source we could isolate biphenyl nitrile **11** in quantitative yield (Scheme 10). On the other hand, with **33** as the palladium source no conversion to product was observed. This is in perfect accordance with the observations shown in Table 2, demonstrating that the presence of aqueous KOAc prevents formation of Pd<sub>3</sub>-complex **33** and that  $K_2[Pd(CN)_4]$ , which is most likely produced from **33** under these conditions, is inactive as a catalyst precursor.†‡



**Scheme 10** Pd-Catalysed cyanation of aryl bromides using **32** instead of  $P(tBu)_3$ -Pd-G3. <sup>a</sup>Chamber A: 4-bromobiphenyl (1.0 mmol), **33** (2.5 mol%) and KOAc (3.0 mmol) in dioxane (1.0 mL) and H<sub>2</sub>O (2.0 mL). Chamber B: KCN (1.5 mmol), ethylene glycol (1.0 mL) and AcOH (9.0 mmol).



**Scheme 11** Robustness screening on the formation of benzonitrile **9**. <sup>a</sup>Chamber A: 4-bromophenol (1.0 mmol), 2-acetyl-5-bromothiophene (1.0 mmol)  $P(tBu)_3$ -Pd-G3 (2.5 mol%) and KOAc (3.0 mmol) in dioxane (1.0 mL) and H<sub>2</sub>O (2.0 mL). Chamber B: KCN (1.5 mmol), ethylene glycol (1.0 mL) and AcOH (9.0 mmol).



**Scheme 12** Formation of  $Pd_2(\mu-Br)_2(P(tBu)_3)_2$  and 5,5'-diacetyl-2,2'-bithiophene from  $Pd(P(tBu)_3)_2$ . <sup>a</sup>Reaction performed on a 0.1 mmol scale.

Finally, while the substrate scope of the Pd-catalysed cyanation of aryl bromides using HCN proved broad, some heteroaryl bromides provided low or even no conversion to product. One of these inactive bromides, 2-acetyl-5-bromothiophenyl bromide, was added to the cyanation reaction of 4-bromophenol, which under our standard condition with HCN and in the absence of the heteroaryl bromide provided the corresponding 4-cyanophenol in high yield as depicted in Scheme 3.<sup>45</sup> §§ However, under the exact same conditions but with the addition of this inactive heteroaryl bromide, the formation of benzonitrile **9** was completely inhibited (Scheme 11). Next, we attempted to isolate the oxidative addition complex of  $Pd(P(tBu)_3)_2$  and 2-acetyl-5-bromothiophene in order to test the stoichiometric reaction with Pd-hydride **32**. Despite extensive experimentation, *i.e.* different solvents and temperatures, we only observed the formation of  $Pd_2(\mu-Br)_2(P(tBu)_3)_2$  (**38**) and 5,5'-diacetyl-2,2'-bithiophene (**39**) in all cases (Scheme 12).

The Pd-complex **38** has previously been synthesised by Hartwig and co-workers by a similar protocol using  $Pd(dba)_2$ ,  $P(tBu)_3$  and a tenfold excess of 2-bromothiophene.<sup>46</sup> The formation of **38** is fast at 25 °C when using either THF or pentane. Given that the developed conditions for the cyanation of aryl bromides operate at 60 °C, production of **38** is fast leading to apparent shutdown of the catalytic system. The formation of complex **38** offers an explanation to why some heteroaromatic bromides fail under the developed cyanation protocol, however further investigations are needed to understand the reasons for this divergence.<sup>47,48</sup>

## Conclusions

In summary, a new protocol for the direct use of stoichiometric gaseous hydrogen cyanide in the Pd-catalysed cyanation of aryl



bromides has been developed. Contrary to previous studies, the use of HCN did not lead to catalytic shutdown, but instead provided a robust and reproducible method. A broad range of aryl bromides and a few heteroaromatic bromides afforded the desired benzonitrile derivatives and given the simple setup, utilising *ex situ* generation of HCN, labelling with  $H^{13}CN$  was also straightforward. The presence of water as co-solvent and the use of the mild base KOAc proved imperative for catalytic efficiency. In particular, the suitability of this weak base indicated that possibly the mechanism in operation deviates from previous catalytic cyanation studies as the concentration of free cyanide would be virtually non-existent. This led to the proposal of a mechanism based on a transmetallation between two Pd-complexes produced from the oxidative addition of Pd(0) into hydrogen cyanide and an aryl bromide. This proposal was based on mechanistic indications that co-aligns with observations reported by Grushin, Beller and others. Further work is now ongoing to examine a similar protocol under nickel catalysis, as well as examining other electrophiles than aryl halides. The results of this work will be reported in due course.

## Conflicts of interest

Troels Skrydstrup and Anders T. Lindhardt are co-owners of SyTracks a/s, which commercialises the two-chamber technology.

## Acknowledgements

This study was funded by the BioValue SPIR, Strategic Platform for Innovation and Research on value added products from biomass, which is co-funded by The Innovation Fund Denmark, case no: 0603-00522B. We are also deeply appreciative of generous financial support from Haldor Topsøe A/S, the Danish National Research Foundation (grant no. DNRF118 and DNRF93) and Aarhus University.

## Notes and references

‡ According to the Sigma-Aldrich catalogue, 5 g of acetone cyanohydrin costs the same as approximately 100 g of potassium cyanide.

§  $H^{13}CN$  was used because this isotope eased the spectroscopic analysis of the complexes formed.

¶ In the non-isotopically labelled version, the multiplicity is seen as a triplet.

|| Complex 33 or the corresponding potassium salt ( $K_2[Pd(^{13}CN)_4]$ ) could not be detected by either  $^{31}P$  NMR or  $^{13}C$  NMR spectroscopic analysis.

\*\* Compound 34 has previously been synthesised by Grushin and co-workers by the reaction of  $Pd(PPh_3)_4$  with  $K^{13}CN$  in the presence of water. See ref. 26.

†† During the reaction, no clear phosphine signals are observed from either the oxidative addition complex (*i.e.*  $P(tBu)_3Pd(Ar)(Br)$ ) or the complex after transmetallation (*i.e.*  $P(tBu)_3Pd(Ar)(CN)$ ). This further indicates that the rate determining step is the oxidative addition into the aryl bromide, since the generated  $P(tBu)_3Pd(Ar)(Br)$  is consumed almost instantaneously (see J. L. Klinckenberg and J. F. Hartwig, *J. Am. Chem. Soc.*, 2012, **134**, 5748, which studies the reductive elimination of  $L_nPd(Ar)(CN)$  complexes).

‡‡  $K_2[Pd(CN)_4]$  was also tested. This resulted in no conversion to benzonitrile 11.

§§ Lowering the amount of 2-acetyl-5-bromothiophenyl bromide to 0.5 equivalents or using other heteroaromatic bromides, such as 5-bromopyridine, gave the same outcome.

- (a) P. Pollak, G. Romeder, F. Hagedorn and H. P. Gelbke, *Nitriles, Ullmann's Encyclopedia of Industrial Chemistry*, Wiley-VCH, June 15, 2000, [http://onlinelibrary.wiley.com/doi/10.1002/14356007.a17\\_363/full](http://onlinelibrary.wiley.com/doi/10.1002/14356007.a17_363/full); (b) J. S. Miller and J. L. Manson, *Acc. Chem. Res.*, 2001, **34**, 563; (c) A. Kleemann, J. Engel, B. Kutscher and D. Reichert, *Pharmaceutical substances: Syntheses, Patents, Applications of the most relevant APIs*, Georg Thieme Verlag, Stuttgart, New York, 2001; (d) L. J. Goujon, A. Khaldi, A. Maziz, C. Plesse, G. T. M. Nguyen, P.-H. Aubert, F. Vidal, C. Chevrot and D. Teyssié, *Macromolecules*, 2011, **44**, 9683; (e) F. F. Fleming, *Nat. Prod. Rep.*, 1999, **16**, 597; (f) C. Torborg and M. Beller, *Adv. Synth. Catal.*, 2009, **351**, 3027.
- Z. Rappoport, *The Chemistry of the Cyano Group*, Interscience Publisher, New York, 1970.
- (a) K. W. Rosenmund and E. Struck, *Chem. Ber.*, 1919, **52**, 1749; (b) J. Lindley, *Tetrahedron*, 1984, **40**, 1433; (c) T. Sandmeyer, *Ber. Dtsch. Chem. Ges.*, 1884, **17**, 1633; (d) H. H. Hodgson, *Chem. Rev.*, 1947, **40**, 251; (e) J. K. Kochi, *J. Am. Chem. Soc.*, 1957, **79**, 2942.
- For a recent review see: (a) Q. Wen, J. Jin, L. Zhang, Y. Luo, P. Lu and P. Wang, *Tetrahedron Lett.*, 2014, **55**, 1271; for selected examples, see: (b) J. Zanon, A. Klapars and S. L. Buchwald, *J. Am. Chem. Soc.*, 2003, **125**, 2890; (c) H.-J. Cristau, A. Ouali, J.-F. Spindler and M. Taillefer, *Chem.-Eur. J.*, 2005, **11**, 2483; (d) T. Schareina, A. Zapf, W. Mägerlein, N. Müller and M. Beller, *Chem.-Eur. J.*, 2007, **13**, 6249; (e) T. Schareina, A. Zapf, A. Cotté, M. Gotta and M. Beller, *Adv. Synth. Catal.*, 2011, **353**, 777.
- (a) L. Cassar, *J. Organomet. Chem.*, 1973, **54**, C57; (b) L. Cassar, S. Ferrara and M. Foa, *Adv. Chem. Ser.*, 1974, **132**, 252; (c) L. Cassar, M. Foa, F. Montanari and G. P. Marinelli, *J. Organomet. Chem.*, 1979, **173**, 335; (d) Y. Sakakibara, F. Okuda, A. Shimobayashi, K. Kirino, M. Sakai, N. Uchino and K. Takagi, *Bull. Chem. Soc. Jpn.*, 1988, **61**, 1985; (e) K. Takagi and Y. Sakakibara, *Chem. Lett.*, 1989, **18**, 1957; (f) K. Sasaki, M. Sakai, Y. Sakakibara and K. Takagi, *Chem. Lett.*, 1991, **20**, 2017; (g) Y. Sakakibara, Y. Ido, K. Sasaki, M. Saki and N. Uchino, *Bull. Chem. Soc. Jpn.*, 1993, **66**, 2776; (h) Y. Sakakibara, K. Sasaki, F. Okuda, A. Hokimoto, T. Ueda, M. Sakai and K. Takagi, *Bull. Chem. Soc. Jpn.*, 2004, **77**, 1013; (i) R. K. Arvela and N. E. Leadbeater, *J. Org. Chem.*, 2003, **68**, 9122; (j) X. Zhang, A. Xia, H. Chen and Y. Liu, *Org. Lett.*, 2017, **19**, 2118.
- (a) K. Takagi, T. Okamoto, Y. Sakakibara and S. Oka, *Chem. Lett.*, 1973, **2**, 471; (b) A. Sekiya and N. Ishikawa, *Chem. Lett.*, 1975, **4**, 277; (c) K. Takagi, T. Okamoto, Y. Sakakibara, A. Ohno, S. Oka and N. Hayama, *Bull. Chem. Soc. Jpn.*, 1975, **48**, 3298; (d) K. Takagi, T. Okamoto, Y. Sakakibara, A. Ohno, S. Oka and N. Hayama, *Bull. Chem. Soc. Jpn.*, 1976, **49**, 3177; (e) Y. Akita, M. Shimazaki and A. Ohta, *Synthesis*, 1981, 974; (f) K. Takagi, K. Sasaki and Y. Sakakibara, *Bull. Chem. Soc. Jpn.*, 1991, **64**, 1118; (g) B. A. Andersen, E. C. Bell, F. O. Ginah, N. K. Harn, L. M. Pagh and J. P. Wepsiec, *J. Org. Chem.*, 1998, **63**, 8224;



- (h) M. Sundermeier, A. Zapf, S. Mutyala, W. Baumann, J. Sans, S. Weiss and M. Beller, *Chem.–Eur. J.*, 2003, **9**, 1828.
- 7 For a recent review on Pd-catalysed cyanation of aryl halides covering the extensive work by Beller, see: P. Anbarasan, T. Scareina and M. Beller, *Chem. Soc. Rev.*, 2011, **40**, 5049; see also references listed herein.
- 8 A. V. Ushkov and V. V. Grushin, *J. Am. Chem. Soc.*, 2011, **133**, 10999.
- 9 (a) T. D. Senecal, W. Shu and S. L. Buchwald, *Angew. Chem., Int. Ed.*, 2013, **52**, 10053; (b) D. T. Cohen and S. L. Buchwald, *Org. Lett.*, 2015, **17**, 202.
- 10 T. Okano, M. Iwahara and J. Kiji, *Synlett*, 1998, 243.
- 11 M. Sundermeier, M. Zapf, M. Beller and J. Sans, *Tetrahedron Lett.*, 2001, **42**, 6707.
- 12 (a) N. Chatani and T. Hanafusa, *J. Org. Chem.*, 1986, **51**, 4714; (b) M. Sundermeier, S. Mutyala, A. Zapf, A. Spannenberg and M. Beller, *J. Organomet. Chem.*, 2003, **684**, 50.
- 13 M. Sundermeier, A. Zapf and M. Beller, *Angew. Chem., Int. Ed.*, 2003, **42**, 1661.
- 14 (a) T. Schareina, A. Zapf and M. Beller, *Chem. Commun.*, 2004, 1388; (b) D. Zhang, H. Sun, L. Zhang, Y. Zhou, C. Li, H. Jiang, K. Chen and H. Liu, *Chem. Commun.*, 2012, **48**, 2909; (c) S. A. Weissman, D. Zewge and C. Chen, *J. Org. Chem.*, 2005, **70**, 1508; (d) O. Grossman and D. Gelman, *Org. Lett.*, 2006, **8**, 1189; (e) T. Schareina, A. Zapf, W. Mägerlein, N. Müller and M. Beller, *Tetrahedron Lett.*, 2007, **48**, 1087; (f) T. Schareina, R. Jackstell, T. Schulz, A. Zapf, A. Cotté, M. Gotta and M. Beller, *Adv. Synth. Catal.*, 2009, **351**, 643.
- 15 (a) D. M. Tschaen, R. Desmond, A. O. King, M. C. Fortin, B. Pipik, S. King and T. R. Verhoeven, *Synth. Commun.*, 1994, **24**, 887; (b) P. E. Maligres, M. S. Waters, F. Fleitz and D. Askin, *Tetrahedron Lett.*, 1999, **40**, 8193; (c) M. Alterman and A. Hallberg, *J. Org. Chem.*, 2000, **65**, 7984; (d) R. Chidambaram, *Tetrahedron Lett.*, 2004, **45**, 1441; (e) R. S. Jensen, A. S. Gajare, K. Toyota, M. Yoshifuji and F. Ozawa, *Tetrahedron Lett.*, 2005, **46**, 8645; (f) M. T. Martin, B. Liu, B. E. Cooley Jr and J. F. Eaddy, *Tetrahedron Lett.*, 2007, **48**, 2555.
- 16 T. Sakamoto and K. Ohsawa, *J. Chem. Soc., Perkin Trans. 1*, 1999, 2323.
- 17 D. Andrussow, *Angew. Chem.*, 1935, **48**, 593.
- 18 H. A. Wittcoff, B. G. Reuben and J. S. Plotkin, *Industrial Organic Chemicals*, John Wiley And Sons Ltd, 2013.
- 19 Areas of large scale-application for HCN includes the DuPont Adiponitrile process (monomer in the synthesis of Nylon 6,6), which annually produces around 1 million tons of adiponitrile, see: L. Bini, C. Müller and D. Vogt, *ChemCatChem*, 2010, **2**, 590.
- 20 For excellent examples on the development and application of transfer hydrocyanations, see: (a) X. Fang, P. Yu and B. Morandi, *Science*, 2016, **351**, 832; (b) X. Fang, P. Yu, G. Prina Cerai and B. Morandi, *Chem.–Eur. J.*, 2016, **22**, 15629; (c) B. N. Bhawal and B. Morandi, *ACS Catal.*, 2016, **6**, 7528.
- 21 H. G. Lee, J. P. Milner, M. S. Placzek, S. L. Buchwald and J. M. Hooker, *J. Am. Chem. Soc.*, 2015, **137**, 648.
- 22 W. Zhao, H. G. Lee, S. L. Buchwald and J. M. Hooker, *J. Am. Chem. Soc.*, 2017, **139**, 7152.
- 23 G. Romeder, *Hydrogen Cyanide, e-EROS Encyclopedia of Reagents for Organic Synthesis*, 2001.
- 24 For a recent review on deactivation of homogeneous transition metal catalysis see: R. H. Crabtree, *Chem. Rev.*, 2015, **115**, 127.
- 25 K. D. Dobbs, W. J. Marshall and V. V. Grushin, *J. Am. Chem. Soc.*, 2007, **129**, 30.
- 26 S. Erhardt, V. V. Grushin, A. H. Kilpatrick, S. A. Macgregor, W. J. Marshall and C. R. Roe, *J. Am. Chem. Soc.*, 2008, **130**, 4828.
- 27 (a) P. Hermange, A. T. Lindhardt, R. H. Taaning, K. Bjerglund, D. Lupp and T. Skrydstrup, *J. Am. Chem. Soc.*, 2011, **133**, 6061; (b) S. D. Friis, A. T. Lindhardt and T. Skrydstrup, *Acc. Chem. Res.*, 2016, **49**, 594.
- 28 (a) A. Modvig, T. L. Andersen, R. H. Taaning, A. T. Lindhardt and T. Skrydstrup, *J. Org. Chem.*, 2014, **79**, 5861; (b) K. T. Neumann, S. Klimczyk, M. N. Burhardt, B. Bang-Andersen, T. Skrydstrup and A. T. Lindhardt, *ACS Catal.*, 2016, **6**, 4710.
- 29 G. K. Min, K. Bjerglund, S. Kramer, T. M. Gøsgis, A. T. Lindhardt and T. Skrydstrup, *Chem.–Eur. J.*, 2013, **19**, 17603.
- 30 Hydrocyanations using HCN: (a) A. L. Casalnuovo, T. V. RajanBabu, T. A. Ayers and T. H. Warren, *J. Am. Chem. Soc.*, 1994, **116**, 9869; (b) L. Bini, C. Müller, J. Wilting, L. von Chrzanowski, A. L. Spek and D. Vogt, *J. Am. Chem. Soc.*, 2007, **129**, 12622; (c) J. Wilting, M. Janssen, C. Müller, M. Lutz, A. L. Spek and D. Vogt, *Adv. Synth. Catal.*, 2007, **349**, 350; (d) L. Bini, E. A. Pidko, C. Müller, R. A. van Santen and D. Vogt, *Chem.–Eur. J.*, 2009, **15**, 8768; (e) A. Falk, A. Cavalieri, G. S. Nichol, D. Vogt and H.-G. Schmalz, *Adv. Synth. Catal.*, 2015, **357**, 3317.
- 31 Hydrocyanations using HCN-surrogates: (a) M. de Greef and B. Breit, *Angew. Chem., Int. Ed.*, 2009, **48**, 551; (b) A. Falk, A.-L. Göderz and H.-G. Schmalz, *Angew. Chem., Int. Ed.*, 2013, **52**, 1576.
- 32 For recent reviews on Ni-catalysed hydrocyanations: (a) L. Bini, C. Müller and D. Vogt, *Chem. Commun.*, 2010, **46**, 8325; (b) T. V. RajanBabu, *Org. React.*, 2011, **75**, 1.
- 33 F. F. Flemming, L. Yao, P. C. Ravikumar, L. Funk and B. C. Shock, *J. Med. Chem.*, 2010, **53**, 7902.
- 34 F. G. Buono, R. Chidambaram, R. H. Mueller and R. E. Waltermire, *Org. Lett.*, 2008, **10**, 5325.
- 35 For a review on palladium hydrides, see: V. V. Grushin, *Chem. Rev.*, 1996, **96**, 2011.
- 36 (a) R. Vilar, D. M. P. Mingos and C. J. Cardin, *J. Chem. Soc., Dalton Trans.*, 1996, 4313; (b) V. Durà-Vilà, D. M. P. Mingos, R. Vilar, A. J. P. White and D. J. Williams, *J. Organomet. Chem.*, 2000, **600**, 198.
- 37 (a) A. B. Goel and S. Goel, *Inorg. Chim. Acta*, 1980, **45**, L85; (b) R. G. Goel and W. O. Ogini, *Organometallics*, 1982, **1**, 654; (c) I. D. Hills and G. C. Fu, *J. Am. Chem. Soc.*, 2004, **126**, 13178.
- 38 E. H. Brooks and F. Glockling, *J. Chem. Soc. A*, 1967, 1030.
- 39 A. G. Sergeev, A. Spanneberg and M. Beller, *J. Am. Chem. Soc.*, 2008, **130**, 15549.



- 40 It has been earlier shown that the oxidative addition to AcOH is a reversible process: C. Amatore, A. Jutland, G. Meyer, I. Carelli and I. Chiarotto, *Eur. J. Inorg. Chem.*, 2000, 1855.
- 41 The oxidative addition of Pd(P(*t*Bu)<sub>3</sub>) into aryl bromides has previously been shown to be a reversible process: (a) A. H. Roy and J. F. Hartwig, *J. Am. Chem. Soc.*, 2001, **123**, 1232; (b) A. H. Roy and J. F. Hartwig, *J. Am. Chem. Soc.*, 2003, **125**, 13944; (c) A. H. Roy and J. F. Hartwig, *Organometallics*, 2004, **23**, 1533.
- 42 A. D. Littke, C. Dai and G. C. Fu, *J. Am. Chem. Soc.*, 2000, **122**, 4020.
- 43 (a) F. Ozawa, M. Fujimori, T. Yamamoto and A. Yamamoto, *Organometallics*, 1986, **5**, 2144; (b) A. L. Casado, J. A. Casares and P. Espinet, *Organometallics*, 1997, **16**, 5730; (c) Y. Tan, F. Barrios-Landeros and J. F. Hartwig, *J. Am. Chem. Soc.*, 2012, **134**, 3683; (d) D. Wang, Y. Izawa and S. S. Stahl, *J. Am. Chem. Soc.*, 2014, **136**, 9914; (e) D. Wang and S. S. Stahl, *J. Am. Chem. Soc.*, 2017, **139**, 5704.
- 44 J. P. Stambuli, C. D. Incarvito, M. Bühl and J. F. Hartwig, *J. Am. Chem. Soc.*, 2004, **126**, 1184.
- 45 K. D. Collins and F. Glorius, *Nat. Chem.*, 2013, **5**, 597.
- 46 M. W. Hooper, M. Utsunomiya and J. F. Hartwig, *J. Org. Chem.*, 2003, **68**, 2861.
- 47 The formation and role of **38** has recently been elucidated by Schoenebeck and Colacot. See: C. C. C. Johansson Seechurn, T. Sperger, T. G. Scrase, F. Schoenebeck and T. J. Colacot, *J. Am. Chem. Soc.*, 2017, **139**, 5194.
- 48 For other examples from the Schoenebeck group exploiting **38** and its iodo-analogue as a Pd(0) precatalyst, see: (a) F. Proutiere, M. Aufiero and F. Schoenebeck, *J. Am. Chem. Soc.*, 2012, **134**, 606; (b) K. J. Bonney, F. Proutiere and F. Schoenebeck, *Chem. Sci.*, 2013, **4**, 4434; (c) G. Yin, I. Kalvet and F. Schoenebeck, *Angew. Chem., Int. Ed.*, 2015, **54**, 6809; (d) M. Aufiero, T. Sperger, A. S.-K. Tsang and F. Schoenebeck, *Angew. Chem., Int. Ed.*, 2015, **54**, 10322; (e) T. Sperger, C. K. Stirner and F. Schoenebeck, *Synthesis*, 2017, **49**, 115; (f) I. Kalvet, G. Magnin and F. Schoenebeck, *Angew. Chem., Int. Ed.*, 2017, **56**, 1581; (g) I. Kalvet, T. Sperger, T. Scattolin, G. Magnin and F. Schoenebeck, *Angew. Chem., Int. Ed.*, 2017, **56**, 7078.

