Cite this: *Soft Matter*, 2017, **13**, 5085

Correction: Phase behavior of colloid–polymer depletion mixtures with unary or binary depletants

Nayoung Park and Jacinta C. Conrad *

DOI: 10.1039/c7sm90120h

Correction for 'Phase behavior of colloid–polymer depletion mixtures with unary or binary depletants' by Nayoung Park *et al.*, *Soft Matter*, 2017, **13**, 2781–2792.

rsc.li/soft-matter-journal

The authors would like to correct errors in Fig. 3 and 4. The x-axes of Fig. 3d–f and 4d–f should have no units, and the x-axis labels of Fig. 4a–c should be $r/2a$. Furthermore, there was a data set missing in Fig. 3f, for $\phi = 0.45$ and $C_p/C_p^* = 2.95$, and a data set missing in Fig. 4d, for $\phi = 0.05$ and $C_{p,L}/C_{p,L}^* = 0$. The correct versions of Fig. 3 and 4 are shown below. These corrections affect neither the calculations nor the conclusions of the manuscript. Finally, Nayoung Park would like to acknowledge partial support from the Houston Endowment Fund.

Fig. 3 (a–c) Radial distribution function $g(r)$ as a function of normalized radial distance $r/2a$ and (d–f) normalized mean squared displacement $MSD/(2a)^2$ as a function of non-dimensionalized delay time $\tau D/a^2$, for PMMA suspensions with various concentrations of 328.9 kDa PS and particle volume fractions of (a and d) $\phi \approx 0.05$, (b and e) $\phi \approx 0.25$, or (c and f) $\phi \approx 0.45$. The dashed lines in (a–c) indicate the limiting value of $g(r)$ at large r . The inset in (d) shows $MSD/(2a)^2$ as a function of delay time τ for suspensions with volume fraction of $\phi \approx 0.05$ without the viscosity correction. The dashed lines in (d–f) indicate the resolution of the measurement $\epsilon^2/(2a)^2$.

Fig. 4 (a–c) Radial distribution function $g(r)$ as a function of normalized radial distance $r/2a$ and (d–f) normalized mean squared displacement $MSD/(2a)^2$ as a function of non-dimensionalized delay time $\tau D/a^2$, for PMMA suspensions with various concentrations of 328.9 kDa PS ($C_{p,L}/C_{p,L}^*$) and fixed concentration of 6400 Da PS ($C_{p,S}/C_{p,S}^*$). The particle volume fractions are (a and d) $\phi \approx 0.05$, (b and e) $\phi \approx 0.25$, or (c and f) $\phi \approx 0.45$. The dashed lines in (a–c) indicate the limiting value of $g(r)$ at large r ; the dashed lines in (d–f) indicate the resolution of the tracking algorithm $\epsilon^2/(2a)^2$.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.