

CORRECTION

[View Article Online](#)
[View Journal](#) | [View Issue](#)



Cite this: *J. Mater. Chem. A*, 2017, 5, 14444

DOI: 10.1039/c7ta90131c
www.rsc.org/MaterialsA

Correction: High surface area, amorphous titania with reactive Ti^{3+} through a photo-assisted synthesis method for photocatalytic H_2 generation

Dennis Zywitzki,^a Hangkun Jing,^b Harun Tüysüz*^a and Candace K. Chan*^b

Correction for 'High surface area, amorphous titania with reactive Ti^{3+} through a photo-assisted synthesis method for photocatalytic H_2 generation' by Dennis Zywitzki *et al.*, *J. Mater. Chem. A*, 2017, 5, 10957–10967.

The authors would like to replace Fig. 4B with the corrected version, shown below.




Fig. 4 Photocatalytic H_2 production rates from samples prepared using $Ti(OEt)_4$. (B) H_2 evolution of Pt-coated DI-OEt samples compared to DI-OEt without Pt.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aDepartment of Heterogeneous Catalysis, Max-Planck-Institut für Kohlenforschung, Mülheim an der Ruhr, Germany. E-mail: tueysuez@kofo.mpg.de

^bMaterials Science and Engineering, School for Engineering of Matter, Transport and Energy, Arizona State University, Tempe, AZ, USA. E-mail: candace.chan@asu.edu