Chemical Science



EDGE ARTICLE

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2018, 9, 634

Gold-catalyzed stereoselective dearomatization/ metal-free aerobic oxidation: access to 3substituted indolines/oxindoles†

Kai Liu, Guangyang Xu and Jiangtao Sun 10 *

An unprecedented dearomatization of indoles with diazoesters has been developed *via* cationic gold(I) catalysis. The functionalization selectively occurs at the C3-position to deliver methylene indole derivatives in good yields with excellent *Z*-selectivity, demonstrating unusual reactivity and selectivity compared with other noble metal catalysis. Importantly, simply followed by silica gel adsorption, an unprecedented metal-free aerobic oxidation occurs for indoles bearing N-electron donating substituents, providing a novel and efficient approach towards 3-substituted indolin-2-ones with a newly formed quaternary stereocenter in excellent stereoselectivity. Notably, these processes afford direct and selective access to a variety of valuable intermediates from abundant feedstock chemicals.

Received 18th September 2017 Accepted 4th November 2017

DOI: 10.1039/c7sc04086e

rsc.li/chemical-science

Introduction


Transition-metal-catalyzed carbene transfer from diazo compounds represents a powerful tool in modern organic synthesis, allowing the rapid assembly of a range of valuable structures which cannot be easily achieved by other methodologies.1 Particularly, different metal-carbenes often exhibit distinct reactivity and selectivity toward the same reaction precursors, which increases molecular complexity. A good example is the reaction of diazoesters with 2,3-nonsubstituted indoles, which often leads to two principal products, namely cyclopropane derivatives (Scheme 1a)² and formal C(sp²)-H insertion products (Scheme 1b).3-7 Indeed, for the addition reaction, the use of rhodium,3 copper,4 iron5 and palladium6 complexes selectively afforded C3-alkylation products. In contrast, C2-alkylation has been observed for 1H-indoles upon exposure to a ruthenium catalyst.7 Additionally, annulation8 and the N-H insertion reaction have also been reported.

On the other hand, recent literature disclosed that the reactivity and chemoselectivity of gold-carbene highly depended on the electronic and steric properties of the ancillary ligand to the gold center, 10,11 as well as the choice of counterion. 12 Variation of the gold catalysts has recently been shown to even allow the formation of gold(III) intermediates from gold(I) precursors. 13 In 2014, Shi and co-workers described a ligand-controlled gold-catalyzed addition of arenes to α -aryl

Jiangsu Key Laboratory of Advanced Catalytic Materials & Technology, School of Petrochemical Engineering, Changzhou University, Changzhou 213164, P. R. China. E-mail: jtsun08@gmail.com; jtsun@cczu.edu.cn

 \dagger Electronic supplementary information (ESI) available. CCDC 1551068, 1551070, 1572912 and 1572913. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7sc04086e

diazoesters.¹⁴ They mentioned one example of indole C3-alkylation in the presence of an electron-deficient phosphite-gold catalyst, which exhibited similar chemo- and site-selectivity to rhodium and copper catalysis (Scheme 1b). However, we recently found that N-heterocyclic carbene (NHC) gold complexes displayed inverse chemoselectivity to phosphite gold catalysts even for the same substrates.^{15b} Thus, we envisioned that, when exposed to different gold catalysts, the reaction of indoles with diazo compounds would probably result in distinct reaction pathways. In continuation with our interests in gold-carbene chemistry,¹⁵ as anticipated but not expected,



Scheme 1 Functionalization of indoles by metal-carbene transfer from diazo compounds: previous reports and our discovery.

Edge Article Chemical Science

herein we report the unprecedented stereoselective dearomatization of indoles with diazoesters under cationic gold(1) catalysis (Scheme 1c). Furthermore, when N-electron-donating substituted indoles are utilized, an unprecedented metal-free aerobic oxidation occurs after the initial dearomatization.

Results and discussion

At the outset, we employed N-boc indole 1a, N-benzyl indole 2a and phenyl diazoacetate 3a as model substrates to investigate the reaction (Table 1). The use of Ph₃PAuCl/AgSbF₆ (5 mol%) in dichloromethane at room temperature afforded C3-insertion product 5a in 70% yield (entry 1), while JohnPhosAuCl/AgSbF₆ gave 16% yield of 5a (entry 2). When t-BuXPhosAuCl was used, 10% yield of 4a was obtained although 5a was still the major product (entry 3). Not surprisingly, the electron-deficient phosphite gold complex only gave 5a as a single product (entry 4). Gratifyingly, the use of IPrAuCl/AgSbF₆ provided 4a in 70% yield (entry 5). We therefore started to survey other NHC


Table 1 Optimization of the reaction conditions^a

Entry	Indole	Catalyst	Solvent	Yield ^b (%)	
				4a(5a)	6a
1	1a	Ph ₃ PAuCl/AgSbF ₆	CH ₂ Cl ₂	0(70)	_
2	1a	JohnPhosAuCl/AgSbF ₆	CH_2Cl_2	0(16)	_
3	1a	t-BuXPhosAuCl/AgSbF ₆	CH_2Cl_2	10(61)	_
4	1a	(ArO) ₃ PAuCl/AgSbF ₆	CH_2Cl_2	0(75)	_
5	1a	IPrAuCl/AgSbF ₆	CH_2Cl_2	70(13)	_
6	1a	IPrAu(PhCN)SbF ₆	CH_2Cl_2	79(10)	_
7	1a	IPrAu(PhCN)BAr _F	CH_2Cl_2	81(8)	_
8	1a	IPrAu(PhCN)BAr _F	DCE	81(9)	_
9	1a	IPrAu(PhCN)BAr _F	CHCl ₃	87(<5)	_
10	1a	IPrAu(PhCN)BAr _F	Toluene	70(15)	_
11	1a	IPrAu(PhCN)BAr _F	THF	25(<5)	_
12	2a	IPrAu(PhCN)BAr _E	CHCl ₃	_ ` ´	77
13	2a	IPrAu(PhCN)BAr _F	CH_2Cl_2	_	70
14	2a	IPrAu(PhCN)BAr _F	DCE	_	65
15	2a	IPrAu(PhCN)BAr _F	THF	_	50

^a Reaction conditions: 1a or 2a (0.2 mmol) and 3a (0.3 mmol) in 2 mL of the solvent were added to a solution of 5 mol% gold catalyst in 2 mL of the solvent via a syringe pump under argon for 2 h. The mixture was stirred at rt for another 2 h. For 6a, silica gel (5 g) adsorption of crude products was performed and was kept in air for 12 h at rt. ^b Isolated yields. (ArO) = (2,4-di-tert-butylphenyl).

gold complexes as well so as to try different counterions. To our delight, the yield of 4a was improved to 79% by IPrAu(PhCN) SbF₆ (entry 6), and was further increased to 81% by IPrAu(PhCN)BAr_F¹⁶ (entry 7). A screen of the solvents revealed chloroform was the best one (entries 8 to 11), providing 4a in 87% yield (entry 9), and no significant side products were detected. The structure of 4a was confirmed with NMR spectroscopy and was further identified using X-ray analysis.17 Notably, when N-benzyl indole 2a was subjected to this reaction, the corresponding 3-methyleneindoline was not obtained and 3-substituted indolin-2-one 6a was obtained instead in 77% yield (entry 12, see ESI† for details), indicating that an aerobic oxidation occurred. Variation of the solvents did not improve the reaction (entries 13 to 15).

We then started to explore the substrate scope for the first dearomatization (Scheme 2). The reaction of indole 1a with a variety of diazoesters was firstly examined. It was observed that methyl phenyl diazoacetate gave the corresponding product in better yield than other esters (4a to 4d). Aromatic diazoesters with different substituents were employed, providing the corresponding 3-methyleneindolines in moderate



Scheme 2 Substrate scope and reaction conditions: to a solution of 5 mol% of IPrAu(PhCN)BAr_F in CHCl₃ (2 mL) a solution of 1 (0.2 mmol) and 3 (0.3 mmol) in 2 mL CHCl₃ was added at rt for 2 h via a syringe pump under argon. The resulting mixture was stirred at rt for another 2 h. Isolated yields have been listed. 4 equiv. of the diazo substrate were used for the preparation of 4q

Chemical Science Edge Article

to excellent yields (4e to 4k). In general, electron-deficient aromatic diazoesters furnished the products (4e to 4i) in higher yields than electron-rich substrates (4j to 4k). Next, the scope of indoles was investigated. Gratifyingly, C5-, C6- and C7substituted N-boc indoles bearing either electron-withdrawing or electron-donating groups were all tolerated, furnishing the corresponding indolines in moderate to excellent yields (41 to 4p). Furthermore, different N-substituted indoles were also examined. The protecting groups such as tosyl (Ts), benzyloxycarbonyl (Cbz), benzoyl (Bz) and acetyl (Ac) were all amenable to the reaction and the corresponding products were obtained in acceptable yields (4q to 4t). This protocol was also amenable to pyrroles. The desired dearomatization products were isolated in good yields (4u to 4x). The structure of 4x was further confirmed by single-crystal X-ray crystallography. 17 It should be noted that all of the methylene derivatives were isolated in the single Z-configuration.

Next, we investigated the substrate scope of the tandem reaction towards the formation of 3-substituted indolin-2-ones (Scheme 3). Generally, the reaction of 2a with aromatic diazoesters either bearing electron-donating or electron-withdrawing substituents proceeded smoothly to afford the corresponding products in moderate to good yields (6a to 6h). A longer oxidation time (24 h) is needed for 6e to 6h. Afterwards, indoles bearing various substituents were examined. The use of Nbenzyl 5-methoxy and 5-bromo indoles provided 6i and 6j in


HO. IPrAu(PhCN)BAr_F (5 mol%) CO₂Me Ar CHCl₃; Ν̈́2 then silica gel, air rt, 12-24 h 3 HO. CO₂Me Ŕп Ь'n 6a, 77% 6b (R = Me), 80%, >20:1 d.r. >20·1 d r **6c** (R = OMe), 82%, >20:1 d.r. 6d (R = Br), 76%, >20:1 d.r = CF₃), 65%, >20:1 d.r. CO₂Me CO₂Me CO₂Me o² Β'n В'n **6g**, 70% >20:1 d.r. 6f. 74% 6h. 51% >20:1 d.r. HO, CO₂Me CO₂Me Β'n 6i (R = OMe), 81%, >20:1 d.r. 6k (R = CI), 75%, >20:1 d.r. 6j (R = Br), 74%, >20:1 d.r. 6I (R = Me), 78%, >20:1 d.r X-ray of 61 HO, HO CO₂Me °O Me ḃn 6m 53% 6n 67% 60 59%

Scheme 3 Tandem reaction of the dearomatization and aerobic oxidation. Reaction conditions: IPrAu(PhCN)BAr_F (5 mol%), 2 (0.2 mmol) and $\bf 3$ (0.3 mmol) in 4 mL CHCl₃ at rt for 4 h. Then SiO₂ (5 g) adsorption in air at rt for 12 to 24 h. Isolated yields have been listed.

81% and 74% yield, while 6-chloro and 6-methyl delivered the corresponding products in 75% and 78% yield, respectively. 7-Methoxy N-benzyl indole was also examined, and 6m was isolated in 53% yield. Finally, N-methyl and N-phenyl indoles were tested and the corresponding products (6n and 6o) were obtained in moderate yields. The structure of 6a and 6l was confirmed by X-ray analysis.

Deuterium labeling and control experiments were conducted to understand the reaction mechanism (Scheme 4). First, 4a cannot be converted to 5a under standard reaction conditions (Scheme 4a), which ruled out the possibility of preferential formation of 5 followed by isomerization and vice versa. Next, the reaction of D-1a with 3a yielded D-4a. The high deuterium incorporation at the 2-position might indicate that a 1,2hydrogen shift is likely involved in the reaction (Scheme 4b). Since there is significant loss of deuterium, we suspected that the reaction may be interfered with by adventitious water present in the reaction mixture. Thus, a control reaction with 3 equivalents of D2O was run. Indeed, significant deuterium incorporation into the product was observed (Scheme 4c), suggesting that there is D/H exchange with the adventitious proton source during the reaction's progress. A study on the kinetic isotope effect (KIE) using intermolecular competition between 1a and D-1a indicated a KIE value of 3.35 (Scheme 4d), which is consistent with the 1,2-hydrogen shift being the ratedetermining step. Furthermore, compound 7 was separately prepared and subjected to standard reaction conditions. Unfortunately, no reaction was observed. The chemical incompetence of 7 rules out its possible role as an intermediate in this process (Scheme 4e).

Next, mechanistic studies for the aerobic oxidation were carried out (Scheme 5). First, the reaction of D-2a with 3a gave D-6a in 73% yield with a low ratio of deuterium labeling (Scheme 5a). Moreover, the ¹⁸O-labeled product **6a**' was



Mechanistic studies on the dearomatization reaction.


Edge Article Chemical Science

Scheme 5 Mechanistic studies on the aerobic oxidation.

obtained in 78% yield under an ¹⁸O₂ atmosphere, indicating that the oxygen atom in 6 came from the air (Scheme 5b). Using TBHP (tert-butyl hydroperoxide) instead of air as an oxidant, the reaction was messy although 6a could still be detected by GC/ MS (Scheme 5c). To determine the role of the silica gel, ptoluene sulfonic acid was added and the C-3 alkylation product 8 was obtained in high yield (Scheme 5d). Moreover, by replacing the silica gel with anhydrous MgSO4, 6a was isolated in 68% yield upon exposure to air for 12 h (Scheme 5d). Therefore, the acidic property of the silica gel may not be critically important for this transformation. Its role is presumably to facilitate the aerobic oxidation by increasing the contact surface of the olefin intermediate with oxygen. Just recently, López and co-workers reported the gold-catalyzed formal insertion of aryl diazoesters into ferrocene to generate functionalized metallocenes.18 They also found that the silica gel promoted the aerobic oxidation leading to tertiary-substituted ferrocenyl alcohols. In that case, they believed the aerobic oxidation might proceed through initial electron-transfer from iron to molecular oxygen, which triggered the radical sequence to give the target product. Thus, to determine the reactive intermediate for this oxidation process, the C3-alkylation product 8 was subjected to silica gel in air, however, no reaction occurred (Scheme 5d). Gratifyingly, although the reaction intermediates are extremely unstable, we successfully isolated intermediate 9 after carefully checking all of the reactions listed in Scheme 3. Furthermore, the treatment of 9 with silica gel in

air provided **6h** in 73% yield (Scheme 5e). This result indicated that gold catalysts did not work during the aerobic oxidation. Without silica gel adsorption, **6h** cannot be obtained either. Different from López's report, this aerobic oxidation does not involve metal participation.

Until now, several reaction mechanisms have been proposed for indole functionalization with diazo compounds in the presence of rhodium, copper or other metal complexes.²⁻⁸ Clearly, this gold-catalyzed dearomatization is quite different from these known processes. 19,20 Although the exact reaction mechanism is not clear at this moment, plausible ones have been proposed (Scheme 6). In view of the unique formation of Zolefins, the carboxylate group may assist the olefin selectivity. Partially analogous to Fox's description of rhodium-catalyzed C-3 alkylation of indoles with diazoesters^{3d} and density functional theory (DFT) calculations reported by Xie et al., 3f the reaction of 3a with the cationic gold catalyst first generates gold carbene species IA or IB, which is followed by nucleophilic attack with indole to produce ylide III via transition state II. Then this ylide intermediate would undergo a 1,2-hydrogen shift to give the final dearomatization intermediate IV, together with catalyst regeneration. The assistance of the ester carbonyl group explains the Z-configuration of the observed product. Next, silica gel-assisted aerobic oxidation occurs. The reaction of IV with molecular oxygen generates intermediate V via the Schenck ene reaction.21 Owing to the high oxidation ability of the



Scheme 6 Proposed reaction mechanisms.

peroxide motif and the electron-rich nature of the indole ring, subsequent internal epoxidation can rapidly take place *via* either **VI** or **VII**. Finally, semi-pinacol rearrangement of **VI**²² or rearrangement of the amino epoxide motif in **VII**²³ leads to the observed amide **6**. Considering the high reactivity of peroxide **V**, the epoxidation step occurs on the same face of the indole plane once it is formed (C–C bond rotation is thus discouraged). In this scenario, the stereochemistry integrity determines the high diastereoselectivity observed in the final product.

Conclusions

In summary, we have developed an unprecedented gold-catalyzed stereoselective dearomatization of indoles with diazoesters, providing 3-methyleneindolines in good to excellent yields with unique *Z*-configuration. Moreover, when N-donating substituent indoles were subjected to the reaction, a tandem reaction sequence occurred including the initial dearomatization and a sequential metal-free aerobic oxidation to produce 3-substituted indolin-2-ones. As a result, molecular oxygen has been successfully inlaid into the final structure. Notably, the use of the cationic gold(i) catalyst IPrAu(PhCN)BAr_F is crucial to the whole process.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We gratefully acknowledge the National Natural Science Foundation of China (21572024), the Natural Science Foundation of Jiangsu Province (BK20151184), the Jiangsu Key Laboratory of Advanced Catalytic Materials and Technology (BM2012110) and the Advanced Catalysis & Green Manufacturing Collaborative Innovation Center for their financial support.

Notes and references

- 1 For selected leading reviews, see: (a) H. M. L. Davies and S. J. Hedley, Chem. Soc. Rev., 2007, 36, 1109; (b) M. P. Doyle, R. Duffy, M. Ratnikov and L. Zhou, Chem. Rev., 2010, 110, 704; (c) X. Guo and W. Hu, Acc. Chem. Res., 2013, 46, 2427; (d) Y. Xia, Y. Zhang and J. Wang, ACS Catal., 2013, 3, 2586; (e) X. Xu and M. P. Doyle, Acc. Chem. Res., 2014, 47, 1396; (f) A. Ford, H. Miel, A. Ring, C. N. Slattery, A. R. Maguire and M. A. McKervey, Chem. Rev., 2015, 115, 9981.
- 2 (a) F. Gnad, M. Poleschak and O. Reiser, *Tetrahedron Lett.*, 2004, 45, 4277; (b) S. J. Hedley, D. L. Ventura, P. M. Dominiak, C. L. Nygren and H. M. L. Davies, *J. Org. Chem.*, 2006, 71, 5349; (c) M. Delgado-Rebollo, A. Prieto and P. J. Pérez, *ChemCatChem*, 2014, 6, 2047 and reference therein; (d) V. Lehner, H. M. L. Davies and O. Reiser, *Org. Lett.*, 2017, 19, 4722.
- 3 (a) J. L. Wood, B. M. Stoltz, H.-J. Dietrich, D. A. Pflum and D. T. Petsch, *J. Am. Chem. Soc.*, 1997, **119**, 9641; (b) R. Gibe

- and M. A. Kerr, *J. Org. Chem.*, 2002, **67**, 6247; (c) Y. Lian and H. M. L. Davies, *Org. Lett.*, 2010, **12**, 924; (d) A. DeAngelis, V. W. Shurtleff, O. Dmitrenko and J. M. Fox, *J. Am. Chem. Soc.*, 2011, **133**, 1650; (e) Y. Lian and H. M. L. Davies, *Org. Lett.*, 2012, **14**, 1934; (f) Q. Xie, X.-S. Song, D. Qu, L.-P. Guo and Z.-Z. Xie, *Organometallics*, 2015, **34**, 3112.
- 4 M. B. Johansen and M. A. Kerr, Org. Lett., 2010, 12, 4956.
- 5 Y. Cai, S.-F. Zhu, G.-P. Wang and Q.-L. Zhou, *Adv. Synth. Catal.*, 2011, 353, 2939.
- 6 X. Gao, B. Wu, W.-X. Huang, M.-W. Chen and Y.-G. Zhou, *Angew. Chem., Int. Ed.*, 2015, **54**, 11956.
- 7 W.-W. Chan, S.-H. Yeung, Z. Zhou, A. S. C. Chan and W.-Y. Yu, *Org. Lett.*, 2010, **12**, 604.
- 8 For selected examples, see: (a) Y. Lian and H. M. L. Davies, J. Am. Chem. Soc., 2010, 132, 440; (b) C. Jing, Q.-Q. Cheng, Y. Deng, H. Arman and M. P. Doyle, Org. Lett., 2016, 18, 4550; (c) M. Li, X. Guo, W. Jin, Q. Zheng, S. Liu and W. Hu, Chem. Commun., 2016, 52, 2736; (d) L. Jiang, W. Jin and W. Hu, ACS Catal., 2016, 6, 6146; for intramolecular reactions, (e) A. R. Reddy, F. Hao, K. Wu, C.-Y. Zhou and C.-M. Che, Angew. Chem., Int. Ed., 2016, 55, 1810; (f) M. J. James, P. O'Brien, R. J. K. Taylor and W. P. Unsworth, Angew. Chem., Int. Ed., 2016, 55, 9671; (g) H. Xu, Y.-P. Li, Y. Cai, G.-P. Wang, S.-F. Zhu and Q.-L. Zhou, J. Am. Chem. Soc., 2017, 139, 7697.
- 9 V. Arredondo, S. C. Hiew, E. S. Gutman, I. D. U. A. Premachandra and D. L. Van Vranken, *Angew. Chem., Int. Ed.*, 2017, **56**, 4156.
- 10 For selected reviews, see: (a) A. S. K. Hashmi and G. J. Hutchings, Angew. Chem., Int. Ed., 2006, 45, 7896; (b) A. Fürstner and P. W. Davies, Angew. Chem., Int. Ed., 2007, 46, 3410; (c) D. J. Gorin, B. D. Sherry and F. D. Toste, Chem. Rev., 2008, 108, 3351; (d) A. Corma, A. Leyva-Pérez and M. J. Sabater, Chem. Rev., 2011, 111, 1657; (e) N. Krause and C. Winter, Chem. Rev., 2011, 111, 1994; (f) R. Dorel and A. M. Echavarren, Chem. Rev., 2015, 115, 9028; (g) Z. Zheng, Z. Wang, Y. Wang and L. Zhang, Chem. Soc. Rev., 2016, 45, 4448.
- 11 For recent representative examples on gold-catalyzed diazo reactions, see: (a) V. V. Pagar, A. M. Jadhav and R.-S. Liu, J. Am. Chem. Soc., 2011, 133, 20728; (b) G. Lonzi and L. A. López, Adv. Synth. Catal., 2013, 355, 1948; (c) J. F. Briones and H. M. L. Davies, J. Am. Chem. Soc., 2012, 134, 11916; (d) J. F. Briones and H. M. L. Davies, J. Am. Chem. Soc., 2013, 135, 13314; (e) S. K. Pawar, C.-D. Wang, S. Bhunia, A. M. Jadhav and R.-S. Liu, Angew. Chem., Int. Ed., 2013, 52, 7559; (f) Z.-Y. Cao, X. Wang, C. Tan, X.-L. Zhao, J. Zhou and K. Ding, J. Am. Chem. Soc., 2013, 135, 8197; (g) Z. Yu, B. Ma, M. Chen, H.-H. Wu, L. Liu and J. Zhang, J. Am. Chem. Soc., 2014, 136, 6904; (h) V. V. Pagar and R.-S. Liu, Angew. Chem., Int. Ed., 2015, 54, 4923; (i) Z. Yu, H. Qiu, L. Liu and J. Zhang, Chem. Commun., 2016, 52, 2257; (j) F.-M. Liao, Z.-Y. Cao, J.-S. Yu and J. Zhou, Angew. Chem., Int. Ed., 2017, 56, 2459; (k) B. Ma, Z. Chu, B. Huang, Z. Liu, L. Liu and J. Zhang, Angew. Chem., Int. Ed., 2017, 56, 2749; (l) F. Zhao, N. Li, T. Zhang, Z.-Y. Han,

S.-W. Luo and L.-Z. Gong, *Angew. Chem., Int. Ed.*, 2017, **56**, 3247; (*m*) S. K. Pawar, M.-C. Yang, M.-D. Su and R.-S. Liu, *Angew. Chem., Int. Ed.*, 2017, **56**, 5035. For reviews, see: (*n*) F. Wei, C. Song, Y. Ma, L. Zhou, C.-H. Tung and Z. Xu, *Sci. Bull.*, 2015, **60**, 1479; (*o*) L. Liu and J. Zhang, *Chem. Soc. Rev.*, 2016, **45**, 506; (*p*) M. R. Fructos, M. M. Díaz-Requejo and P. J. Pérez, *Chem. Commun.*, 2016, **52**, 7326.

12 M. Jia and M. Bandini, ACS Catal., 2015, 5, 1638.

Edge Article

- (a) C.-Y. Wu, T. Horibe, C. B. Jacobsen and F. D. Toste, *Nature*, 2015, 517, 449; (b) L. Huang, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Angew. Chem., Int. Ed.*, 2016, 55, 4808; (c) L. Huang, F. Rominger, M. Rudolph and A. S. K. Hashmi, *Chem. Commun.*, 2016, 52, 6435; (d) S. Witzel, J. Xie, M. Rudolph and A. S. K. Hashmi, *Adv. Synth. Catal.*, 2017, 359, 1522; (e) P. T. Bohan and F. D. Toste, *J. Am. Chem. Soc.*, 2017, 139, 11016.
- 14 Y. Xi, Y. Su, Z. Yu, B. Dong, E. J. McClain, Y. Lan and X. Shi, *Angew. Chem.*, *Int. Ed.*, 2014, 53, 9817.
- 15 (a) D. Zhang, G. Xu, D. Ding, C. Zhu, J. Li and J. Sun, Angew. Chem., Int. Ed., 2014, 53, 11070; (b) G. Xu, C. Zhu, W. Gu, J. Li and J. Sun, Angew. Chem., Int. Ed., 2015, 54, 883; (c) C. Zhu, G. Xu and J. Sun, Angew. Chem., Int. Ed., 2016, 55, 11867.
- 16 N. Huguet, D. Lebœuf and A. M. Echavarren, *Chem.-Eur. J.*, 2013, 19, 6581.
- 17 CCDC 1551068 (4a), 1551070 (4x), 1572912 (6a) and 1572913 (6l).†
- 18 When we prepared this work, López and co-workers reported a different iron-promoted aerobic oxidation of ferrocene

- derivatives: E. López, J. Borge and L. A. López, *Chem.-Eur. I.*, 2017, 23, 3091.
- 19 For DFT studies on gold-catalyzed direct arene C-H bond functionalization, see: (a) Y. Liu, Z. Yu, J. Z. Zhang, L. Liu, F. Xia and J. Zhang, *Chem. Sci.*, 2016, 7, 1988; (b) M. R. Fructos, M. Besora, A. A. C. Braga, M. M. Díaz-Requejo, F. Maseras and P. J. Pérez, *Organometallics*, 2017, 36, 172.
- 20 For leading examples on the cyclopropanation reaction, see:
 (a) M. P. Doyle, J. H. Griffin, V. Bagheri and R. L. Dorow, Organometallics, 1984, 3, 53; (b) M. P. Doyle, V. Bagheri, T. J. Wandless, N. K. Harn, D. A. Brinker, C. T. Eagle and K. L. Loh, J. Am. Chem. Soc., 1990, 112, 1906; (c) M. P. Doyle, Q.-L. Zhou, A. B. Dyatkin and D. A. Ruppar, Tetrahedron Lett., 1995, 36, 7579; (d) D. T. Nowlan III, T. M. Gregg, H. M. L. Davies and D. A. Singleton, J. Am. Chem. Soc., 2003, 125, 15902 and reference therein.
- 21 According to the reviewers' suggestions, for selected examples, see: (a) M. Prein and W. Adam, Angew. Chem., Int. Ed., 1996, 35, 477; (b) H. M. R. Hoffmann, Angew. Chem., Int. Ed., 1969, 8, 556; (c) B. M. Snider, Acc. Chem. Res., 1980, 13, 426.
- 22 T. J. Snap, Chem. Soc. Rev., 2007, 36, 1823.
- 23 Y. Zhang, Y. Zou, N. L. Brock, T. Huang, Y. Lan, X. Wang, Z. Deng, Y. Tang and S. Lin, *J. Am. Chem. Soc.*, 2017, 139, 11887.