ChemComm

Chemical Communications

rsc.li/chemcomm

ISSN 1359-7345

COMMUNICATION

Yizhou Liu *et al.*

Enhanced measurement of residual chemical shift anisotropy for small molecule structure elucidation

Cite this: *Chem. Commun.*, 2018, 54, 4254

Received 22nd January 2018,
Accepted 31st January 2018

DOI: 10.1039/c8cc00552d

rsc.li/chemcomm

Enhanced measurement of residual chemical shift anisotropy for small molecule structure elucidation†

Yizhou Liu, ^a Ryan D. Cohen, ^a Kirk R. Gustafson, ^b Gary E. Martin ^a and R. Thomas Williamson ^a

A method is introduced to measure residual chemical shift anisotropies conveniently and accurately in the mesophase of poly- γ -(benzyl-L-glutamate). The alignment amplitude is substantially enhanced over common methods which greatly benefits measurements particularly on sp^3 carbons. The approach offers significant improvements in data accuracy and utility for small molecule structure determination.

Residual chemical shift anisotropy (RCSA) arises from incomplete rotational averaging of chemical shift anisotropy (CSA) as a molecule undergoes partial ordering in an alignment medium.^{1–5} RCSA provides structural information in the form of relative orientations between different CSA tensors in a molecule. Such information is particularly valuable for proton-deficient compounds in which one-bond ^{13}C – ^1H residual dipolar couplings (RDCs) are scarce. Historically, accurate RCSA measurements were plagued by the change in the isotropic component of the chemical shift tensor ($\Delta\delta_{\text{iso}}$) as a molecule is aligned by the addition of an alignment medium. Strategies have recently been introduced to eliminate or correct for $\Delta\delta_{\text{iso}}$ for RCSA data collected in polymeric gels.^{6–8} While RCSAs from these methods, either alone or in conjunction with RDC data, have been successfully applied to the determination of constitution and configuration of a number of small molecules,^{7–11} a limitation exists in that RCSA values tend to be very small for sp^3 -hybridized carbons under these alignment conditions, and differentiation of subtle structural variations involving these carbons is expected to be challenging in the absence of RDC data.

Increasing the alignment amplitude is a logical step to address the problem of small RCSAs. In fact, certain liquid crystal (LC) media, such as that formed by poly- γ -(benzyl-L-glutamate)

(PBLG),¹² are known to generate alignment amplitudes over an order of magnitude larger than constrained polymeric gels.¹³ For RDC measurement, a stronger alignment is not necessarily beneficial, as it simultaneously enlarges undesired interactions such as the homonuclear dipolar couplings, which severely broaden proton lines and degrade the performance of HSQC-type experiments typically used for RDC measurements. As a result, PBLG has only occasionally been used as an alignment medium for acquiring RDC data in structure elucidation studies.^{14–19} In contrast, RCSA measurement only utilizes the simple $^{13}\text{C}\{^1\text{H}\}$ experiment in which inhomogeneous ^{13}C – ^1H dipolar interactions and homogenous interactions from proton spin diffusion are both eliminated by proton decoupling. Therefore, relatively sharp ^{13}C lines can still be obtained even with strong alignment, making accurate RCSA measurement feasible. In fact, the RCSA phenomenon has previously been utilized in the chiral nematic (cholesteric) phase of PBLG for the qualitative differentiation of enantiomeric mixtures.^{20–23} Here we describe a simple but highly effective strategy for the accurate quantitative measurement of RCSA in PBLG mesophase for the purpose of small molecule structural elucidation. Data were collected for strychnine, retrorsine, and caulamidine A (Fig. 1a–c), for which RCSA data were also available for comparison from stretched poly-methylmethacrylate (PMMA) or poly-2-hydroxyethyl-methacrylate (PHEMA) gels.^{8,11}

The challenge for accurate RCSA measurement is to correct for $\Delta\delta_{\text{iso}}$ associated with the PBLG phase transition at different concentrations. Our correction method is based on the premise

Fig. 1 Structures of strychnine (1a), retrorsine (1b), and caulamidine A (1c).

^a Structure Elucidation Group, Process and Analytical Research and Development, Merck & Co. Inc., 126 East Lincoln Avenue, Rahway, NJ 07065, USA.
E-mail: Yizhou.Liu@merck.com

^b Molecular Targets Program, Center for Cancer Research, National Cancer Institute, Frederick, Maryland 21702-1201, USA

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c8cc00552d

that the $\Delta\Delta\delta_{\text{iso}}$ dependence on PBLG concentration is identical in both phases, such that the $\Delta\Delta\delta_{\text{iso}}$ in the mesophase can be corrected for by extrapolating its value from its trend in the isotropic phase. The rationale and limitation of this approach are discussed in ESI†.

We successively added small amounts of PBLG to a strychnine solution and acquired a $^{13}\text{C}\{^1\text{H}\}$ spectrum after each addition (Fig. 2). Tetramethylsilane (TMS) at 4% (v/v) was added for ^{13}C chemical shift referencing. The effect of the bulk susceptibility change at different PBLG concentrations is eliminated by TMS referencing. Although vibration-rotation coupling can generate anisotropic effects in a tetrahedral molecule,^{24–27} TMS aligns much more weakly than most compounds of interest and can therefore be used as a convenient reference in most cases.⁸ Alternatively, RCSA measurements can also be referenced by an arbitrary carbon in the compound itself, following a previously described procedure.⁷ As shown in Fig. 2a, the C21 sp^2 resonance of strychnine gradually shifted upfield as PBLG was added up to 8.8% (w/v) due to $\Delta\Delta\delta_{\text{iso}}$, and no CDCl_3 residual quadrupolar coupling (RQC) was observed consistent with an isotropic solution. As the PBLG concentration reached 11.4%, a sudden large downfield resonance jump occurred due to RCSA (spectrum 5) concomitant with a CDCl_3 RQC of 215.5 Hz, indicating the sample entered the mesophase. Additional PBLG increased both RCSA and RQC as the alignment strengthened. As shown in Fig. S3 (ESI†), $\Delta\Delta\delta_{\text{iso}}$ follows a nearly linear relationship with PBLG concentration within the isotropic phase, although not strictly. The linearity is actually expected if the analyte and PBLG

do not bind (*i.e.*, the binding K_D is very large). This relationship is then used to remove the $\Delta\Delta\delta_{\text{iso}}$ contribution in the mesophase. Importantly, because $\Delta\Delta\delta_{\text{iso}}$ from PBLG addition is small in comparison to RCSA even for sp^3 carbons (Fig. 2b), ignoring the surface interference effect, which is a higher order effect as described in the ESI†, is not expected to cause large errors, at least when the PBLG concentration is not much higher than the mesophase critical concentration (C_{crit}).

Based on the linear relationship, we propose a simplified three-measurement approach, using the terminology: I0–I1–A1. I0 is an isotropic measurement without PBLG, I1 is a second isotropic measurement at a PBLG concentration below but close to C_{crit} , and A1 is an anisotropic measurement at a PBLG concentration above C_{crit} . RCSA can be extracted from these measurements by the simple equation:

$$\text{RCSA}_i = (\delta_{\text{A1}}^i - \delta_{\text{A1}}^{\text{TMS}}) - (\delta_{\text{I0}}^i - \delta_{\text{I0}}^{\text{TMS}}) - \frac{[\text{PBLG}]_{\text{A1}}}{[\text{PBLG}]_{\text{I1}}} [(\delta_{\text{I1}}^i - \delta_{\text{I1}}^{\text{TMS}}) - (\delta_{\text{I0}}^i - \delta_{\text{I0}}^{\text{TMS}})] \quad (1)$$

The first and second terms in parentheses on the right side of eqn (1) represent the TMS-referenced chemical shifts of carbon i in A1 and I0, respectively. Their difference defines the RCSA of carbon i , as well as $\Delta\Delta\delta_{\text{iso}}$, represented by the third term. Note that all chemical shift terms on the right are determined experimentally and the PBLG concentrations are also known. Therefore, the RCSA value of each carbon can be extracted.

As shown in Fig. 2c, the RCSA in PBLG is much larger than the corresponding RCSA in a stretched PMMA gel.⁸ The ~ 2 -fold increase in line-width is amply compensated for by the over 34-fold enhancement of the measured RCSA value. Consequently, these large RCSAs greatly improve the quality of data and structural analysis, as demonstrated later. One noteworthy point is that the proton decoupling used for an isotropic sample may be insufficient to fully decouple the much larger couplings present in mesophase. For example, the alignment with 34.5% PBLG can generate ^{13}C – ^1H RDCs of over 1 kHz so higher decoupling power and a decoupling sequence with wide broadband coverage, such as GARP or WURST, should be used for PBLG mesophase (see ESI† for experimental details).

The data quality is evaluated in Table 1. The generalized degree of order (GDO) was employed to measure the overall magnitude of molecular ordering.²⁸ For a mostly rigid molecule, the GDO measures the alignment amplitude. Our previously reported RCSA data collected in PMMA gels were used for comparison. Clearly, even at relatively low PBLG concentrations, the GDOs are ~ 5 – 6 times larger than in PMMA gels; at the high end of this study, a GDO of 2% was obtained for strychnine, *i.e.*, the anisotropic NMR interactions were measured in a liquid sample at 2% of the solid-state values. To the best of our knowledge, anisotropic interactions of this magnitude have not been previously demonstrated in high-resolution NMR for complex molecules.

RCSAs from these strong alignments yield significantly improved structural analysis, as reflected by the Q -factors. As shown in Table 1, Q -factors with all carbon atoms considered (Q_{all}) are substantially better in PBLG mesophases than in PMMA gels. For example,

Fig. 2 Representative spectra from RCSA measurement. Labelled spectra (0–10) were collected with 0, 2.1, 4.1, 6.6, 8.8, 11.4, 12.9, 15.5, 18.4, 22.9, and 34.5% (w/v) of PBLG in CDCl_3 , respectively. Samples 0–4 were acquired with PBLG below its LC-forming critical concentration and hence are isotropic solution spectra, whereas the data for samples 5–10 were collected with PBLG above its C_{crit} and are increasingly anisotropic in nature. A quaternary sp^2 carbon of strychnine (**1a**) (C21) is shown in (a), and the only quaternary sp^3 carbon in strychnine (C7) is shown in (b). Clearly RCSA shifts in traces 5–10 are significantly larger than the change due to $\Delta\Delta\delta_{\text{iso}}$ seen in traces 0–4, even for C7 (b), which has a very small DFT-computed CSA of 30 ppm. (c) Top traces: C21 of strychnine in weakly (red) and strongly (blue) stretched PMMA gel;⁸ the RCSA value corresponds to the separation between red and blue spectra. Bottom traces: C21 of strychnine with 0% [red, “0” in (a)] and 34.5% PBLG [blue, “10” in (a)]; the actual RCSA value is actually slightly larger than the separation between red and blue spectra after correcting for $\Delta\Delta\delta_{\text{iso}}$. The red spectra in the PMMA gel and PBLG LC solvent are vertically aligned to illustrate the difference in the RCSA size. Although eleven PBLG concentrations were used here to study the trend of chemical shift changes, in practice only three concentrations are needed, *e.g.*, samples 0, 4, and 8 are sufficient for RCSA data extraction by the I0–I1–A1 method described below.

Table 1 RCSA data quality evaluation

	Strychnine (9 sp ² and 12 sp ³ carbons)							Retrorsine ^a (6 sp ² and 10 sp ³ carbons)		
	Stretched PMMA	11.4% PBLG	12.9% PBLG	15.5% PBLG	18.4% PBLG	22.9% PBLG	34.5% PBLG	Stretched PMMA	12.6% PBLG	15.1% PBLG
GDO	5.56×10^{-4}	3.47×10^{-3}	4.67×10^{-3}	6.58×10^{-3}	8.58×10^{-3}	1.21×10^{-2}	2.05×10^{-2}	9.76×10^{-4}	5.04×10^{-3}	7.63×10^{-3}
Q_{all}^b	0.049	0.032	0.032	0.029	0.028	0.029	0.030	0.104	0.040	0.033
$Q_{\text{sp}^3}^c$	0.122	0.060	0.072	0.078	0.074	0.080	0.086	0.184	0.104	0.095
θ^d	12.0°	7.0°	7.4°	7.1°	6.1°	6.6°	7.6°	13.1°	14.0°	4.4°

^a Only data from the more rigid part of retrorsine were used for calculations in this table. RCSAs of C23 and C11 are affected by the rotameric rotation of the primary alcohol across the C23–C11 bond, as will be described in a follow-up study, and therefore were omitted here. ^b Q_{all} represents the Q -factor obtained using all available carbon RCSA data. ^c Q_{sp^3} represents the Q -factor obtained using only sp³ carbon RCSA data. ^d The angle θ represents the intertensor angle between the alignment tensor determined using all carbon RCSA data and that determined using only sp³ carbon RCSA data.

a Q_{all} of 0.029 in 15.5% PBLG is a significant improvement over a Q_{all} of 0.049 in the stretched PMMA gel for strychnine. Similarly, RCSAs from the rigid part of retrorsine produced a vastly improved Q_{all} of 0.033 in 15.1% PBLG, over a Q_{all} of 0.104 in stretched PMMA.

This strong alignment is particularly beneficial for compounds constituted primarily of sp³ carbons. We demonstrate this point by omitting all RCSA data from sp² carbons while using only those data from sp³ carbons to calculate the Q -factors (Q_{sp^3}) and alignment tensor parameters. As shown in Table 1, the improvement in Q_{sp^3} followed the trend of Q_{all} for data in PBLG vs. PMMA gels, with Q_{sp^3} from the former being lower than 0.1 in most cases. To further evaluate the reliability of using only sp³ carbon RCSAs for structural analysis, we compared the agreement of the alignment tensor parameters determined with all of the data and those determined with only sp³ data through calculations of their intertensor angles (θ).²⁹ As shown in Table 1, the intertensor angles are substantially smaller in PBLG solutions than in PMMA gels, with θ well below 10° in most cases, confirming that the reliability of using only sp³ carbon data significantly improved. One exception was observed for retrorsine in 12.6% PBLG, in which θ was slightly higher than in the PMMA gel, presumably due to inaccuracy of the alignment tensor determination with only a small number of sp³ carbons. From Table 1, it is also

apparent that the Q_{sp^3} -factor under a larger alignment for strychnine at the high end of PBLG concentration did not improve, but instead slightly degraded, whereas the Q_{all} -factor barely changes. This divergence between Q_{all} and Q_{sp^3} is possibly due to smaller sp³ carbon RCSA values being more susceptible to imperfections in our $\Delta\Delta\delta_{\text{iso}}$ elimination method at high PBLG concentrations where surface interference is expected to be more pronounced. Based on this observation, for a sp³ carbon-rich compound, a PBLG concentration of about 15% is advisable based on available data.

In a recent study, we successfully differentiated 13 possible diastereomers of strychnine using only RCSA data collected in a PMMA gel.⁸ Here we conducted the same analysis with RCSAs collected in 15.5% PBLG as a representative example (Fig. 3a), and compared the performance of PBLG vs. PMMA gel data for this differentiation (Fig. 3b). The stereochemical definition follows that in our previous work. The correct diastereomer *RSSRRS* with a Q -factor of 0.029, shown on the left, is clearly distinguished from all other diastereomers that exhibited much larger Q -factors (Fig. 3a). The confidence level of differentiation can be quantitatively assessed by the ratio: Q/Q_{best} , where Q is the Q -factor of an isomer being evaluated, and Q_{best} represents the lowest Q -factor in the pool,³⁰ which, in this example, is Q_{RSSRRS} . The larger this Q -factor ratio, the higher the level of confidence for the differentiation. Fig. 3b displays the Q -factor ratios of all 13 diastereomers over Q_{RSSRRS} , for both PBLG data (blue) and PMMA gel data (red). Clearly, the confidence level is about twice as good with PBLG data for those less discriminated isomers, such as *RSRRRS* and *SSRSR*, and overall much better, in comparison to the results obtained with PMMA gels. We should mention that the data quality from the stretched PMMA gel, with a Q_{RSSRRS} of 0.049, was actually quite good. However, that result is superseded by the enhanced quality afforded by the measurement in PBLG.

Next, as a more challenging example, we used RCSA data to differentiate caulamidine A (Fig. 1c) from its C26-inverted diastereomer. In a recent collaboration, we reported the structure of caulamidine A using NMR data including RDC and RCSA collected in a PHEMA gel.¹¹ Although a combination of RDC and RCSA clearly favoured the proposed structure over other constitutional and stereoisomers, the weak alignment of caulamidine A in PHEMA clearly left room for further improvement. Indeed, alignment with 13.1% PBLG generated a large RCSA range of –40 to

Fig. 3 Stereochemical differentiation of strychnine by RCSA data. (a) Differentiation by Q -factors using all carbon RCSA data collected in 15.5% PBLG (w/v). (b) Comparison of confidence levels of differentiation between 15.5% PBLG data in (a) (blue) and stretched PMMA gel data in ref. 2c (red). (c) Differentiation by Q -factors using sp³ carbon RCSA data collected in 15.5% PBLG. (d) Comparison of confidence levels of differentiation between using all carbon RCSAs (Q_{all}) and using only sp³ carbon RCSAs (Q_{sp^3}) with data collected in 15.5% PBLG.