



Cite this: *Polym. Chem.*, 2018, 9, 3959


## Correction: An injectable hyaluronic acid/PEG hydrogel for cartilage tissue engineering formed by integrating enzymatic crosslinking and Diels–Alder “click chemistry”

Feng Yu,<sup>a,b</sup> Xiaodong Cao,<sup>\*a,b,c</sup> Yuli Li,<sup>a,b,c</sup> Lei Zeng,<sup>a,c</sup> Bo Yuan<sup>a,b</sup> and Xiaofeng Chen<sup>\*a,b,c</sup>

DOI: 10.1039/c8py90106f  
rsc.li/polymers

Correction for ‘An injectable hyaluronic acid/PEG hydrogel for cartilage tissue engineering formed by integrating enzymatic crosslinking and Diels–Alder “click chemistry” by Feng Yu et al., *Polym. Chem.*, 2014, 5, 1082–1090.

The authors regret that the incorrect images were used in Fig. 8A, B and C to represent three different culture time points. The correct version of Fig. 8 is shown below.




**Fig. 8** Live–dead and CCK-8 assay of ATDC-5 cells encapsulated in HA/PEG DS1 hydrogel (living cells appearing green, 4 magnifications). (A–C) The live–dead assay of ATDC-5 cells encapsulated in hydrogels after 1, 3, 7 days culture with 40% image brightness increasing. (D) A CCK-8 cytotoxicity assay was performed to evaluate the cells proliferation after 1, 3, 5 and 7 days culture.

<sup>a</sup>School of Materials Science and Engineering, South China University of Technology, Guangzhou, 510641, PR China. E-mail: caoxd@scut.edu.cn; Fax: +86-20-87111752; Tel: +86-20-22236066

<sup>b</sup>National Engineering Research Centre for Tissue Restoration and Reconstruction, Guangzhou, 510006, PR China. E-mail: chenxf@scut.edu.cn; Fax: +86-20-22236083; Tel: +86-20-22236283

<sup>c</sup>Guangdong Province Key Laboratory of Biomedical Engineering, South China University of Technology, Guangzhou 510640, PR China





**Fig. S1** Live–dead assay of encapsulated ATDC-5 cells in HA/PEG DS1 hydrogel (living cells appearing green, 4 magnifications). (A–I) The live–dead assay of ATDC-5 cells encapsulated in hydrogels after 1, 3, 7 days culture with different brightness. A–C are the original images. D–F are the images with 20% brightness increasing. G–I are the images with 40% brightness increasing.

**Table S1** CCK-8 assay results after different time points

OD value	1 day	3 days	5 days	7 days
Parallel group 1	0.68284	0.981505	1.39048	1.848
Parallel group 2	0.723215	1.02393	1.431505	1.94505
Parallel group 3	0.731945	0.921341	1.429301	1.79301
Parallel group 4	0.703852	0.843448	1.321341	1.96406
Average value	0.710463	0.942556	1.393157	1.88753

In order to further prove the authenticity of the experimental data, additional supplementary information is also provided here.

Fig. S1(A–C) are the original images generated using fluorescence microscopy. Because of the high cell culture density encapsulated in the hydrogel ( $3 \times 10^6$  cells per ml), the phenomenon of cell proliferation after 1 d, 3 d and 7 d culture cannot be distinguished and observed especially in low brightness. Therefore, the image brightness was increased to 20% and 40% to make sure that more cells (marked with green fluorescence) could appear in the image and so that the cell proliferation after different culture days could be observed by the naked eye (Fig. S1(D–I)).

In addition, a CCK-8 assay is an efficient and quantitative way to prove the cell proliferation. Here, the detailed data are also provided in Table S1 to further supplement the data shown in Fig. 8D.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

