PAPER

View Article Online
View Journal | View Issue Cite this: *RSC Adv.*, 2018, 8, 3899Received 17th December 2017
Accepted 15th January 2018

DOI: 10.1039/c7ra13391j

rsc.li/rsc-advances

Synthesis of C14–C21 acid fragments of cytochalasin Z₈ via *anti*-selective aldol condensation and *B*-alkyl Suzuki–Miyaura cross-coupling†

Weiwei Han ^{*ab}

An efficient synthesis of the C14–C21 acid fragment of cytochalasin Z₈ was accomplished in 10 steps with 14% overall yield. Boron-mediated *anti*-selective aldol condensation and Pd(OAc)₂–Aphos–Y-catalysed *B*-alkyl Suzuki–Miyaura cross-coupling were employed to construct the requisite C17 and C18 stereogenic centres and alkene subunit.

Cytochalasins are secondary fungal metabolites with a wide range of biological activities that target cytoskeletal processes.¹ Cytochalasins Z₇–Z₉ (1–3, Chart 1) were isolated from the marine-derived fungus *Spicaria elegans*, and their structures and absolute configurations were established by Zhu *et al.*² Cytochalasin Z₈ (2, Chart 1) is structurally related to cytochalasins Z₇ and Z₉ and features highly substituted hydroisoindol-1-one fused with a 12-membered macrolactone ring at the C-8 and C-9 positions. Cytochalasin Z₈ has been reported to exert cytotoxicity against P388 and A-549 cell lines with IC₅₀ values of 56 and 21 μM, respectively, and therefore has significant potential in cell biology and medicine. A number of laboratories have worked towards total synthesis of the cytochalasin family and developed linear³ or convergent⁴ strategies for their total synthesis. Total synthesis of cytochalasin congeners was accomplished by the laboratories of Stork,^{3a,4a} Thomas,^{3b,3c,3e,3f}

Trost,^{4d} Vedejs (zygospurin E),^{4b,4c,4e} Myers,⁵ Liu and Tang (periconiasins A–E)⁶ and Nay (periconiasin G).⁷ To the best of our knowledge, total synthesis of cytochalasin with a 12-membered macrocyclic ring has not been reported. The intriguing molecular architecture and potent biological activity of cytochalasin Z₈ prompted us to pursue its total synthesis and render it to be readily available for biological investigations.

The retrosynthetic strategy is depicted in Scheme 1. Intramolecular ring-closing metathesis (RCM) strategy⁸ which is a promising tool for constructing macrolactone is often used for synthesising macrolides.⁹ We envisioned an RCM reaction at C13 and C14 positions and an esterification for assembling a 12-membered macrolactone. Thus, acid fragment 4 was required for the total synthesis of 2. Our strategy was flexible and it allowed rapid access to structural analogues. In this study, we report the synthesis of C14–C21 acid fragment 4 via a highly *anti*-selective aldol condensation¹⁰ of aldehyde 6 with Abiko's chiral norephedrine-derived propionate (1*R*,2*S*)-7 (ref. 11) and *B*-alkyl Suzuki–Miyaura cross-coupling¹² of chiral alkyl iodide 5 with (Z)-1-bromoprop-1-ene.

Our first task was to construct C16–C18 *syn-anti* stereotriad.¹³ The aldehyde functionality in 6 was expected to undergo an *anti*-selective aldol reaction with the (*E*)-boron enolate generated from Abiko's chiral propionate 7 for installing C17–C18 *anti* stereochemistry according to our synthetic strategy in Scheme 1. We initially prepared crude aldehyde 6 from commercially available (*S*)-methyl 3-hydroxy-2-methyl propionate (Roche ester)¹⁴ by tosylation and partial ester reduction¹⁵ (Scheme 2). The unstable crude aldehyde 6, without column chromatographic purification, was immediately used with the (*E*)-boron enolate derived from 7 for *anti*-selective aldol reaction to secure the *syn/anti* stereotriad in 8. The key intermediate 8 was prepared in high diastereoselectivity of 98 : 2 (determined by proton nuclear magnetic resonance spectroscopy) and in the desired absolute configuration as predicted by the chiral

Chart 1 Structures of cytochalasin Z₇–Z₉.

^aCollege of Chemistry and Chemical Engineering, Xi'an Shiyou University, Xi'an, 710065, P. R. China. E-mail: vivien2014@xsyu.edu.cn; Tel: +86-29-88382703

^bLaboratory of Asymmetric Catalysis and Synthesis, Department of Chemistry, Zhejiang University, Hangzhou 310027, P. R. China

† Electronic supplementary information (ESI) available: Experimental details and scanned copies of ¹H and ¹³C NMR spectra for new compounds. See DOI: 10.1039/c7ra13391j

Scheme 1 Retrosynthetic bond disconnections of cytochalasin **2** yielding C14–C21 acid fragment **4** and hydroisindol-1-one fragment.

Scheme 2 Synthesis of alkyl iodide **6**.

auxiliary in **7**. The influence of the stereogenic centre of aldehyde **6** on the stereochemical course of the aldol reaction was not observed. The hydroxyl group in **8** was then protected as TES ether **9** (TESOTf, 2, 6-lutidine, 98% yield). Iodide replacement of the tosylate group in **9** with LiI–THF furnished alkyl iodide **5** in 95% yield (Scheme 2).

The cross-coupling reaction of chiral alkyl iodide **5** with (*Z*)-1-bromoprop-1-ene was performed under the established conditions¹⁶ for the '9-MeO-9-BBN variant' of the *B*-alkyl Suzuki–Miyaura cross-coupling reaction.^{12f,17} Alkyl iodide **5** was treated with *t*-BuLi in the presence of 9-MeO-9-BBN in Et₂O–THF to form the corresponding borinate species which was subjected to Pd(OAc)₂–Aphos-Y-catalysed^{16,18} cross-coupling reaction with (*Z*)-

1-bromoprop-1-ene in the presence of K₃PO₄·3H₂O as the base in THF–H₂O at room temperature to furnish **11** in 15% yield along with cyclopentanol **10** and deiodinated byproduct **12** (entry 1, Table 1). We speculated that cyclopentanol byproduct **10** would be formed in the following pathway. Treatment of **5** with *t*-BuLi formed alkylolithium which underwent an intramolecular cycloaddition to form cyclopentanone; cyclopentanol **10** was formed by the addition of *t*-BuLi (Scheme 3). These results suggested that the formation of **10** could be suppressed by controlling reaction temperature. The first step reaction was maintained under low temperatures for a long time before warming up. After adding *t*-BuLi and THF, the reaction temperature was sequentially kept at –78 °C for 30 min, at –40 °C for 30 min, at –20 °C for 30 min and

Table 1 Results of cross coupling of chiral alkyl iodide **5** with (*Z*)-1-bromoprop-1-ene

Entry	Conditions	Conditions	Yield ^a (%)
	Step 1	Step 2	
1	2.8 eq. <i>t</i> -BuLi, 3.0 eq. 9-MeO-9-BBN, Et ₂ O/THF, –78 °C then r.t. for 2 h	5.0 mol% Pd(OAc) ₂ , 7.5 mol% Aphos-Y, 3.0 eq. K ₃ PO ₄ ·3H ₂ O, 18.0 eq. H ₂ O, THF, r.t. (14 h)	15; (21; 15)
2	3.8 eq. <i>t</i> -BuLi, 5.0 eq. 9-MeO-9-BBN, Et ₂ O/THF, –78 °C (30 min), –40 °C (30 min), –20 °C (90 min), r.t. (2 h)	5.0 mol% Pd(OAc) ₂ , 7.5 mol% Aphos-Y, 3.0 eq. K ₃ PO ₄ ·3H ₂ O, 18.0 eq. H ₂ O, THF, r.t. (14 h)	32; (19; 32)
3	4.0 eq. <i>t</i> -BuLi, 4.5 eq. 9-MeO-9-BBN, Et ₂ O/THF, –78 °C (30 min), –40 °C (30 min), –20 °C (90 min), r.t. (2 h)	10 mol% Pd(OAc) ₂ , 15 mol% Aphos-Y, 3.0 eq. K ₃ PO ₄ ·3H ₂ O, 18.0 eq. H ₂ O, THF, r.t. (12 h)	40; (17; 10)

^a Isolated yield of product **11**. Data in the parentheses are the isolated yields of cyclopentanol **10** and deiodinated byproduct **12**, respectively.

Scheme 3 Cross-coupling of chiral alkyl iodide 5 with (Z)-1-bromoprop-1-ene.

Scheme 4 Synthesis of C14–C21 acid fragment 4.

at room temperature for 2 h. The newly formed borinate species was subjected to coupling reaction with (Z)-1-bromoprop-1-ene. The yield was improved to 40% (entry 3, Table 1), and deiodinated byproduct 12 was inhibited to a large extent but could not be eliminated (in 10% yield). This condition might be associated with the steric hindrance imposed by the bulky TES and Abiko's chiral ester moieties of 5.

The completion of the total synthesis of acid fragment 4 is illustrated in Scheme 4. Reduction of 11 with DIBAL-H provided the resultant primary alcohol 13 in 75% isolated yield. Dess–Martin periodinane oxidation¹⁹ in the presence of NaHCO₃ converted 13 into the corresponding aldehyde 14. Aldehyde 14 was subjected to Wittig olefination with the stabilised ylide, Ph₃P=CHCO₂Me, in toluene at 60 °C to produce α,β -unsaturated ester 15 with exclusive *E* configuration for the newly formed disubstituted double bond. The hydrolysis of methyl ester 15 with LiOH in THF/H₂O at room temperature furnished the target C14–C21 acid fragment 4 in 72% overall yield for the three steps.

Conclusions

We developed a concise synthesis of the C14–C21 acid fragment 4 of cytochalasin Z₈. The *anti*-selective aldol reaction of 6 with the (*E*)-boron enolate derived from Abiko's chiral propionate 7

achieved the desired C16–C18 *syn/anti* stereotriad in high diastereoselectivity presumably attained *via* a reagent control process. (Z)-Alkene functionality was introduced by the Pd(OAc)₂–Aphos-Y-catalysed C(sp²)–C(sp³) bond formation reaction of chiral alkyl iodide 5 with (Z)-1-bromoprop-1-ene. The target acid fragment 4 could be prepared from chiral (*S*)-Roche ester by a 10-step sequence in an overall yield of 14%. The strategy is concise and flexible to produce additional analogues of cytochalasin Z₈ in enantiomerically pure form. Efforts to achieve this goal are ongoing in our laboratory.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We are grateful for financial support from the National Natural Science Foundation of China (Project no. 21172191).

Notes and references

- (a) M. Binder and C. Tamm, *Angew. Chem., Int. Ed.*, 1973, **12**, 370; (b) F. Wang, H. Wei, T. Zhu, D. Li, Z. Lin and Q. Gu, *Chem. Biodiversity*, 2011, **8**, 887; (c) J. Wang, Z. Wang, Z. Ju, J. Wan, S. Liao, X. Lin, T. Zhang, X. Zhou, H. Chen and

- Z. Tu, *Planta Med.*, 2015, **81**, 160; (d) J. Xu, *RSC Adv.*, 2015, **5**, 841.
- 2 R. Liu, Q. Gu, W. Zhu, C. Cui, G. Fan, Y. Fang, T. Zhu and H. Liu, *J. Nat. Prod.*, 2006, **69**, 871.
- 3 (a) G. Stork and E. Nakamura, *J. Am. Chem. Soc.*, 1983, **105**, 5510; (b) H. Dyke, R. Sauter, P. Steel and E. J. Thomas, *J. Chem. Soc., Chem. Commun.*, 1986, 1447; (c) E. J. Thomas and J. W. F. Whitehead, *J. Chem. Soc., Chem. Commun.*, 1986, 727; (d) E. J. Thomas and J. W. F. Whitehead, *J. Chem. Soc., Perkin Trans. 1*, 1989, 499; (e) E. Merifield and E. J. Thomas, *J. Chem. Soc., Chem. Commun.*, 1990, 464; (f) E. Merifield and E. J. Thomas, *J. Chem. Soc., Perkin Trans. 1*, 1999, 3269.
- 4 (a) G. Stork, Y. Nakahara, Y. Nakahara and W. J. Greenlee, *J. Am. Chem. Soc.*, 1978, **100**, 7775; (b) E. Vedejs and J. G. Reid, *J. Am. Chem. Soc.*, 1984, **106**, 4617; (c) E. Vedejs, J. D. Rodgers and S. J. Wittenberger, *J. Am. Chem. Soc.*, 1988, **110**, 4822; (d) B. M. Trost, M. Ohmori, S. A. Boyd, H. Okawara and S. J. Brickner, *J. Am. Chem. Soc.*, 1989, **111**, 8281; (e) E. Vedejs, J. G. Reid, J. D. Rodgers and S. J. Wittenberger, *J. Am. Chem. Soc.*, 1990, **112**, 4351.
- 5 A. M. Haidle and A. G. Myers, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 12048.
- 6 C. Tian, X. Lei, Y. Wang, Z. Dong, G. Liu and Y. Tang, *Angew. Chem., Int. Ed.*, 2016, **55**, 6992.
- 7 M. Zaghouani, C. Kunz, L. Guédon, F. Blanchard and B. Nay, *Chem.-Eur. J.*, 2016, **22**, 15257.
- 8 (a) R. H. Grubbs and S. Chang, *Tetrahedron*, 1998, **54**, 4413; (b) A. Fürstner, *Angew. Chem., Int. Ed.*, 2000, **39**, 3012; (c) T. M. Trnka and R. H. Grubbs, *Acc. Chem. Res.*, 2001, **34**, 18; (d) R. R. Schrock and A. H. Hoveyda, *Angew. Chem., Int. Ed.*, 2003, **42**, 4592; (e) S. P. Nolan and H. Clavier, *Chem. Soc. Rev.*, 2010, **39**, 3305; (f) J. W. Herndon, *Coord. Chem. Rev.*, 2010, **254**, 103; (g) G. C. Vougioukalakis and R. H. Grubbs, *Chem. Rev.*, 2010, **110**, 1746; (h) A. H. Hoveyda, S. J. Malcolmson, S. J. Meek and A. R. Zhugralin, *Angew. Chem., Int. Ed.*, 2010, **49**, 34; (i) A. Fürstner, *Chem. Commun.*, 2011, **47**, 6505; (j) S. Kotha and M. K. Dipak, *Tetrahedron*, 2012, **68**, 397; (k) A. Fürstner, *Science*, 2013, **341**, 1229713.
- 9 B. Thirupathi and D. K. Mohapatra, *RSC Adv.*, 2014, **4**, 8027.
- 10 (a) B. Schetter and R. Mahrwald, *Angew. Chem., Int. Ed.*, 2006, **45**, 7506; (b) L. M. Geary and P. G. Hultin, *Tetrahedron: Asymmetry*, 2009, **20**, 131.
- 11 (a) A. Abiko, J.-F. Liu and S. Masamune, *J. Am. Chem. Soc.*, 1997, **119**, 2586; (b) T. Inoue, J.-F. Liu, D. C. Buske and A. Abiko, *J. Org. Chem.*, 2002, **67**, 5250; (c) A. Abiko, *Acc. Chem. Res.*, 2004, **37**, 387.
- 12 (a) S. R. Chemler, D. Trauner and S. J. Danishefsky, *Angew. Chem., Int. Ed.*, 2001, **40**, 4544; (b) K. C. Nicolaou, P. G. Bulger and D. Sarlah, *Angew. Chem., Int. Ed.*, 2005, **44**, 4442; (c) S. Kotha and K. Mandal, *Chem.-Asian J.*, 2009, **4**, 354; (d) A. Suzuki, *Angew. Chem., Int. Ed.*, 2011, **50**, 6722; (e) R. Jana, T. P. Pathak and M. S. Sigman, *Chem. Rev.*, 2011, **111**, 1417; (f) G. Seidel and A. Fürstner, *Chem. Commun.*, 2012, **48**, 2055; (g) M. M. Heravi and E. Hashemi, *Tetrahedron*, 2012, **68**, 9145.
- 13 R. W. Hoffmann, *Angew. Chem., Int. Ed. Engl.*, 1987, **26**, 489.
- 14 (a) C. Aïssa, R. Riveiros, J. Ragot and A. Fürstner, *J. Am. Chem. Soc.*, 2003, **125**, 15512; (b) D. A. Kummer, J. B. Brenneman and S. F. Martin, *Org. Lett.*, 2005, **7**, 4621.
- 15 H. Li, J. Wu, J. Luo and W.-M. Dai, *Chem.-Eur. J.*, 2010, **16**, 11530.
- 16 (a) N. Ye and W. M. Dai, *Eur. J. Org. Chem.*, 2013, **2013**, 831; (b) Y. Wu, Y. Lai and W.-M. Dai, *ChemistrySelect*, 2016, **1**, 1022.
- 17 (a) A. Fürstner and G. Seidel, *Tetrahedron*, 1995, **51**, 11165; (b) J. A. Soderquist, K. Matos, A. Rane and J. Ramos, *Tetrahedron Lett.*, 1995, **36**, 2401.
- 18 J. Jin, Y. Chen, Y. Li, J. Wu and W.-M. Dai, *Org. Lett.*, 2007, **9**, 2585.
- 19 (a) D. B. Dess and J. C. Martin, *J. Org. Chem.*, 1983, **48**, 4155; (b) S. D. Meyer and S. L. Schreiber, *J. Org. Chem.*, 1994, **59**, 7549.